


3.3 Differentiation Rules



Colorado National Monument

Photo by Vickie Kelly, 2003

Greg Kelly, Hanford High School, Richland, Washington

A quick review


If f has a derivative at $x = a$, then f is continuous at $x = a$.

Since a function must be continuous to have a derivative, if it has a derivative then it is continuous.



Intermediate Value Theorem for Derivatives

If a and b are any two points in an interval on which f is differentiable, then f' takes on every value between $f'(a)$ and $f'(b)$.



Between a and b , f' must take on every value between $\frac{1}{2}$ and 3 .

If the derivative of a function is its slope, then for a constant function, the derivative must be zero.

$$\frac{d}{dx}(c) = 0$$

example: $y = 3$
 $y' = 0$

The derivative of a constant is zero.



We saw that if $y = x^2$, $y' = 2x$.

This is part of a pattern.

$$\frac{d}{dx}(x^n) = nx^{n-1}$$



power rule

examples:

$$f(x) = x^4$$

$$y = x^8$$

$$f'(x) = 4x^3$$

$$y' = 8x^7$$



constant multiple rule:

$$\frac{d}{dx}(cu) = c \frac{du}{dx}$$

examples:

$$\frac{d}{dx} cx^n = cnx^{n-1}$$

$$\frac{d}{dx} 7x^5 = 7 \cdot 5x^4 = 35x^4$$



constant multiple rule:

$$\frac{d}{dx}(cu) = c \frac{du}{dx}$$

sum and difference rules:

$$\frac{d}{dx}(u - v) = \frac{du}{dx} - \frac{dv}{dx}$$

$$y = x^4 + 12x$$

$$y' = 4x^3 + 12$$

$$y = x^4 - 2x^2 + 2$$

$$\frac{dy}{dx} = 4x^3 - 4x$$



Find the derivative of each function below

$$y = x^3 + x + 2$$

$$y = 2x^{3/2} + x^{-2}$$

$$y' = 3x^2 + 1$$

$$\frac{dy}{dx} = 3x^{1/2} - 2x^{-3}$$

Example:

Find the horizontal tangents of: $y = x^4 - 2x^2 + 2$

$$\frac{dy}{dx} = 4x^3 - 4x$$

Horizontal tangents occur when slope = zero.

$$4x^3 - 4x = 0$$

$$x^3 - x = 0$$

$$x(x^2 - 1) = 0$$

$$x(x+1)(x-1) = 0$$


$$x = 0, -1, 1$$


Plugging the x values into the original equation, we get:

$$y = 2, \quad y = 1, \quad y = 1$$


(The function is even, so we only get two horizontal tangents.)








$$y = x^4 - 2x^2 + 2$$



$$y = x^4 - 2x^2 + 2$$


$$y = 2$$




$$y = x^4 - 2x^2 + 2$$

$$y = 2$$

$$y = 1$$



$$y = x^4 - 2x^2 + 2$$



$$y = x^4 - 2x^2 + 2$$

$$\frac{dy}{dx} = 4x^3 - 4x$$

First derivative
(slope) is zero at:

$$x = 0, -1, 1$$



$$\frac{d}{dx} \left[(x^2 + 3)(2x^3 + 5x) \right]$$

$$\frac{d}{dx} (2x^5 + 5x^3 + 6x^3 + 15x)$$

$$\frac{d}{dx} (2x^5 + 11x^3 + 15x) = 10x^4 + 33x^2 + 15$$



product rule:

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

Notice that this is not just the product of two derivatives.

This is sometimes memorized as: $d(uv) = u \, dv + v \, du$

$$\frac{d}{dx}[(x^2 + 3)(2x^3 + 5x)] = (x^2 + 3)(6x^2 + 5) + (2x^3 + 5x)(2x)$$

$$6x^4 + 5x^2 + 18x^2 + 15 + 4x^4 + 10x^2$$

$$10x^4 + 33x^2 + 15$$

(1st)(Derivative of 2nd) + (2nd)(Derivative of 1st)

(Lefty(D-righty) + (Righty)(D-lefty))



quotient rule:

$$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

or

$$d \left(\frac{u}{v} \right) = \frac{v \, du - u \, dv}{v^2}$$

$$\frac{d}{dx} \frac{2x^3 + 5x}{x^2 + 3} = \frac{(x^2 + 3)(6x^2 + 5) - (2x^3 + 5x)(2x)}{(x^2 + 3)^2}$$

(Bottom)(Derivative of Top) – (Top)(Derivative of Bottom) all divided by the bottom squared

(Low)(D-high) – (High)(D-low) all divided by low squared



Higher Order Derivatives:

$y' = \frac{dy}{dx}$ is the first derivative of y with respect to x .

$y'' = \frac{dy'}{dx} = \frac{d}{dx} \frac{dy}{dx} = \frac{d^2 y}{dx^2}$ is the second derivative.
(y double prime)

$y''' = \frac{dy''}{dx}$ is the third derivative.

$y^{(4)} = \frac{d}{dx} y'''$ is the fourth derivative.

We will learn
later what these
higher order
derivatives are
used for.

Product Rule

(1st)(Derivative of 2nd) + (2nd)(Derivative of 1st)

(Lefty)(D-righty) + (Righty)(D-lefty)

Quotient Rule

(Bottom)(Derivative of Top) – (Top)(Derivative of Bottom) all
divided by the bottom squared

(Low)(D-high) – (High)(D-low) all
divided by low squared

24. Suppose u and v are differentiable functions at $x = 2$

$$u(2) = 3, \quad u'(2) = -4, \quad v(2) = 1, \quad v'(2) = 2$$

Find

$$a) \frac{d}{dx}(uv) \qquad b) \frac{d}{dx}\left(\frac{u}{v}\right) \qquad c) \frac{d}{dx}\left(\frac{v}{u}\right)$$

$$d) \frac{d}{dx}(3u - 2v + 2uv)$$

