Jasper City Schools 8th Grade Pre-Algebra Pacing Guide 08.14.2018

- Thoughtful and effective <u>planning</u> throughout the school year is crucial for student mastery of standards.
- Once a standard is introduced, it is understood that the standard is continuously taught and/or reviewed throughout the <u>entire</u> school year
 ONGOING standards listed after pacing guide

Fourth Nine Weeks First Nine Weeks **Second Nine Weeks Third Nine Weeks** 3.) Know and apply the properties of **1.**) Know that numbers that are not 1.) Know the formulas for the **16.)** Verify experimentally the integer exponents to generate properties of rotations, reflections, and rational are called irrational. volumes of cones, cylinders, and equivalent numerical expressions. spheres, and use them to solve real-Understand informally that every translations: [8-G1] number has a decimal expansion; for [8-EE1] world and mathematical problems. a. Lines are taken to lines, and line rational numbers show that the [8-G9] segments are taken to line segments of the same length. [8-G1a] decimal expansion repeats 5.) Use numbers expressed in the eventually, and convert a decimal form of a single digit times an integer 2.) Use rational approximations of b. Angles are taken to angles of expansion which repeats eventually power of 10 to estimate very large irrational numbers to compare the the same measure. [8-G1b] into a rational number. [8-NS1] or very small quantities, and to size of irrational numbers, locate c. Parallel lines are taken to express how many times as much them approximately on a number line parallel lines. [8-G1c] one is than the other. [8-EE3] diagram, and estimate the value of 9.) Solve linear equations in one expressions (e.g., π 2). [8-NS2] 17.) Understand that a two dimensional variable. [8-EE7] **6.)** Perform operations with numbers figure is congruent to another if the a. Give examples of linear equations expressed in scientific notation, 10.) Analyze and solve pairs of second can be obtained from the first in one variable with one solution. including problems where both simultaneous linear equations. by a sequence of rotations, reflections, infinitely many solutions, or no decimal and scientific notation are [8-EE8] and translations; given two congruent solutions. Show which of these a. Understand that solutions to a used. Use scientific notation and figures, describe a sequence that possibilities is the case by exhibits the congruence between them. successively transforming the given choose units of appropriate size for system of two linear equations in two measurements of very large or very variables correspond to points of [8-G2] equation into simpler forms until an small quantities (e.g., use millimeters intersections of their graphs because equivalent equation of the form x = a, per year for seafloor spreading). points of intersection satisfy both 18.) Describe the effect of dilations, a = a, or a = b results (where a and b equations simultaneously. [8-EE8a] Interpret scientific notation that has translations, rotations, and reflections are different numbers). [8-EE7a] been generated by technology. [8b. Solve systems of two linear on two-dimensional figures using b. Solve linear equations with rational number coefficients. EE41 equations in two variables algebraically, coordinates. [8-G3] and estimate solutions by graphing the including equations whose solutions equations. Solve simple cases by require expanding expressions, inspection. [8-EE8b] using the distributive property 11.) Understand that a function is a 7.) Graph proportional relationships, **19.)** Understand that a two dimensional and collecting like terms. [8-EE7b] interpreting the unit rate as the slope rule that assigns to each input exactly figure is similar to another if the second

- **4.)** Use square root and cube root symbols to represent solutions to equations of the form $x_2 = p$ and $x_3 = p$, where p is a positive rational number. Evaluate square roots of small perfect squares and cube roots of small perfect cubes. Know that $\sqrt{2}$ is irrational. [8-EE2]
- 25.) Construct and interpret scatter plots for bivariate measurement data to investigate patterns of association between two quantities. Describe patterns such as clustering, outliers, positive or negative association, linear association, and nonlinear association. [8-SP1]
- **26.)** Know that straight lines are widely used to model relationships between two quantitative variables. For scatter plots that suggest a linear association, informally fit a straight line, and informally assess the model fit by judging the closeness of the data points to the line. [8-SP2]

- of the graph. Compare two different proportional relationships represented in different ways. [8-EE5]
- **8.)** Use similar triangles to explain why the slope m is the same between any two distinct points on a non-vertical line in the coordinate plane; derive the equation y = mx for a line through the origin and the equation y = mx + b for a line intercepting the vertical axis at b. [8-EE6]
- **20.)** Use informal arguments to establish facts about the angle sum and exterior angle of triangles, about the angles created when parallel lines are cut by a transversal, and the angle-angle criterion for similarity of triangles. [8-G5]

- one output. The graph of a function is the set of ordered pairs consisting of an input and the corresponding output. (Function notation is not required in Grade 8.) [8-F1]
- **12.)** Compare properties of two functions, each represented in a different way (algebraically, graphically, numerically in tables, or by verbal descriptions). [8-F2]
- **13.)** Interpret the equation y = mx + b as defining a linear function whose graph is a straight line; give examples of functions that are not linear. [8-F3]
- **14.)** Construct a function to model a linear relationship between two quantities. Determine the rate of change and initial value of the function from a description of a relationship or from two (*x*, *y*) values, including reading these from a table or from a graph. Interpret the rate of change and initial value of linear function in terms of the situation it models and in terms of its graph or a table of values. [8-F4]
- **15.)** Describe qualitatively the functional relationship between two quantities by analyzing a graph (e.g., where the function is increasing or decreasing, linear or nonlinear). Sketch a graph that exhibits the qualitative features of a function that has been described verbally. [8-F5]
- **21.)** Explain a proof of the Pythagorean Theorem and its converse. [8-G6]
- **22.)** Apply the Pythagorean Theorem to determine unknown side lengths in right triangles in real-world and mathematical problems in two and

- can be obtained from the first by a sequence of rotations, reflections, translations, and dilations; given two similar two-dimensional figures, describe a sequence that exhibits the similarity between them. [8-G4]
- **24.)** Know the formulas for the volumes of cones, cylinders, and spheres, and use them to solve real-world and mathematical problems. [8-G9]
- 28.) Understand that patterns of association can also be seen in bivariate categorical data by displaying frequencies and relative frequencies in a two-way table. Construct and interpret a two-way table summarizing data on two categorical variables collected from the same subjects. Use relative frequencies calculated for rows or columns to describe possible association between the two variables. [8-SP4]

three dimensions. [8-G7]	
23.) Apply the Pythagorean Theorem to find the distance between two points in a coordinate system. [8-G8]	
27.) Use the equation of a linear model to solve problems in the context of bivariate measurement data, interpreting the slope and intercept. [8-SP3]	

Power Standards

** power standards are indicated with an asterisk **

These standards are those that are essential for student grade-level success.