


1

## **LESSON 10-4**



a. 
$$\sqrt{x} - 5 = 4$$
  
 $\sqrt{x} - 5 = 4$   
 $\sqrt{x} = 9$  Isolate the radical on the left side of the equation.  
 $(\sqrt{x})^2 = 9^2$  Square each side.  
 $x = 81$ 

Check: 
$$\sqrt{x} - 5 = 4$$
  
 $\sqrt{81} - 5 \stackrel{?}{=} 4$  Substitute 81 for *x*.  
9 - 5  $\stackrel{?}{=} 4$   
4 = 4 ✓





PEARSON

Prentice Hall



EXAMPLE (continued)

**b.** 
$$\sqrt{x-5} = 4$$
  
 $(\sqrt{x-5})^2 = 4^2$ Square each side

x - 5 = 16 Solve for x.

*x* = 21


Check: 
$$\sqrt{x-5} = 4$$
  
 $\sqrt{21-5} = 4$ Substitute 21 for x.  
 $\sqrt{16} = 4$   
 $4 = 4$ 







**Definition** I be a second of a car at the top of the loop.








#### LESSON 10-4





The loop on a roller coaster ride has a radius of 18 ft. Your car has a velocity of 120 ft/s at the top of the loop. How high is the hill of the loop you have just come down before going into the loop?



Solve  $v = 8\sqrt{h-2r}$  for h when v = 120 and r = 18.  $120 = 8\sqrt{h-2(18)}$  Substitute 120 for v and 18 for r.  $\frac{120}{8}$  Divide Gach Side by 8 to isolate the radical.  $15 = \sqrt{h-36}$ Simplify.  $(15)^2 = (\sqrt{h-36})^2$  Square both sides. 225 = h - 36 261 = hThe hill is 261 ft high.





## **Additional Examples**



3 EXAMPLE 
$$3x - 4 \neq 2x + 3.\sqrt{}$$

 $(\sqrt{3x-4})^2 = (\sqrt{2x+3})^2$ Square both sides.

3x - 4 = 2x + 3Simplify.

3x = 2x + 7Add 4 to each side.

x = 7Subtract 2x from each side.

Check: 
$$\sqrt{3x-4} = \sqrt{2x+3}$$
  
 $\sqrt{3(7)-4} \ge \sqrt{2(7)+3}$ Substitute 7 for x.  
 $\sqrt{17} = \sqrt{17}$ 

The solution is 7.





# **Solving Radical Equations**

EXAM

#### **LESSON 10-4**

### **Additional Examples**



ple : 
$$x + 12.\sqrt{}$$

$$(x)^{2} = (\sqrt{x+12})^{2}$$
$$x^{2} = x + 12$$

Square both sides.

 $x^2 - x - 12 = 0$ 

$$(x-4)(x+3) = 0$$

$$(x-4) = 0$$
 or  $(x + 3) = 0$   
 $x = 4$  or  $x = -3$ 

Simplify.

Solve the quadratic equation by factoring.

Use the Zero–Product Property. Solve for *x*.

**Check:** 
$$x = \sqrt{x + 12}$$
  
 $4 \stackrel{?}{=} \sqrt{4 + 12}$   $-3 \stackrel{?}{=} \sqrt{-3 + 12}$   
 $4 = 4 \stackrel{\checkmark}{=} 3$ 

The solution to the original equation is 4. The value –3 is an extraneous solution.





**Solving Radical Equations** 

## **LESSON 10-4**

## **Additional Examples**

5 EXAMPLE 
$$3x + 8 \neq \sqrt{2}$$
.

$$\sqrt{3x} = -6$$
Subtract 8 from each side.  

$$(\sqrt{3x})^2 = (-6)^2$$
Square both sides.  

$$3x = 36$$

$$x = 12$$
Check:  $\sqrt{3x} + 8 = 2$   
 $\sqrt{3(12)} + 8 \stackrel{?}{=} 2$ 
Substitute 12 for x.  
 $\sqrt{36} + 8 \stackrel{?}{=} 2$   
 $6 + 8 \neq 2$ 
 $x = 12$  does not solve the original equation

 $\sqrt{3x}$  + 8 = 2 has no solution.





PEARSON

Prentice Hall