

2.5

The Fundamental Theorem of Algebra



What You Should Learn

- Use the Fundamental Theorem of Algebra to determine the number of zeros of a polynomial function.
- Find all zeros of polynomial functions, including complex zeros.
- Find conjugate pairs of complex zeros.
- Find zeros of polynomials by factoring.



The Fundamental Theorem of Algebra



The Fundamental Theorem of Algebra

We know that an n th-degree polynomial can have at most n real zeros. In the complex number system, this statement can be improved. That is, **in the complex number system, every n th-degree polynomial function has precisely n zeros.**

This important result is derived from the **Fundamental Theorem of Algebra**, first proved by the German mathematician Carl Friedrich Gauss (1777–1855).

The Fundamental Theorem of Algebra

If $f(x)$ is a polynomial of degree n , where $n > 0$, then f has at least one zero in the complex number system.



The Fundamental Theorem of Algebra


Using the Fundamental Theorem of Algebra and the equivalence of zeros and factors, you obtain the **Linear Factorization Theorem**.

Linear Factorization Theorem

If $f(x)$ is a polynomial of degree n , where $n > 0$, then f has precisely n linear factors

$$f(x) = a_n(x - c_1)(x - c_2) \cdot \cdot \cdot (x - c_n)$$

where $c_1, c_2, \cdot \cdot \cdot, c_n$ are complex numbers.



Example 1 – *Zeros of Polynomial Functions*


a. The first-degree polynomial $f(x) = x - 2$ has exactly *one* zero: $x = 2$.

b. Counting multiplicity, the second-degree polynomial function

$$f(x) = x^2 - 6x + 9$$

$$= (x - 3)(x - 3)$$

has exactly *two* zeros: $x = 3$ and $x = 3$ (This is called a *repeated zero*.)



Example 1 – *Zeros of Polynomial Functions*

cont'd

c. The third-degree polynomial function

$$\begin{aligned}f(x) &= x^3 + 4x \\&= x(x^2 + 4) \\&= x(x - 2i)(x + 2i)\end{aligned}$$

has exactly *three* zeros: $x = 0$, $x = 2i$, and $x = -2i$

d. The fourth-degree polynomial function

$$\begin{aligned}f(x) &= x^4 - 1 \\&= (x - 1)(x + 1)(x - i)(x + i)\end{aligned}$$

has exactly *four* zeros: $x = 1$, $x = -1$, $x = i$, and $x = -i$




Finding Zeros of a Polynomial Function



Finding Zeros of a Polynomial Function

Remember that the n zeros of a polynomial function can be real or complex, and they may be repeated. Example 2 illustrates several cases.



Example 2 – *Real and Complex Zeros of a Polynomial Function*

Confirm that the third-degree polynomial function

$$f(x) = x^3 + 4x$$

has exactly three zeros: $x = 0$, $x = 2i$, and $x = -2i$.

Solution:

Factor the polynomial completely as $x(x - 2i)(x + 2i)$. So, the zeros are

$$x(x - 2i)(x + 2i) = 0$$

$$x = 0$$

$$x - 2i = 0$$


$$x = 2i$$



$$x + 2i = 0$$



$$: -2i$$



Example 2 – Solution

cont'd

In the graph in Figure 2.33, only the real zero $x = 0$ appears as an x-intercept.




Figure 2.33



Conjugate Pairs




Conjugate Pairs

The two complex zeros are conjugates. That is, they are of the forms $a + bi$ and $a - bi$.

Complex Zeros Occur in Conjugate Pairs

Let $f(x)$ be a polynomial function that has *real coefficients*. If $a + bi$, where $b \neq 0$, is a zero of the function, then the conjugate $a - bi$ is also a zero of the function.

Be sure you see that this result is true only when the polynomial function has *real coefficients*. For instance, the result applies to the function $f(x) = x^2 + 1$, but not to the function $g(x) = x - i$.



Example 4 – *Finding a Polynomial with Given Zeros*

Find a fourth-degree polynomial function with real coefficients that has -1 , -1 and $3i$ as zeros.

Solution:

Because $3i$ is a zero and the polynomial is stated to have real coefficients, you know that the conjugate $-3i$ must also be a zero.

So, from the Linear Factorization Theorem, $f(x)$ can be written as

$$f(x) = a(x + 1)(x + 1)(x - 3i)(x + 3i)$$



Example 4 – *Solution*

cont'd

For simplicity, let $a = 1$ to obtain

$$\begin{aligned} f(x) &= (x^2 + 2x + 1)(x^2 + 9) \\ &= x^4 + 2x^3 + 10x^2 + 18x + 9. \end{aligned}$$



Factoring a Polynomial



Factoring a Polynomial

Factors of a Polynomial

Every polynomial of degree $n > 0$ with real coefficients can be written as the product of linear and quadratic factors with real coefficients, where the quadratic factors have no real zeros.

A quadratic factor with no real zeros is said to be **prime** or **irreducible over the reals**. Be sure you see that this is not the same as being irreducible over the rationals. For example, the quadratic

$$x^2 + 1 = (x - i)(x + i)$$

is irreducible over the reals (and therefore over the rationals).



Factoring a Polynomial

On the other hand, the quadratic

$$x^2 - 2 = (x - \sqrt{2})(x + \sqrt{2})$$

is irreducible over the rationals, but *reducible* over the reals.



Example 6 – *Factoring a Polynomial*

Write the polynomial $f(x) = x^4 - x^2 - 20$

- a. as the product of factors that are irreducible over the *rationals*,
- b. as the product of linear factors and quadratic factors that are irreducible over the reals, and
- c. in completely factored form.



Example 6 – *Solution*

- a.** Begin by factoring the polynomial into the product of two quadratic polynomials.

$$x^4 - x^2 - 20 = (x^2 - 5)(x^2 + 4)$$

Both of these factors are irreducible over the rationals.

- b.** By factoring over the reals, you have

$$x^4 - x^2 - 20 = (x^2 + 4)(x + \sqrt{5})(x - \sqrt{5})$$

where the quadratic factor is irreducible over the reals.


- c.** In completely factored form, you have

$$x^4 - x^2 - 20 = (x - 2i)(x + 2i)(x + \sqrt{5})(x - \sqrt{5})$$



Factoring a Polynomial

The results and theorems have been stated in terms of zeros of polynomial functions. Be sure you see that the same results could have been stated in terms of solutions of polynomial equations.



Example 7 – *Finding The Zeros of Polynomial Function*

Find all the zeros of

$$f(x) = x^4 - 3x^3 + 6x^2 + 2x - 60$$

given that $1 + 3i$ is a zero of f .

Solution:

Because complex zeros occur in conjugate pairs, you know that $1 - 3i$ is also a zero of f . This means that both

$$x - (1 + 3i) \text{ and } x - (1 - 3i)$$

are factors of f .



Example 7 – *Solution*

cont'd

Multiplying these two factors produces

$$\begin{aligned}[x - (1 + 3i)][x - (1 - 3i)] &= [(x - 1) - 3i][(x - 1) + 3i] \\ &= (x - 1)^2 - 9i^2 \\ &= x^2 - 2x + 10.\end{aligned}$$

Using long division, you can divide $x^2 - 2x + 10$ into f to obtain the following.



Example 7 – Solution

cont'd

Using long division, you can divide $x^2 - 2x + 10$ into f to obtain the following.

$$\begin{array}{r} \overline{x^4 - 3x^3 + 6x^2 + 2x - 60} \\ \underline{x^4 - 2x^3 + 10x^2} \\ -x^3 - 4x^2 + 2x \\ \underline{-x^3 + 2x^2 - 10x} \\ -6x^2 + 12x - 60 \\ \underline{-6x^2 + 12x - 60} \\ 0 \end{array}$$



Example 7 – *Solution*

cont'd

So, you have

$$\begin{aligned} f(x) &= (x^2 - 2x + 10)(x^2 - x - 6) \\ &= (x^2 - 2x + 10)(x - 3)(x + 2) \end{aligned}$$

and you can conclude that the zeros of f are

$$x = 1 + 3i, x = 1 - 3i, x = 3, x = -2.$$