

7.5

Operations with Matrices




What You Should Learn

- Decide whether two matrices are equal.
- Add and subtract matrices and multiply matrices by scalars.
- Multiply two matrices.
- Use matrix operations to model and solve real-life problems.



Equality of Matrices



Equality of Matrices

This section introduces some fundamentals of matrix theory. It is standard mathematical convention to represent matrices in any of the following three ways. **Do not copy.**

Representation of Matrices

1. A matrix can be denoted by an uppercase letter such as

A , B , or C .

2. A matrix can be denoted by a representative element enclosed in brackets, such as

$[a_{ij}]$, $[b_{ij}]$, or $[c_{ij}]$.

3. A matrix can be denoted by a rectangular array of numbers such as

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \cdots & a_{mn} \end{bmatrix}.$$




Equality of Matrices

Two matrices

$$A = [a_{ij}] \quad \text{and} \quad B = [b_{ij}]$$

are **equal** when they have the same dimension ($m \times n$) and all of their corresponding entries are equal.



Example 1 – *Equality of Matrices*

Solve for a_{11} , a_{12} , a_{21} , and a_{22} in the following matrix equation. You do not need to copy this.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ -3 & 0 \end{bmatrix}$$

Solution:

Because two matrices are equal only when their corresponding entries are equal, you can conclude that

$$a_{11} = 2, \quad a_{12} = -1, \quad a_{21} = -3, \quad \text{and} \quad a_{22} = 0.$$



Equality of Matrices

Be sure you see that for two matrices to be equal, they must have the same dimension *and* their corresponding entries must be equal. You do not need to copy this again.

For instance,

$$\begin{bmatrix} 2 & -1 \\ \sqrt{4} & \frac{1}{2} \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ 2 & 0.5 \end{bmatrix} \quad \text{but} \quad \begin{bmatrix} 2 & -1 \\ 3 & 4 \\ 0 & 0 \end{bmatrix} \neq \begin{bmatrix} 2 & -1 \\ 3 & 4 \end{bmatrix}.$$



Matrix Addition and Scalar Multiplication



Matrix Addition and Scalar Multiplication


You can add two matrices (of the same dimension) by adding their corresponding entries.

Definition of Matrix Addition

If $A = [a_{ij}]$ and $B = [b_{ij}]$ are matrices of dimension $m \times n$, then their sum is the $m \times n$ matrix given by

$$A + B = [a_{ij} + b_{ij}].$$

The sum of two matrices of different dimensions is undefined.



Example 2 – Addition of Matrices

a.
$$\begin{bmatrix} -1 & 2 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 3 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} -1 + 1 & 2 + 3 \\ 0 + (-1) & 1 + 2 \end{bmatrix} = \begin{bmatrix} 0 & 5 \\ -1 & 3 \end{bmatrix}$$

b.
$$\begin{bmatrix} 1 \\ -3 \\ -2 \end{bmatrix} + \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

c. The sum of

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 4 & 0 & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 1 \\ -1 & 3 \end{bmatrix}$$

is undefined because A is of dimension 2×3 and B is of dimension 2×2 .



Matrix Addition and Scalar Multiplication

In operations with matrices, numbers are usually referred to as **scalars**. In this text, scalars will always be real numbers. **You can multiply a matrix A by a scalar c by multiplying each entry in A by c .**

Definition of Scalar Multiplication

If $A = [a_{ij}]$ is an $m \times n$ matrix and c is a scalar, then the **scalar multiple** of A by c is the $m \times n$ matrix given by

$$cA = [ca_{ij}].$$



Example 3 – *Scalar Multiplication*

For the following matrix, find $3A$.

$$A = \begin{bmatrix} 2 & 2 & 4 \\ -3 & 0 & -1 \\ 2 & 1 & 2 \end{bmatrix}$$

Solution:

$$3A = 3 \begin{bmatrix} 2 & 2 & 4 \\ -3 & 0 & -1 \\ 2 & 1 & 2 \end{bmatrix} = \begin{bmatrix} 3(2) & 3(2) & 3(4) \\ 3(-3) & 3(0) & 3(-1) \\ 3(2) & 3(1) & 3(2) \end{bmatrix} = \begin{bmatrix} 6 & 6 & 12 \\ -9 & 0 & -3 \\ 6 & 3 & 6 \end{bmatrix}$$



Matrix Addition and Scalar Multiplication

Please read slides #13-17 for important information, but you do not need to copy them down.

The properties of matrix addition and scalar multiplication are similar to those of addition and multiplication of real numbers.

One important property of addition of real numbers is that the number 0 is the additive identity.

That is, $c + 0 = c$ for any real number. For matrices, a similar property holds.



Matrix Addition and Scalar Multiplication

For matrices, a similar property holds. That is, if A is an $m \times n$ matrix and O is the $m \times n$ **zero matrix** consisting entirely of zeros, then $A + O = A$.

In other words, O is the **additive identity** for the set of all $m \times n$ matrices. For example, the following matrices are the additive identities for the sets of all 2×3 and 2×2 matrices.

and $O = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

2×3 zero matrix

$$O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

2×2 zero matrix



Matrix Addition and Scalar Multiplication

Properties of Matrix Addition and Scalar Multiplication

Let A , B , and C be $m \times n$ matrices and let c and d be scalars.

1. $A + B = B + A$

Commutative Property of Matrix Addition

2. $A + (B + C) = (A + B) + C$

Associative Property of Matrix Addition

3. $(cd)A = c(dA)$

Associative Property of Scalar Multiplication

4. $1A = A$

Scalar Identity

5. $A + O = A$


Additive Identity

6. $c(A + B) = cA + cB$

Distributive Property

7. $(c + d)A = cA + dA$

Distributive Property



Example 5 – *Using the Distributive Property*

$$\begin{aligned} 3\left(\begin{bmatrix} -2 & 0 \\ 4 & 1 \end{bmatrix} + \begin{bmatrix} 4 & -2 \\ 3 & 7 \end{bmatrix}\right) &= 3\begin{bmatrix} -2 & 0 \\ 4 & 1 \end{bmatrix} + 3\begin{bmatrix} 4 & -2 \\ 3 & 7 \end{bmatrix} \\ &= \begin{bmatrix} -6 & 0 \\ 12 & 3 \end{bmatrix} + \begin{bmatrix} 12 & -6 \\ 9 & 21 \end{bmatrix} \\ &= \begin{bmatrix} 6 & -6 \\ 21 & 24 \end{bmatrix} \end{aligned}$$



Matrix Addition and Scalar Multiplication

The algebra of real numbers and the algebra of matrices have many similarities. For example, compare the following solutions.

Real Numbers $m \times n$ *Matrices*

(Solve for x .) *(Solve for X .)*

$$x + a = b \quad X + A = B$$

$$x + a + (-a) = b + (-a)$$


$$x + 0 = b - a$$

$$x = b - a$$

$$X + A + (-A) = B + (-A)$$

$$X + O = B - A$$

$$X = B - A$$



Example 6 – Solving a Matrix Equation


Solve for X in the equation

$$3X + A = B$$

where

$$\text{and } A = \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}$$

$$B = \begin{bmatrix} -3 & 4 \\ 2 & 1 \end{bmatrix}.$$



Example 6 – *Solution*

Begin by solving the equation for X to obtain

$$3X = B - A$$

$$X = \frac{1}{3}(B - A)$$

Now, using the A matrices and B , you have

$$X = \frac{1}{3} \left(\begin{bmatrix} -3 & 4 \\ 2 & 1 \end{bmatrix} - \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix} \right)$$

Substitute the matrices

$$= \frac{1}{3} \begin{bmatrix} -4 & 6 \\ 2 & -2 \end{bmatrix}$$

Subtract matrix A from matrix B

$$= \begin{bmatrix} -\frac{4}{3} & 2 \\ \frac{2}{3} & -\frac{2}{3} \end{bmatrix}$$

Multiply the resulting matrix by

$\frac{1}{3}$



Matrix Multiplication



Matrix Multiplication

Another basic matrix operation is **matrix multiplication**. You will see later, that this definition of the product of two matrices has many practical applications.

Definition of Matrix Multiplication

If $A = [a_{ij}]$ is an $m \times n$ matrix and $B = [b_{ij}]$ is an $n \times p$ matrix, then the product AB is an $m \times p$ matrix given by

$$AB = [c_{ij}]$$

where $c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + a_{i3}b_{3j} + \cdots + a_{in}b_{nj}$.



Matrix Multiplication

The definition of matrix multiplication indicates a *row-by-column multiplication*, where the entry in the i th row and j th column of the product AB is obtained by multiplying the entries in the i th row A of by the corresponding entries in the j th column of B and then adding the results.

Look at next slide so you can multiply matrices, but do not copy it.




Matrix Multiplication

The general pattern for matrix multiplication is as follows.

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \vdots & & \vdots \\ a_{i1} & a_{i2} & a_{i3} & \dots & a_{in} \\ \vdots & \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1j} & \dots & b_{1p} \\ b_{21} & b_{22} & \dots & b_{2j} & \dots & b_{2p} \\ b_{31} & b_{32} & \dots & b_{3j} & \dots & b_{3p} \\ \vdots & \vdots & & \vdots & & \vdots \\ b_{n1} & b_{n2} & \dots & b_{nj} & \dots & b_{np} \end{bmatrix} = \begin{bmatrix} c_{11} & c_{12} & \dots & c_{1j} & \dots & c_{1p} \\ c_{21} & c_{22} & \dots & c_{2j} & \dots & c_{2p} \\ \vdots & \vdots & & \vdots & & \vdots \\ c_{i1} & c_{i2} & \dots & c_{ij} & \dots & c_{ip} \\ \vdots & \vdots & & \vdots & & \vdots \\ c_{m1} & c_{m2} & \dots & c_{mj} & \dots & c_{mp} \end{bmatrix}$$

$a_{i1}b_{1j} + a_{i2}b_{2j} + a_{i3}b_{3j} + \dots + a_{in}b_{nj} = c_{ij}$




Example 7 – Finding the Product of Two Matrices

Find the product AB using $A = \begin{bmatrix} -1 & 3 \\ 4 & -2 \\ 5 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} -3 & 2 \\ -4 & 1 \end{bmatrix}$.

Solution:

First, note that the product is defined because the number of columns of A is equal to the number of rows of B . Moreover, the product AB has dimension 3×2 .

To find the entries of the product, multiply each row of A by each column of B .



Example 7 – Solution

cont'd

$$AB = \begin{bmatrix} -1 & 3 \\ 4 & -2 \\ 5 & 0 \end{bmatrix} \begin{bmatrix} -3 & 2 \\ -4 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} (-1)(-3) + (3)(-4) & (-1)(2) + (3)(1) \\ (4)(-3) + (-2)(-4) & (4)(2) + (-2)(1) \\ (5)(-3) + (0)(-4) & (5)(2) + (0)(1) \end{bmatrix}$$

$$= \begin{bmatrix} -9 & 1 \\ -4 & 6 \\ -15 & 10 \end{bmatrix}$$



Matrix Multiplication

The general pattern for matrix multiplication is as follows, but you do not need to write it or the next slide down.

Properties of Matrix Multiplication

Let A , B , and C be matrices and let c be a scalar.

1. $A(BC) = (AB)C$

Associative Property of Matrix Multiplication

2. $A(B + C) = AB + AC$

Left Distributive Property

3. $(A + B)C = AC + BC$

Right Distributive Property

4. $c(AB) = (cA)B = A(cB)$

Associative Property of Scalar Multiplication



Matrix Multiplication

Definition of Identity Matrix

The $n \times n$ matrix that consists of 1's on its main diagonal and 0's elsewhere is called the **identity matrix of dimension $n \times n$** and is denoted by

$$I_n = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}. \quad \text{Identity matrix}$$

Note that an identity matrix must be *square*. When the dimension is understood to be $n \times n$, you can denote I_n simply by I .



Matrix Multiplication

If A is an $n \times n$ matrix, then the identity matrix has the property that $AI_n = A$ and $I_nA = A$.

For example,

$$\begin{bmatrix} 3 & -2 & 5 \\ 1 & 0 & 4 \\ -1 & 2 & -3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -2 & 5 \\ 1 & 0 & 4 \\ -1 & 2 & -3 \end{bmatrix} \quad AI = A$$