dent's Name:D.			Place child's
pol: Tea			photo here
tory:			
Asthma: YES (Higher risk for severe reaction)	□NO TREATMEN	T	-
Any SEVERE SYMPTOMS after suspected or known			HRINE IMMEDIATEL
ingestion:  One or more of the following: LUNG: Short of breath, wheeze, repetitive cough HEART: Pale, blue, faint, weak pulse, dizzy, confuse THROAT: Tight, hoarse, trouble breathing/swallowing MOUTH: Obstructive swelling (tongue and/or lips) SKIN: Many hives over body  Or combination of symptoms from different body areas SKIN: Hives, itchy rashes, swelling (e.g., eyes, lips)		<ul> <li>2. Call 911</li> <li>3. Begin monitoring</li> <li>4. Give additional m</li> <li>Antihistamine</li> <li>Inhaler (quick</li> <li>*Antihistamine &amp; quick</li> </ul>	(see box below) nedications:* e k relief) if asthma ck relief inhalers are no o treat a severe reaction
GUT: Vomiting, crampy pain		1. GIVE ANTIHIST	
MILD SYMPTOMS ONLY:  MOUTH: Itchy mouth  SKIN: A few hives around mouth/face, mild itch  GUT: Mild nausea/discomfort		<ul> <li>2. Stay with student; alert healthcare professionals and parent</li> <li>3. If symptoms progress (see above), USEPINEPHRINE</li> <li>4. Begin monitoring</li> </ul>	
<u>DOSAGE</u>			
<b>Epinephrine</b> : inject intramuscularly using autoinjector	-		mg
Administer 2 <sup>nd</sup> dose if symptoms do not improve in <b>Antihistamine</b> : (brand and dose)			
If Asthmatic: (brand and dose)			
Student's condition warrants accommodations from	food servic	e (nlesse complete the hotte	om of the back nage)
Student has been instructed and is capable of carrying			<u></u>
		Phone Number:	
Provider's Signature:		Date:	
♦ STEP 2: EME		_	
<ol> <li>If epinephrine given, call 911. State that an alle epinephrine, oxygen, or other medications may</li> </ol>	be needed	d.	
2. Parent:			
		Number(s)	
a			
	1\	2)	

Parent/Guardian's Signature:	Date:
School Nurse:	Date:

Student Name:	DOB:
TRAINED/DELEGATED ST	Room
<ul> <li>Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10.</li> <li>Remove the EpiPen® unit and massage the injection area for 10 seconds.</li> </ul>	<ul> <li>Pull off Red safety guard.</li> <li>Place black end against the middle of the outer thigh (through clothing, if needed), then press firmly, and hold in</li> </ul>
Once epinephrine i Student should remain lying down Additional information:	n or in a comfortable position.
MEDICAL STATEMENT FOR SCHO  I certify that the student's food allergy rises to the level of a d  1. List foods to be omitted:  2. Indicate food modification/substitutions:	isability & qualifies for school meal modification.
Signature by a licensed MD or DO only	Date

C.R.S. 22-2-135(3)(b) 11/2012