

9.4

Parametric Equations



What You Should Learn

- Evaluate sets of parametric equations for given values of the parameter
- Graph curves that are represented by sets of parametric equations
- Rewrite sets of parametric equations as single rectangular equations by eliminating the parameter



Plane Curves



Plane Curves

Up to this point, you have been representing a graph by a single equation involving *two* variables such as x and y .

In this section, you will study situations in which it is useful to introduce a *third* variable to represent a curve in the plane.

To see the usefulness of this procedure, consider the path of an object that is propelled into the air at an angle of 45° .



Plane Curves

When the initial velocity of the object is 48 feet per second, it can be shown that the object follows the parabolic path

$$y = -\frac{x^2}{72} + x$$

Rectangular equation

as shown in Figure 9.42.


Rectangular equation:

$$y = -\frac{x^2}{72} + x$$

Parametric equations:

$$x = 24\sqrt{2}t$$

$$y = -16t^2 + 24\sqrt{2}t$$



Curvilinear motion: two variables for position, one variable for time

Figure 9.42



Plane Curves

However, this equation does not tell the whole story. Although it does tell you *where* the object has been, it does not tell you *when* the object was at a given point (x, y) on the path.

To determine this time, you can introduce a third variable t , **called a parameter**. It is possible to write both x and y as functions of t to obtain the **parametric equations**

$$x = 24\sqrt{2}t$$

Parametric equation for x

$$y = -16t^2 + 24\sqrt{2}t.$$

Parametric equation for y



Plane Curves

From this set of equations you can determine that at time $t = 0$, the object is at the point $(0, 0)$.

Similarly, at time $t = 1$, the object is at the point

$$(24\sqrt{2}, 24\sqrt{2} - 16)$$

and so on.



Plane Curves

Definition of a Plane Curve

If f and g are continuous functions of t on an interval I , then the set of ordered pairs

$$(f(t), g(t))$$

is a **plane curve** C . The equations given by

$$x = f(t) \quad \text{and} \quad y = g(t)$$

are **parametric equations** for C , and t is the **parameter**.



Graphs of Plane Curves




Graphs of Plane Curves

One way to sketch a curve represented by a pair of parametric equations is to plot points in the xy -plane.

Each set of coordinates (x, y) is determined from a value chosen for the parameter t .

By plotting the resulting **points in the order of *increasing* values of t , you trace the curve in a specific direction.** This is called the **orientation** of the curve.




Example 1 – *Sketching a Plane Curve*

Sketch the curve given by the parametric equations

$$x = t^2 - 4 \quad \text{and} \quad y = \frac{t}{2}, \quad -2 \leq t \leq 3.$$

Describe the orientation of the curve.



Example 1 – *Solution*

Using values of t in the interval, the parametric equations yield the points (x, y) shown in the table.

t	-2	-1	0	1	2	3
x	0	-3	-4	-3	0	5
y	-1	$-\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{3}{2}$

Example 1 – Solution

cont'd

By plotting these points in the order of increasing t , you obtain the curve shown in Figure 9.43.





Figure 9.43



Example 1 – *Solution*

cont'd

The **arrows on the curve indicate its orientation** as t increases from -2 to 3 .

So, when a particle moves on this curve, it would start at $(0, -1)$ and then move along the curve to the point $(5, \frac{3}{2})$.



Eliminating the Parameter



Eliminating the Parameter

Many curves that are represented by sets of parametric equations have graphs that can also be represented by rectangular equations (in x and y). The process of finding the rectangular equation is called **eliminating the parameter (using substitution)**.

Parametric equations



Solve for t in one equation.



Substitute in second equation.



Rectangular equation

$$x = t^2 - 4$$

$$t = 2y$$

$$x = (2y)^2 - 4$$

$$x = 4y^2 - 4$$


$$y = \frac{1}{2}t$$



Eliminating the Parameter

Now you can recognize that the equation $x = 4y^2 - 4$ represents a parabola with a horizontal axis and vertex at $(-4, 0)$.

When converting equations from parametric to rectangular form, you may need to alter the domain of the rectangular equation so that its graph matches the graph of the **parametric equations**. This situation is demonstrated in Example 3.



Example 3 – *Eliminating the Parameter*

Identify the curve represented by the equations

$$x = \frac{1}{\sqrt{t+1}} \quad \text{and} \quad y = \frac{t}{t+1}.$$

Solution:

Solving for t in the equation for x produces

$$x^2 = \frac{1}{t+1} \quad \Rightarrow \quad \frac{1}{x^2} = t+1 \quad \Rightarrow \quad \frac{1}{x^2} - 1 = t.$$

Example 3 – Solution

cont'd

Substituting in the equation for y , you obtain the rectangular equation

$$y = \frac{t}{t+1} = \frac{\frac{1}{x^2} - 1}{\frac{1}{x^2} - 1 + 1} = \frac{1 - x^2}{x^2} \cdot \frac{x^2}{x^2} = 1 - x^2.$$

From the rectangular equation, you can recognize that the curve is a parabola that opens downward and has its vertex at $(0, 1)$, as shown in Figure 9.54.




Figure 9.54

Example 3 – Solution

cont'd

The rectangular equation is defined for all values of x . The parametric equation for x however, is defined only when $t > -1$.

From the graph of the parametric equations, you can see that x is always positive, as shown in Figure 9.55.




Figure 9.55

Example 3 – Solution

cont'd

So, you should restrict the domain of x to positive values, as shown in Figure 9.56.




Figure 9.56



Finding Parametric Equations for a Graph



Finding Parametric Equations for a Graph

You have been studying techniques for sketching the graph represented by a set of parametric equations.

Now consider the *reverse* problem—that is, how can you find a set of parametric equations for a given graph or a given physical description?

From the discussion following Example 1, you know that such a representation is not unique (this means that there can be more than one correct answer).



Finding Parametric Equations for a Graph

That is, the equations

$$x = 4t^2 - 4 \quad \text{and} \quad y = t, \quad -1 \leq t \leq \frac{3}{2}$$

produced the same graph as the equations

$$x = t^2 - 4 \quad \text{and} \quad y = \frac{t}{2}, \quad -2 \leq t \leq 3.$$

Example 4 – Solution

cont'd

The graph of these equations is shown in Figure 9.57.




Figure 9.57