

**11.2**

## **Techniques for Evaluating Limits**



# What You Should Learn

- Use the dividing out technique to evaluate limits of functions
- Use the rationalizing technique to evaluate limits of functions
- Use technology to approximate limits of functions graphically and numerically



# What You Should Learn

- Evaluate one-sided limits of functions
- Evaluate limits of difference quotients from calculus



# Dividing Out Technique



# Dividing Out Technique

We have studied several types of functions whose limits can be evaluated by direct substitution.

In this section, you will study several techniques for evaluating limits of functions for which direct substitution fails.

Suppose you were asked to find the following limit.


$$\lim_{x \rightarrow -3} \frac{x^2 + x - 6}{x + 3}$$



# Dividing Out Technique

Direct substitution fails because  $-3$  is a zero of the denominator. By using a table, however, it appears that the limit of the function as  $x$  approaches  $-3$  is  $-5$ .

|                             |         |          |           |      |           |          |         |
|-----------------------------|---------|----------|-----------|------|-----------|----------|---------|
| $x$                         | $-3.01$ | $-3.001$ | $-3.0001$ | $-3$ | $-2.9999$ | $-2.999$ | $-2.99$ |
| $\frac{x^2 + x - 6}{x + 3}$ | $-5.01$ | $-5.001$ | $-5.0001$ | $?$  | $-4.9999$ | $-4.999$ | $-4.99$ |



# Example 1 – *Dividing Out Technique*

Find the limit.


$$\lim_{x \rightarrow -3} \frac{x^2 + x - 6}{x + 3}$$

**Solution:**

Begin by factoring the numerator and dividing out any common factors.

$$\lim_{x \rightarrow -3} \frac{x^2 + x - 6}{x + 3} = \lim_{x \rightarrow -3} \frac{(x - 2)(x + 3)}{x + 3}$$

Factor numerator.



# Example 1 – Solution

cont'd

$$= \lim_{x \rightarrow -3} \frac{(x - 2)(\cancel{x + 3})}{\cancel{x + 3}}$$

Divide out common factor.

$$(x - 2) \quad = \lim_{x \rightarrow -3}$$

Simplify.

$$= -3 - 2$$

Direct substitution

$$= -5$$

Simplify.





# Dividing Out Technique

This procedure for evaluating a limit is called the **dividing out technique**.

The validity of this technique stems from the fact that when two functions agree at all but a single number  $c$ , they must have identical limit behavior at  $x = c$ .

In Example 1, the functions given by

$$f(x) \quad \text{and} \quad g = \frac{x^2 + x - 6}{x + 3}$$

agree at all values of  $x$  other than  $x = -3$ .

So, you can use  $g(x)$  to find the limit of  $f(x)$ .



# Dividing Out Technique

The dividing out technique should be applied only when direct substitution produces 0 in both the numerator *and* the denominator.

An expression such as  $\frac{0}{0}$  has no meaning as a real number.

It is called an **indeterminate form** because you cannot, from the form alone, determine the limit.



# Dividing Out Technique

When you try to evaluate a limit of a rational function by direct substitution and encounter this form, you can conclude that the numerator and denominator must have a common factor.

After factoring and dividing out, you should try direct substitution again.



# One-Sided Limits



# One-Sided Limits

The limit of  $f(x)$  as  $x \rightarrow c$  does not exist when the function  $f(x)$  approaches a different number from the left side of  $c$  than it approaches from the right side of  $c$ .

This type of behavior can be described more concisely with the concept of a **one-sided limit**.

$$\lim_{x \rightarrow c^-} f(x) = L_1 \text{ or } f(x) \rightarrow L_1 \text{ as } x \rightarrow c^-$$

Limit from the left

$$\lim_{x \rightarrow c^+} f(x) = L_2 \text{ or } f(x) \rightarrow L_2 \text{ as } x \rightarrow c^+$$

Limit from the right

## Example 6 – Evaluating One-Sided Limits

Find the limit as  $x \rightarrow 0$  from the left and the limit as  $x \rightarrow 0$  from the right for

$$f(x) = \frac{|2x|}{x}$$

**Solution:**

From the graph of  $f$ , shown in Figure 11.15, you can see that  $f(x) = -2$  for all  $x < 0$ .





Figure 11.15



# Example 6 – *Solution*

cont'd

So, the limit from the left is

$$= -2. \quad \lim_{x \rightarrow 0^-} \frac{|2x|}{x}$$

Limit from the left

Because  $f(x) = 2$  for all  $x > 0$ , the limit from the right is

$$= 2. \quad \lim_{x \rightarrow 0^+} \frac{|2x|}{x}$$

Limit from the right



# One-Sided Limits

## Existence of a Limit

If  $f$  is a function and  $c$  and  $L$  are real numbers, then

$$\lim_{x \rightarrow c} f(x) = L$$

if and only if both the left and right limits *exist* and are *equal* to  $L$ .





# A Limit from Calculus



# A Limit from Calculus

A Limit from Calculus In the next section, you will study an important type of limit from calculus—the limit of a *difference quotient*.



## Example 9 – *Evaluating a Limit from Calculus*


For the function given by  $f(x) = x^2 - 1$ , find

$$\lim_{h \rightarrow 0} \frac{f(3 + h) - f(3)}{h}$$

**Solution:**

Direct substitution produces an indeterminate form.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(3 + h) - f(3)}{h} &= \lim_{h \rightarrow 0} \frac{[(3 + h)^2 - 1] - [(3)^2 - 1]}{h} \\ &= \lim_{h \rightarrow 0} \frac{9 + 6h + h^2 - 1 - 9 + 1}{h} \end{aligned}$$




# Example 9 – Solution

cont'd

$$\begin{aligned} &= \lim_{h \rightarrow 0} \frac{6h + h^2}{h} \\ &= \frac{0}{0} \end{aligned}$$

By factoring and dividing out, you obtain the following.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(3 + h) - f(3)}{h} &= \lim_{h \rightarrow 0} \frac{6h + h^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(6 + h)}{h} \end{aligned}$$



# Example 9 – *Solution*

cont'd

$$= (6 + h) \quad \lim_{h \rightarrow 0}$$

$$= 6 + 0$$

$$= 6$$

So, the limit is 6.