

Name \_\_\_\_\_

Solve the problem.

1) Suppose that  $h$  is continuous and that  $\int_{-2}^4 h(x) dx = 2$  and  $\int_4^8 h(x) dx = -8$ . Find  $\int_{-2}^8 h(x) dx$  and

$$\int_8^{-2} h(x) dx$$

2) Suppose that  $g$  is continuous and that  $\int_4^7 g(x) dx = 9$  and  $\int_4^9 g(x) dx = 14$ .

Find  $\int_9^7 g(x) dx$  and Find  $\int_4^4 f(x) dx$ .

3) Suppose that  $f$  and  $g$  are continuous and that  $\int_7^{11} f(x) dx = -2$  and  $\int_7^{11} g(x) dx = 9$ .

Find  $\int_7^{11} [5f(x) + g(x)] dx$ .

Find the average value over the given interval. SHOW ALL WORK.

4)  $y = \frac{1}{x}$ ;  $[1, e]$

Find  $dy/dx$ .

5) If  $y = \int_{x^4}^1 12t^5 dt$  find  $dy/dx$

6)  $y = \int_{\cos x}^{\sin x} \frac{1}{9 - t^2} dt$  find  $dy/dx$

7) If  $\int_1^3 f(x)dx = 10$ , find  $\int_1^3 (f(x) + 5) dx$

Evaluate the definite integral using areas or antiderivatives. SHOW ALL WORK.

8)  $\int_{-1}^6 6 dx$

9)  $\int_1^2 (2x^3 - 6x^{-2}) dx$

Evaluate the integral. SHOW ALL WORK.

$$10) \int_0^{\pi/2} 17 \sin x \, dx$$

$$11) \int_0^1 (x^4 - x^{\frac{1}{5}}) dx$$


$$12) \int_{\pi/4}^{3\pi/4} 8 \sec \theta \tan \theta \, d\theta$$

$$13) \int_1^2 (2e^x - 8x^{-2}) \, dx$$

$$14) \int_1^2 \frac{1-x}{x^2} \, dx$$

15) The graph of the function,  $f$ , is given below with position defined as follows.

$$g = \int_0^t f(x) dx$$



- a) Determine the relative minimum of  $g$ . Justify your answer.
  
- b) Find the absolute maximum of  $g$  on the interval  $[0, 10]$ ? Justify your answer.
  
- c) Determine when  $g$  is concave down on the interval  $[0, 10]$ ? Justify your answers.
  
- d) Determine the intervals where  $g$  is increasing. Justify your answer.
  
- e) Write the equation of the tangent line of  $g$  at  $t = 10$ .

Solve the problem.

- 16) Use the data below to approximate the area under the curve using Midpoint Riemann Sums with 3 sub-intervals.

T	0	2	4	6	8	10	12
P(t)	0	46	53	57	60	62	63

- 17) Let  $f$  be a function that is twice differentiable for all real numbers. The table gives values of  $f$  for  $s$  points in the closed interval  $2 \leq x \leq 13$

x	2	3	5	8	13
f(x)	1	4	-2	3	6

Use a trapezoid approximation to find  $\int_2^{13} f(x)$

18)

$t$ (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

Johanna jogs along a straight path. For  $0 \leq t \leq 40$ , Johanna's velocity is given by a differential function  $v$ . Selected values of  $v(t)$ , where  $t$  is measured in minutes and  $v(t)$  measured in meters per minute, are given in the table above.

A) Use the data in the table to estimate the value of  $v'(22)$

B) Approximate the value of  $\frac{1}{40} \int_0^{40} v(t) dt$  using a right Riemann sum with four sub-intervals indicated in the table. Using correct units, explain the meaning of the definite integral  $\frac{1}{40} \int_0^{40} v(t) dt$  in the context of the problem.

C) Bob is riding his bicycle along the same path. For  $0 \leq t \leq 10$ , Bob's velocity is modeled by  $B(t) = t^3 - 6t^2 + 300$ , where  $t$  is measured in minutes and  $B(t)$  is measured in meters per minute. Find Bob's acceleration at time  $t = 4$ .

D) Based on the model  $B$  from part (c), find Bob's average velocity during the interval  $0 \leq t \leq 5$ .

Determine the intervals of Increase and Decrease. Then use this information to determine any Local Extrema. Justify your explanation

19)  $f(x) = x^3 - 3x^2 - 9x + 3$

At the given point, find the equation of the line that is tangent to the curve.

20)  $x^2 + y^2 - 2x + 4y = 8$ , tangent at (2, 4)

21) Find  $dy/dx$  when

$$y = \frac{\sin(7x)}{5x}$$