

Chapter 10 FRQ Classwork

1.

At time  $t \geq 0$ , a particle moving along a curve in the  $xy$ -plane has position  $(x(t), y(t))$  with velocity vector  $v(t) = (\cos(t^2), e^{0.5t})$ . At  $t = 1$ , the particle is at the point  $(3, 5)$ .

- (a) Find the  $x$ -coordinate of the position of the particle at time  $t = 2$ .
- (b) For  $0 < t < 1$ , there is a point on the curve at which the line tangent to the curve has a slope of 2. At what time is the object at that point?
- (c) Find the time at which the speed of the particle is 3.
- (d) Find the total distance traveled by the particle from time  $t = 0$  to time  $t = 1$ .

$$(a) \quad x(2) = 3 + \int_1^2 \cos(t^2) dt = 2.557 \text{ (or 2.556)}$$

$$3 : \begin{cases} 1 : \text{integral} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$$

$$(b) \quad \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{e^{0.5t}}{\cos(t^2)}$$

$$\frac{e^{0.5t}}{\cos(t^2)} = 2$$

$$t = 0.840$$

$$2 : \begin{cases} 1 : \text{slope in terms of } t \\ 1 : \text{answer} \end{cases}$$

$$(c) \quad \text{Speed} = \sqrt{\cos^2(t^2) + e^t}$$

$$\sqrt{\cos^2(t^2) + e^t} = 3$$

$$t = 2.196 \text{ (or 2.195)}$$


$$2 : \begin{cases} 1 : \text{speed in terms of } t \\ 1 : \text{answer} \end{cases}$$

$$(d) \quad \text{Distance} = \int_0^1 \sqrt{\cos^2(t^2) + e^t} dt = 1.595 \text{ (or 1.594)}$$

$$2 : \begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$$

2.

The graphs of the polar curves  $r = 3$  and  $r = 4 - 2\sin \theta$  are shown in the figure above. The curves intersect when  $\theta = \frac{\pi}{6}$  and  $\theta = \frac{5\pi}{6}$ .



- (a) Let  $S$  be the shaded region that is inside the graph of  $r = 3$  and also inside the graph of  $r = 4 - 2\sin \theta$ . Find the area of  $S$ .
- (b) A particle moves along the polar curve  $r = 4 - 2\sin \theta$  so that at time  $t$  seconds,  $\theta = t^2$ . Find the time  $t$  in the interval  $1 \leq t \leq 2$  for which the  $x$ -coordinate of the particle's position is  $-1$ .
- (c) For the particle described in part (b), find the position vector in terms of  $t$ . Find the velocity vector at time  $t = 1.5$ .

(a)  $\text{Area} = 6\pi + \frac{1}{2} \int_{\pi/6}^{5\pi/6} (4 - 2\sin \theta)^2 d\theta = 24.709$  (or 24.708)

3 : { 1 : integrand  
1 : limits and constant  
1 : answer

(b)  $x = r \cos \theta \Rightarrow x(\theta) = (4 - 2\sin \theta) \cos \theta$   
 $x(t) = (4 - 2\sin(t^2)) \cos(t^2)$   
 $x(t) = -1$  when  $t = 1.428$  (or 1.427)

3 : { 1 :  $x(\theta)$  or  $x(t)$   
1 :  $x(\theta) = -1$  or  $x(t) = -1$   
1 : answer

(c)  $y = r \sin \theta \Rightarrow y(\theta) = (4 - 2\sin \theta) \sin \theta$   
 $y(t) = (4 - 2\sin(t^2)) \sin(t^2)$

3 : { 2 : position vector  
1 : velocity vector

Position vector =  $\langle x(t), y(t) \rangle$   
 $= \langle (4 - 2\sin(t^2)) \cos(t^2), (4 - 2\sin(t^2)) \sin(t^2) \rangle$

$v(1.5) = \langle x'(1.5), y'(1.5) \rangle$   
 $= \langle -8.072, -1.673 \rangle$  (or  $\langle -8.072, -1.672 \rangle$ )

3.

A particle is moving along a curve so that its position at time  $t$  is  $(x(t), y(t))$ , where  $x(t) = t^2 - 4t + 8$  and  $y(t)$  is not explicitly given. Both  $x$  and  $y$  are measured in meters, and  $t$  is measured in seconds. It is known that  $\frac{dy}{dt} = te^{t-3} - 1$ .

- (a) Find the speed of the particle at time  $t = 3$  seconds.  
 (b) Find the total distance traveled by the particle for  $0 \leq t \leq 4$  seconds.  
 (c) Find the time  $t$ ,  $0 \leq t \leq 4$ , when the line tangent to the path of the particle is horizontal. Is the direction of motion of the particle toward the left or toward the right at that time? Give a reason for your answer.  
 (d) There is a point with  $x$ -coordinate 5 through which the particle passes twice. Find each of the following.  
 (i) The two values of  $t$  when that occurs  
 (ii) The slopes of the lines tangent to the particle's path at that point  
 (iii) The  $y$ -coordinate of that point, given  $y(2) = 3 + \frac{1}{e}$

(a) Speed =  $\sqrt{(x'(3))^2 + (y'(3))^2} = 2.828$  meters per second

1 : answer

(b)  $x'(t) = 2t - 4$

Distance =  $\int_0^4 \sqrt{(2t - 4)^2 + (te^{t-3} - 1)^2} dt = 11.587$  or  $11.588$  meters

2 :  $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(c)  $\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = 0$  when  $te^{t-3} - 1 = 0$  and  $2t - 4 \neq 0$

This occurs at  $t = 2.20794$ .

Since  $x'(2.20794) > 0$ , the particle is moving toward the right at time  $t = 2.207$  or  $2.208$ .

3 :  $\begin{cases} 1 : \text{considers } \frac{dy}{dx} = 0 \\ 1 : t = 2.207 \text{ or } 2.208 \\ 1 : \text{direction of motion with reason} \end{cases}$

(d)  $x(t) = 5$  at  $t = 1$  and  $t = 3$

At time  $t = 1$ , the slope is  $\left. \frac{dy}{dx} \right|_{t=1} = \left. \frac{dy/dt}{dx/dt} \right|_{t=1} = 0.432$ .


At time  $t = 3$ , the slope is  $\left. \frac{dy}{dx} \right|_{t=3} = \left. \frac{dy/dt}{dx/dt} \right|_{t=3} = 1$ .

$y(1) = y(3) = 3 + \frac{1}{e} + \int_2^3 \frac{dy}{dt} dt = 4$

3 :  $\begin{cases} 1 : t = 1 \text{ and } t = 3 \\ 1 : \text{slopes} \\ 1 : y\text{-coordinate} \end{cases}$

4.

The graph of the polar curve  $r = 1 - 2\cos \theta$  for  $0 \leq \theta \leq \pi$  is shown above. Let  $S$  be the shaded region in the third quadrant bounded by the curve and the  $x$ -axis.



- (a) Write an integral expression for the area of  $S$ .
- (b) Write expressions for  $\frac{dx}{d\theta}$  and  $\frac{dy}{d\theta}$  in terms of  $\theta$ .
- (c) Write an equation in terms of  $x$  and  $y$  for the line tangent to the graph of the polar curve at the point where  $\theta = \frac{\pi}{2}$ . Show the computations that lead to your answer.

(a)  $r(0) = -1$ ;  $r(\theta) = 0$  when  $\theta = \frac{\pi}{3}$ .

$$\text{Area of } S = \frac{1}{2} \int_0^{\pi/3} (1 - 2\cos \theta)^2 d\theta$$

2 :  $\begin{cases} 1 : \text{limits and constant} \\ 1 : \text{integrand} \end{cases}$

(b)  $x = r \cos \theta$  and  $y = r \sin \theta$

$$\frac{dr}{d\theta} = 2 \sin \theta$$

$$\frac{dx}{d\theta} = \frac{dr}{d\theta} \cos \theta - r \sin \theta = 4 \sin \theta \cos \theta - \sin \theta$$

$$\frac{dy}{d\theta} = \frac{dr}{d\theta} \sin \theta + r \cos \theta = 2 \sin^2 \theta + (1 - 2 \cos \theta) \cos \theta$$

4 :  $\begin{cases} 1 : \text{uses } x = r \cos \theta \text{ and } y = r \sin \theta \\ 1 : \frac{dr}{d\theta} \\ 2 : \text{answer} \end{cases}$

(c) When  $\theta = \frac{\pi}{2}$ , we have  $x = 0$ ,  $y = 1$ .

$$\left. \frac{dy}{dx} \right|_{\theta=\frac{\pi}{2}} = \left. \frac{dy/d\theta}{dx/d\theta} \right|_{\theta=\frac{\pi}{2}} = -2$$

The tangent line is given by  $y = 1 - 2x$ .

3 :  $\begin{cases} 1 : \text{values for } x \text{ and } y \\ 1 : \text{expression for } \frac{dy}{dx} \\ 1 : \text{tangent line equation} \end{cases}$