




What you'll Learn About

- Right Triangle Trigonometry/ Two Famous Triangles
- Evaluating Trig Functions with a calculator/Applications of right triangle trig

The six trigonometric functions

Find the values of all six trigonometric functions.



Assume that θ is an acute angle in a right triangle satisfying the given conditions. Evaluate the remaining trigonometric functions.

A) $\sin \theta = \frac{4}{9}$

B) $\cos \theta = \frac{2}{9}$


C) $\tan \theta = \frac{4}{9}$


D) $\cot \theta = \frac{2}{9}$

E) $\csc \theta = \frac{10}{7}$

F) $\sec \theta = \frac{4}{3}$

45-45-90 Triangle





30-60-90 Triangle

Evaluate using a calculator. Make sure your calculator is in the correct mode. Give answers to 3 decimal places and then draw the triangle that represents the situation.

A) $\sin 53^\circ$

B) $\cos \frac{2\pi}{5}$

C) $\tan 154^\circ$


D) $\cot \frac{\pi}{9}$

E) $\csc 220^\circ$


F) $\sec \frac{8\pi}{5}$

Solve the triangle for the variable shown.

9)



10)



Solve the triangle ABC for all of its unknown parts. Assume C is the right angle.

$$\alpha = 40^\circ \quad a = 10$$

Solve the triangle ABC for all of its unknown parts. Assume C is the right angle.

$$\beta = 62^\circ \quad a = 7$$

Example 6: From a point 340 feet away from the base of the Peachtree Center Plaza in Atlanta, Georgia, the angle of elevation to the top of the building is 65° . Find the height of the building.

--	--

What you'll Learn About

- Trig functions of any angle/Trig functions of real numbers
- Periodic Functions/The Unit Circle

Point P is on the terminal side of angle θ . Evaluate the six trigonometric functions for θ .

A) (5, 4)

B) (-3, 4)

C) (-2, -5)

D) (-4, -1)

E) (0, -3)

F) (3, 0)

Determine the sign (+ or -) of the given value without the use of a calculator.

A) $\sin 53^\circ$

B) $\cos \frac{2\pi}{5}$


C) $\tan 154^\circ$

D) $\cot \frac{\pi}{9}$

E) $\csc 220^\circ$

F) $\sec \frac{8\pi}{5}$

Unit Circle, Fill in the blank



Evaluate without using a calculator by using ratios in a reference triangle.

A) $\sin 120^\circ$

B) $\cos \frac{2\pi}{3}$

C) $\tan \frac{13\pi}{4}$

D) $\cot \frac{-13\pi}{6}$

E) $\csc \frac{7\pi}{4}$

F) $\sec \frac{23\pi}{6}$

Find sine, cosine, and tangent for the given angle.

A) 90°

B) $-\frac{\pi}{2}$

C) 6π

D) $\frac{-7\pi}{2}$

Evaluate without using a calculator

A) Find $\sin \theta$ and $\tan \theta$ if $\cos \theta = \frac{3}{4}$ and $\cot \theta < 0$


B) Find $\sec \theta$ and $\csc \theta$ if $\cot \theta = \frac{-6}{5}$ and $\sin \theta > 0$

--	--


What you'll Learn About

- The basic waves revisited/Sinusoids and Transformations
- Modeling

The graph of $y = \sin x$



The graph of $y = \cos x$



Find the amplitude of the function and use the language of transformations to describe how the graph of the function is related to the graph of $y = \sin x$

A) $y = 3\sin x$

B) $y = \frac{3}{4}\sin x$

C) $y = -5\sin x$

Find the period of the function and use the language of transformations to describe how the graph of the function is related to the graph of $y = \cos x$

A) $y = \cos(2x)$

B) $y = \cos \frac{x}{2}$

C) $y = \cos\left(\frac{-3x}{4}\right)$

Graph 1 period of the function without using your calculator.

A) $y = 3\sin \frac{x}{2}$

$y = 5\cos 2x$

Identify the maximum and minimum values and the zeros of the function in the interval $[-2\pi, 2\pi]$. Use your understanding of transformations, not your calculator.

A) $y = 4\sin x$

B) $y = -2\cos \frac{x}{3}$

Determine the phase shift for the function and the sketch the graph.

A) $y = \cos\left(x - \frac{\pi}{6}\right)$

B) $y = \sin\left(x + \frac{\pi}{3}\right)$

Determine the vertical shift for the function and the sketch the graph.

A) $y = \cos x - 2$

B) $y = \sin x + 3$

Determine the vertical shift and phase shift of the function and then sketch the graph

A) $y = \cos\left(x + \frac{\pi}{6}\right) - 1$

B) $y = \sin\left(x - \frac{\pi}{3}\right) + 2$

State the Amplitude and period of the sinusoid, and relative to the basic function, the phase shift and vertical translation.

A) $y = 3\sin\left(x - \frac{\pi}{4}\right) + 2$

B) $y = -2\cos\left(3x - \frac{\pi}{4}\right) - 4$

C) $y = 5\sin 4\pi x + 6$

$$Amp = A = \frac{Max - Min}{2}$$

$$Vertical = (C) = \frac{Max + Min}{2}$$

$$period = p$$

Horizontal Stretch/Shrink

$$B = \frac{2\pi}{p}$$

How to choose an appropriate model based on the behavior at some given time, T .

$y = A \cos B(t - T) + C$
if at time T the function
attains a maximum value

$y = -A \cos B(t - T) + C$
if at time T the function
attains a minimum value

$y = A \sin B(t - T) + C$
if at time T the function
halfway between a
minimum and a maximum
value

$y = -A \sin B(t - T) + C$
if at time T the function
halfway between a
maximum and a minimum
value

Construct a sinusoid with the given amplitude and period that goes through the given point.

A) Amp: 4, period 4π , point $(0, 0)$

B) Amp: 2.5, period $\frac{\pi}{5}$, point (2, 0)

$$\text{Amp} = A = \frac{\text{Max} - \text{Min}}{2}$$

$$\text{Vertical} = (C) = \frac{\text{Max} + \text{Min}}{2}$$

$$\text{period} = p$$

Horizontal Stretch/Shrink

$$B = \frac{2\pi}{p}$$

How to choose an appropriate model based on the behavior at some given time, T.

$y = A \cos B(t - T) + C$
if at time T the function attains a maximum value

$y = -A \cos B(t - T) + C$
if at time T the function attains a minimum value

$y = A \sin B(t - T) + C$
if at time T the function halfway between a minimum and a maximum value

$y = -A \sin B(t - T) + C$
if at time T the function halfway between a maximum and a minimum value

Example 7: Calculating the Ebb and Flow of Tides

One particular July 4th in Galveston, TX, high tide occurred at 9:36 am. At that time the water at the end of the 61st Street Pier was 2.7 meters deep. Low tide occurred at 3:48 p.m, at which time the water was only 2.1 meters deep. Assume that the depth of the water is a sinusoidal function of time with a period of half a lunar day (about 12 hrs 24 min)

a) Model the depth, D, as a sinusoidal function of time, t, algebraically then graph the function.

b) At what time on the 4th of July did the first low tide occur.

c) What was the approximate depth of the water at 6:00 am and at 3:00 pm?

d) What was the first time on July 4th when the water was 2.4 meters deep?

80) Temperature Data: The normal monthly Fahrenheit temperatures in Helena, MT, are shown in the table below (month 1 = January)

Model the temperature T as a sinusoidal function of time using 20 as the minimum value and 68 as the maximum value. Support your answer graphically by graphing your function with a scatter plot.


M	1	2	3	4	5	6	7	8	9	10	11	12
T	20	26	35	44	53	61	68	67	56	45	31	21

--	--


What you'll Learn About

- The graphs of the other 4 trig functions


The graph of $y = \csc x$




The graph of $y = \sec x$



The graph of $y = \tan x$



The graph of $y = \cot x$



Describe the graph of the function in terms of a basic trigonometric function. Locate the vertical asymptotes and graph 2 periods of the function.

A) $y = 2\tan(3x)$

B) $y = -\cot(2x)$

C) $y = \sec(4x)$

D) $y = -\csc\left(\frac{x}{3}\right)$

Describe the transformations required to obtain the graph of the given function from a basic trigonometric graph.

A) $y = 5 \tan x$

B) $y = -3 \cot\left(\frac{x}{2}\right)$

C) $y = 2 \sec \frac{4x}{3}$


D) $y = -4 \csc 2\pi x - 3$

--	--


What you'll Learn About

- Inverse Trigonometric Functions and their Graphs


The graph of $y = \sin x$




The graph of $y = \sin^{-1} x = \arcsin x$




The Unit Circle and Inverse Functions




The graph of $y = \cos x$




The graph of $y = \cos^{-1} x = \arccos x$





The Unit Circle and Inverse Functions



The graph of $y = \tan x$



The graph of $y = \tan^{-1} x = \arctan x$



Find the exact value

A) $\cos^{-1} \frac{\sqrt{3}}{2}$

B) $\cos^{-1} \frac{1}{2}$

C) $\cos^{-1} \left(\frac{-1}{2} \right)$

D) $\sin^{-1} \frac{-\sqrt{3}}{2}$

E) $\sin^{-1} \frac{1}{2}$

F) $\sin^{-1} \left(\frac{1}{\sqrt{2}} \right)$

G) $\tan^{-1}(1)$

H) $\tan^{-1}(\sqrt{3})$

I) $\tan^{-1} \left(\frac{-1}{\sqrt{3}} \right)$

J) $\cos^{-1}(0)$

K) $\sin^{-1}(-1)$

L) $\tan^{-1}(0)$

Use a calculator to find the approximate value in degrees. Draw the triangle that represents the situation.

A) $\arccos(.456)$

B) $\arcsin(-.456)$

C) $\arctan(-5.768)$

Use a calculator to find the approximate value in radians. Draw the triangle that represents the situation.

A) $\arcsin(.456)$

B) $\arccos(-.456)$

C) $\arctan(-5.768)$

Find the exact value without a calculator.

A) $\sin(\cos^{-1}(1/2))$

B) $\cos(\tan^{-1}(0))$

C) $\tan\left(\sin^{-1}\left(\frac{\sqrt{2}}{2}\right)\right)$

D) $\sin(\tan^{-1}(-\sqrt{3}))$

E) $\cos^{-1}\left(\sin\left(\frac{\pi}{4}\right)\right)$

F) $\sin^{-1}\left(\cos\left(\frac{\pi}{6}\right)\right)$

--	--

PRE-CALCULUS: by Finney, Demana, Watts and Kennedy
Solving Trigonometric Equations

What you'll Learn About

Solve each trigonometric equation for θ on the interval $[0, 2\pi]$. Then give a formula for all possible angles that could be a solution of the equation.

A) $\sin \theta = \frac{\sqrt{2}}{2}$

B) $\cos \theta = \frac{-1}{2}$

C) $\sin \theta = 1$

D) $\cos \theta = 0$

E) $\tan \theta = \sqrt{3}$

F) $\tan \theta = -1$

Solve each trigonometric equation for θ on the interval $[0, 2\pi]$.

A) $\cos 2\theta = \frac{1}{2}$

B) $\sin 3\theta = \frac{1}{2}$

C) $\cos \frac{\theta}{3} = \frac{\sqrt{3}}{2}$

D) $\tan\left(\frac{\theta}{2} + \frac{\pi}{3}\right) = 1$

E) $\sin \theta = .4$

F) $\cos \theta = -.2$

A) $\sqrt{2} \cos \theta - 1 = 0$

B) $\sqrt{3} \csc \theta - 2 = 0$

C) $4 \sin^2 \theta - 1 = 0$

D) $(3 \cot^2 \theta - 1)(\cot^2 \theta - 3) = 0$

E) $3 \tan^2 \theta - 1 = 0$

F) $\cos^2 \theta = 3 \sin^2 \theta$

$$\text{G) } 2\cos^2\theta + \cos\theta = 0$$

$$\text{H) } 2\sin\theta\cos\theta = \cos\theta$$

$$\text{I) } \csc^2\theta - \csc\theta = 2$$

$$\text{J) } \sin^3\theta = \sin\theta$$

--	--

Reciprocal Identities

$$\begin{aligned} \csc x &= \frac{1}{\sin x} & \sec x &= \frac{1}{\cos x} & \cot x &= \frac{1}{\tan x} \\ \sin x &= \frac{1}{\csc x} & \cos x &= \frac{1}{\sec x} & \tan x &= \frac{1}{\cot x} \end{aligned}$$

Quotient Identities

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x}$$

Even/Odd Identities

$$\begin{aligned} \sin(-x) &= -\sin x & \csc(-x) &= -\csc x \\ \cos(-x) &= \cos x & \sec(-x) &= \sec x \\ \tan(-x) &= -\tan x & \cot(-x) &= -\cot x \end{aligned}$$

Pythagorean Identities

$$\begin{aligned} \sin^2 x + \cos^2 x &= 1 & \tan^2 x + 1 &= \sec^2 x & 1 + \cot^2 x &= \csc^2 x \\ \sin^2 x &= 1 - \cos^2 x & \tan^2 x &= \sec^2 x - 1 & \csc^2 x - \cot^2 x &= 1 \\ \cos^2 x &= 1 - \sin^2 x & \sec^2 x - \tan^2 x &= 1 & \cot^2 x &= \csc^2 x - 1 \end{aligned}$$

Co-function

$$\begin{aligned} \sin\left(\frac{\pi}{2} - x\right) &= \cos x & \cos\left(\frac{\pi}{2} - x\right) &= \sin x & \tan\left(\frac{\pi}{2} - x\right) &= \cot x \\ \csc\left(\frac{\pi}{2} - x\right) &= \sec x & \sec\left(\frac{\pi}{2} - x\right) &= \csc x & \cot\left(\frac{\pi}{2} - x\right) &= \tan x \end{aligned}$$

Equation of Unit Circle $x^2 + y^2 = 1$	Use trig ratios to prove that $\cos^2 \theta + \sin^2 \theta = 1$ is the equation of the unit circle Use basic identities to simplify the expression to a different trig function or a product of two trig functions 10. $\cot x \tan x$ B. $\sin x - \sin^3 x$ A. $\frac{1 - \sin^2 \theta}{\cos \theta}$ C. $\frac{\sin^2 x + \cot^2 x + \cos^2 x}{\csc x}$
---	--

Simplify the expression to either 1 or -1

17. $\sin x \csc(-x)$

19. $\cot(-x)\cot\left(\frac{\pi}{2}-x\right)$

21. $\sin^2(-x) + \cos^2(-x)$

Simplify the expression to either a constant or a basic trig function.

A) $\frac{\cot\left(\frac{\pi}{2}-x\right)\sec x}{\sec^2 x}$

B) $\frac{1+\cot x}{1+\tan x}$

C) $\tan^2 x + \cot^2 x - (\sec^2 x + \csc^2 x)$

Use the basic identities to change the expression to one involving only sines and cosines. Then simplify to a basic trig function.

28) $\sin \theta - \tan \theta \cos \theta + \cos \left(\frac{\pi}{2} - \theta \right)$

30) $\frac{(\sec y - \tan y)(\sec y + \tan y)}{\sec y}$

31. $\frac{\tan x}{\csc^2 x} + \frac{\tan x}{\sec^2 x}$

Combine the fractions and simplify to a multiple of a power of a basic trig function

A) $\frac{1}{1 - \cos x} + \frac{1}{1 + \cos x}$

35. $\frac{\sin x}{\cot^2 x} - \frac{\sin x}{\cos^2 x}$

Write each expression in factored form as an algebraic expression of a single trig function

A) $\sin^2 x + 2 \sin x + 1$

B) $1 - 2 \cos x + \cos^2 x$

C) $\sin x - 2 \cos^2 x + 1$

45. $4 \tan^2 x - \frac{4}{\cot x} + \sin x \csc x$

Write each expression as an algebraic expression of a single trigonometric function

A) $\frac{1 - \cos^2 x}{1 + \cos x}$

B) $\frac{\cot^2 \alpha - 1}{1 + \cot x}$

C) $\frac{\cos^2 x}{1 + \sin x}$

D) $\frac{\cot^2 x}{1 + \csc x}$

What you'll Learn About

$$12. \sin x (\cot x + \cos x \tan x) = \cos x + \sin^2 x$$

$$14. (\cos x - \sin x)^2 = 1 - 2 \sin x \cos x$$

$$18. \frac{\sec^2 \theta - 1}{\sin \theta} = \frac{\sin \theta}{1 - \sin^2 \theta}$$

$$20. \frac{1}{1 - \cos x} + \frac{1}{1 + \cos x} = 2 \csc^2 x$$

$$22. \sin^2 \alpha - \cos^2 \alpha = 1 - 2\cos^2 \alpha$$

$$26. \frac{\sec x + 1}{\tan x} = \frac{\sin x}{1 - \cos x}$$

$$30. \quad \tan^2 \theta - \sin^2 \theta = \tan^2 \theta \sin^2 \theta$$

$$35. \quad \frac{\tan x}{\sec x - 1} = \frac{\sec x + 1}{\tan x}$$

$$37. \quad \frac{\sin x - \cos x}{\sin x + \cos x} = \frac{2 \sin^2 x - 1}{1 + 2 \sin x \cos x}$$

--	--

