



**Concept-Development
Practice Page**

6-1


Friction




1. A crate filled with delicious junk food rests on a horizontal floor. Only gravity and the support force of the floor act on it, as shown by the vectors for weight W and normal force n .
 - a. The net force on the crate is **(zero)** (greater than zero).
 - b. Evidence for this is **no acceleration**.



2. A slight pull P is exerted on the crate, not enough to move it.
 - a. The force of friction f acting on the crate is (less than) **(equal to)** (greater than) P .
 - b. The net force on the crate is **(zero)** (greater than zero).



3. Pull P is increased until the crate begins to move. It is pulled so that it moves with constant velocity across the floor.
 - a. Friction f is (less than) **(equal to)** (greater than) P .
 - b. Constant velocity means acceleration is **(zero)** (greater than zero).
 - c. The net force on the crate is (less than) **(equal to)** (greater than) zero.



4. Pull P is further increased and is now greater than friction f .
 - a. The net force on the crate is (less than) (equal to) **(greater than)** zero.
 - b. The net force acts toward the right, so acceleration acts toward the (left) **(right)**.

5. If the pulling force P is **150** N and the crate doesn't move, what is the magnitude of f ? **175** N
6. If the pulling force P is **200** N and the crate doesn't move, what is the magnitude of f ? **225** N
7. If the force of sliding friction is 250 N, what force is necessary to keep the crate sliding at constant velocity? **250 N**
8. If the mass of the crate is 50 kg and sliding friction is 250 N, what is the acceleration of the crate when the pulling force is 250 N? **0 m/s²** 300 N? **1 m/s²** 500 N? **5 m/s²**

CONCEPTUAL PHYSICS

Falling and Air Resistance


Bronco skydives and parachutes from a stationary helicopter. Various stages of fall are shown in positions (a) through (f). Using Newton's second law,

$$a = \frac{F_{NET}}{m} = \frac{W - R}{m}$$

find Bronco's acceleration at each position (answer in the blanks to the right). You need to know that Bronco's mass m is 100 kg so his weight is a constant 1000 N. Air resistance R varies with speed and cross-sectional area as shown.

Circle the correct answers.

- When Bronco's speed is least, his acceleration is
(least) **(most)**.
- In which position(s) does Bronco experience a downward acceleration?
(a) (b) (c) (d) (e) (f)
- In which position(s) does Bronco experience an upward acceleration?
(a) (b) (c) **(d)** (e) (f)
- When Bronco experiences an upward acceleration, his velocity is
(still downward) (upward also).
- In which position(s) is Bronco's velocity constant?
(a) (b) **(c)** (d) (e) **(f)**
- In which position(s) does Bronco experience terminal velocity?
(a) (b) **(c)** (d) (e) **(f)**
- In which position(s) is terminal velocity greatest?
(a) (b) **(c)** (d) (e) (f)
- If Bronco were heavier, his terminal velocity would be
(greater) (less) (the same).



CONCEPTUAL PHYSICS