

### Question 1

$t$ (minutes)	0	4	9	15	20
$W(t)$ (degrees Fahrenheit)	55.0	57.1	61.8	67.9	71.0

The temperature of water in a tub at time  $t$  is modeled by a strictly increasing, twice-differentiable function  $W$ , where  $W(t)$  is measured in degrees Fahrenheit and  $t$  is measured in minutes. At time  $t = 0$ , the temperature of the water is  $55^\circ\text{F}$ . The water is heated for 30 minutes, beginning at time  $t = 0$ . Values of  $W(t)$  at selected times  $t$  for the first 20 minutes are given in the table above.

- Use the data in the table to estimate  $W'(12)$ . Show the computations that lead to your answer. Using correct units, interpret the meaning of your answer in the context of this problem.
- Use the data in the table to evaluate  $\int_0^{20} W'(t) dt$ . Using correct units, interpret the meaning of  $\int_0^{20} W'(t) dt$  in the context of this problem.
- For  $0 \leq t \leq 20$ , the average temperature of the water in the tub is  $\frac{1}{20} \int_0^{20} W(t) dt$ . Use a left Riemann sum with the four subintervals indicated by the data in the table to approximate  $\frac{1}{20} \int_0^{20} W(t) dt$ . Does this approximation overestimate or underestimate the average temperature of the water over these 20 minutes? Explain your reasoning.
- For  $20 \leq t \leq 25$ , the function  $W$  that models the water temperature has first derivative given by  $W'(t) = 0.4\sqrt{t} \cos(0.06t)$ . Based on the model, what is the temperature of the water at time  $t = 25$ ?

### Question 2


For  $t \geq 0$ , a particle is moving along a curve so that its position at time  $t$  is  $(x(t), y(t))$ . At time  $t = 2$ , the particle is at position  $(1, 5)$ . It is known that  $\frac{dx}{dt} = \frac{\sqrt{t+2}}{e^t}$  and  $\frac{dy}{dt} = \sin^2 t$ .

- Is the horizontal movement of the particle to the left or to the right at time  $t = 2$ ? Explain your answer. Find the slope of the path of the particle at time  $t = 2$ .
- Find the  $x$ -coordinate of the particle's position at time  $t = 4$ .
- Find the speed of the particle at time  $t = 4$ . Find the acceleration vector of the particle at time  $t = 4$ .
- Find the distance traveled by the particle from time  $t = 2$  to  $t = 4$ .

### Question 3

Let  $f$  be the continuous function defined on  $[-4, 3]$  whose graph, consisting of three line segments and a semicircle centered at the origin, is given above. Let  $g$  be the function given by  $g(x) = \int_1^x f(t) dt$ .

- Find the values of  $g(2)$  and  $g(-2)$ .
- For each of  $g'(-3)$  and  $g''(-3)$ , find the value or state that it does not exist.
- Find the  $x$ -coordinate of each point at which the graph of  $g$  has a horizontal tangent line. For each of these points, determine whether  $g$  has a relative minimum, relative maximum, or neither a minimum nor a maximum at the point. Justify your answers.
- For  $-4 < x < 3$ , find all values of  $x$  for which the graph of  $g$  has a point of inflection. Explain your



Graph of  $f$

### Question 4

$x$	1	1.1	1.2	1.3	1.4
$f'(x)$	8	10	12	13	14.5

The function  $f$  is twice differentiable for  $x > 0$ , with  $f(1) = 15$  and  $f''(1) = 20$ . Values of  $f'$ , the derivative of  $f$ , are given for selected values of  $x$  in the table above.

- Write an equation for the line tangent to the graph of  $f$  at  $x = 1$ . Use this line to approximate  $f(1.4)$ .
- Use a midpoint Riemann sum with two subintervals of equal length and values from the table to approximate  $\int_1^{1.4} f'(x) dx$ . Use the approximation for  $\int_1^{1.4} f'(x) dx$  to estimate the value of  $f(1.4)$ . Show the computations that lead to your answer.
- Use Euler's method, starting at  $x = 1$  with two steps of equal size, to approximate  $f(1.4)$ . Show the computations that lead to your answer.
- Write the second-degree Taylor polynomial for  $f$  about  $x = 1$ . Use the Taylor polynomial to approximate  $f(1.4)$ .


### Question 5

The rate at which a baby bird gains weight is proportional to the difference between its adult weight and its current weight. At time  $t = 0$ , when the bird is first weighed, its weight is 20 grams. If  $B(t)$  is the weight of the bird, in grams, at time  $t$  days after it is first weighed, then

$$\frac{dB}{dt} = \frac{1}{5}(100 - B).$$

Let  $y = B(t)$  be the solution to the differential equation above with initial condition  $B(0) = 20$ .

- Is the bird gaining weight faster when it weighs 40 grams or when it weighs 70 grams? Explain your reasoning.
- Find  $\frac{d^2B}{dt^2}$  in terms of  $B$ . Use  $\frac{d^2B}{dt^2}$  to explain why the graph of  $B$  cannot resemble the following graph.
- Use separation of variables to find  $y = B(t)$ , the particular solution to the differential equation with initial condition  $B(0) = 20$ .



### Question 6

The function  $g$  has derivatives of all orders, and the Maclaurin series for  $g$  is

$$\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+3} = \frac{x}{3} - \frac{x^3}{5} + \frac{x^5}{7} - \dots$$

- Using the ratio test, determine the interval of convergence of the Maclaurin series for  $g$ .
- The Maclaurin series for  $g$  evaluated at  $x = \frac{1}{2}$  is an alternating series whose terms decrease in absolute value to 0. The approximation for  $g\left(\frac{1}{2}\right)$  using the first two nonzero terms of this series is  $\frac{17}{120}$ . Show that this approximation differs from  $g\left(\frac{1}{2}\right)$  by less than  $\frac{1}{200}$ .
- Write the first three nonzero terms and the general term of the Maclaurin series for  $g'(x)$ .