

yes
 Since $C(t)$ is differentiable from $(3,6)$ the MVT guarantees that the Avg. Rate of Change will equal the Instantaneous Rate of Change

2013 BC3

Hot water is dripping through a coffeemaker, filling a large cup with coffee. The amount of coffee in the cup at time t , $0 \leq t \leq 6$, is given by a differentiable function C , where t is measured in minutes. Selected values of $C(t)$, measured in ounces, are given in the table.

t (minute s)	0	1	2	3	4	5	6
$C(t)$ ounces	0	5.3	8.8	11.2	12.8	13.8	14.2


Is there a time t , $3 \leq t \leq 6$, at which $C'(t) = 1$. Justify your answer.

Avg Rate of Change

$$\frac{14.2 - 11.2}{6 - 3} = 1$$

Instantaneous rate of change is 1
 Slope of tangent line is 1

Let g be a continuous function with $g(2) = 5$. The graph of the piecewise-linear function g' , the derivative of g , is shown for $-3 \leq x \leq 7$.



Find the average rate of change of $g(x)$, on the interval $-3 \leq x \leq 1$. Does the Mean Value Theorem applied on the interval $-3 \leq x \leq 1$ guarantee a value of c , for $-3 < c < 1$, such that $g'(c)$ is equal to this average rate of change? Why or why not?


$(-3, -4)$ $(1, 1)$

A.R.O.C. of $g(x) = \frac{-4 - 1}{-3 - 1} = \frac{-5}{-4} = \frac{5}{4}$

No, because $g'(x)$ is not differentiable from $(-3, 1)$
 $-3 < x < 1$

2005 AB5

A car is traveling on a straight road. For $0 \leq t \leq 24$ seconds, the car's velocity $v(t)$, in meters per second, is modeled by the piecewise-linear function defined by the graph



$(0,0)$ $(16,20)$

$$\frac{20-0}{16-0} = \frac{20}{16} = \frac{5}{4}$$

- i) Find the average rate of change of v over the interval $0 \leq t \leq 16$. Does the Mean Value guarantee a value of c , for $0 < c < 16$, such that $v'(t)$ is equal to this average rate of change? Why or why not?

No. B/c $v(t)$ is not differential from $0 < c < 16$

2004 BCB3

A test plane flies in a straight line with positive velocity $v(t)$, in miles per minute at time t minutes, where v is a differentiable function of t . Selected values of $v(t)$ are shown.

t (min)	0	5	10	15	20	25	30	35	40
$v(t)$ (mpm)	7	9.2	9.5	9.2	4.5	2.4	4.5	4.9	7.3

Based on the values in the table, what is the smallest number of instances at which the acceleration of the plane could equal zero on the open interval $0 < t < 40$? Justify your answer


Two. Since $v(t)$ is differentiable $0 < t < 40$ the MVT guarantees the avg. acceleration equals the instantaneous acceleration

$$\frac{9.2-9.2}{15-5} = 0$$

$$\frac{4.5-4.5}{30-20} = 0$$

2009 BC3

A continuous function f is defined on the closed interval $-4 \leq x \leq 6$. The graph of f consists of a line segment and a curve that is tangent to the x -axis at $x = 3$, as shown in the figure above. On the interval $0 < x < 6$, the function f is twice differentiable, with $f'(x) > 0$.



$(0, 2)$ $(6, 1)$

$$\frac{2-1}{0-6} = -\frac{1}{6}$$

Is there a value a , for which the Mean Value Theorem, applied to the interval $[a, 6]$, guarantees a value c , $a < c < 6$, at which $f'(c) = -\frac{1}{6}$? Justify your answer.

Yes. $a=0$. Since f is differentiable from $(0, 6)$ the MVT guarantees that $AROC = IROC$

2011 BCB5

Ben rides a unicycle back and forth along a straight east-west track. The twice-differentiable function B models Ben's position of the track, measured in meters from the western end of the track, at time t , measured in seconds from the start of the ride. The table gives values of $B(t)$ and Ben's velocity, $v(t)$, measured in meters per second, at selected times t .

t (seconds)	0	15	40	60
$B(t)$ (meters)	100	136	9	46
$V(t)$ meters per second	2	2.3	2.5	4.6

For $15 \leq t \leq 60$, must there be a time t when Ben's velocity is -2 meters per second? Justify your answer.


$$\frac{46-136}{60-15} = \frac{-90}{45} = -2 \text{ m/s}$$

yes. since $B(t)$ is differentiable from $0 < t < 60$
the MVT guarantees $AROC$ of $B(t) = IROC$ of $B(t)$

What you'll Learn About:
 How to use derivatives to find limits in an indeterminate form

Why L'Hopitals Works

Sketch the graph of two curves with the following characteristic $f(2) = g(2) = 0$.



a) Write the tangent line for $f(x)$

$$(2, 0) \quad f'(2)$$

$$y = 0 + f'(2)(x-2) \rightarrow f(x)$$

b) Write the tangent line for $g(x)$

$$(2, 0) \quad g'(2)$$

$$y = 0 + g'(2)(x-2)$$

c) $\lim_{x \rightarrow 2} \frac{f(x)}{g(x)} = \frac{f(2)}{g(2)} = \frac{0}{0}$

$$= \frac{0 + f'(2)(x-2)}{0 + g'(2)(x-2)} = \frac{f'(2)}{g'(2)} = \lim_{x \rightarrow 2} \frac{f(x)}{g(x)}$$

d) $\lim_{x \rightarrow 0} \frac{2x^2}{x^2} = 2$

$$\lim_{x \rightarrow 0} \frac{2x^2}{x^2} = \frac{0}{0}$$

$$\lim_{x \rightarrow 0} \frac{4x}{2x} = \frac{0}{0}$$

$$\lim_{x \rightarrow 0} \frac{4}{2} = 2$$

2) $\lim_{x \rightarrow 0} \frac{\sin(5x)}{x} = \frac{0}{0}$

$$\lim_{x \rightarrow 0} \frac{\cos(5x) \cdot 5}{1} = 5$$

Indeterminate Form

$x^{1/3}$

$$4) \lim_{x \rightarrow 1} \frac{\sqrt[3]{x} - 1}{x - 1} = \frac{0}{0}$$

$$\lim_{x \rightarrow 1} \frac{\frac{1}{3}x^{-2/3}}{1}$$

$$\lim_{x \rightarrow 1} \frac{1}{3x^{2/3}} = \frac{1}{3}$$

$$A) \lim_{x \rightarrow \infty} \frac{x^3 - 1}{4x^3 - x - 3}$$

$$35) \lim_{x \rightarrow \infty} \frac{\log_2(x)}{\log_3(x+3)}$$

$$49) \lim_{x \rightarrow 1} \frac{x^3 - 1}{4x^3 - x - 3} = \frac{0}{0}$$

$$\lim_{x \rightarrow 1} \frac{3x^2}{12x^2 - 1} = \frac{3}{11}$$

$$27) \lim_{x \rightarrow \infty} \frac{\ln(x^5)}{x}$$

$$33) \lim_{x \rightarrow 0} \frac{\sin(x^2)}{x} = \frac{0}{0}$$

$$\lim_{x \rightarrow 0} \frac{\cos(x^2) \cdot 2x}{1} = \frac{0}{1}$$

$$\lim_{x \rightarrow 0} \frac{\sin(x^2)}{x} = 0$$

