

Copyright © Cengage Learning. All rights reserved.

What You Should Learn

- Find the dot product of two vectors and use the properties of the dot product.
- Find the angle between two vectors and determine whether two vectors are orthogonal.
- Write vectors as the sums of two vector components.



The Dot Product of Two Vectors



So far you have studied two vector operations—vector addition and multiplication by a scalar—each of which yields another vector.

In this section, you will study a third vector operation, the **dot product.** This product yields a scalar, rather than a vector.

The Dot Product of Two Vectors

Definition of Dot Product

The **dot product** of $\mathbf{u} = \langle u_1, u_2 \rangle$ and $\mathbf{v} = \langle v_1, v_2 \rangle$ is given by

 $\mathbf{u}\cdot\mathbf{v}=u_1v_1+u_2v_2.$

Properties of the Dot Product

Let **u**, **v**, and **w** be vectors in the plane or in space and let *c* be a scalar.

1. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$ 2. $\mathbf{0} \cdot \mathbf{v} = \mathbf{0}$ 3. $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$ 4. $\mathbf{v} \cdot \mathbf{v} = \|\mathbf{v}\|^2$ 5. $c(\mathbf{u} \cdot \mathbf{v}) = c\mathbf{u} \cdot \mathbf{v} = \mathbf{u} \cdot c\mathbf{v}$

Let $\mathbf{u} = \langle -1, 3 \rangle$, $\mathbf{v} = \langle 2, -4 \rangle$, and $\mathbf{w} = \langle 1, -2 \rangle$. Use the vectors and the properties of the dot product to find the indicated quantity.

a. $(\mathbf{u} \cdot \mathbf{v})\mathbf{w}$ b. $\mathbf{u} \cdot 2\mathbf{v}$

Solution:

Begin by finding the dot product of **u** and **v**.

$$\mathbf{u} \cdot \mathbf{v} = \langle -1, 3 \rangle \cdot \langle 2, -4 \rangle$$

= (-1)(2) + 3(-4)
= -14



a.
$$(\mathbf{u} \cdot \mathbf{v})\mathbf{w} = -14\langle 1, -2 \rangle$$

= $\langle -14, 28 \rangle$
b. $\mathbf{u} \cdot 2\mathbf{v} = 2(\mathbf{u} \cdot \mathbf{v})$
= $2(-14)$
= -28

cont'd



The Angle Between Two Vectors



The **angle between two nonzero vectors** is the angle θ , $0 \le \theta \le \pi$, between their respective standard position vectors, as shown in Figure 6.36.



Figure 6.36

This angle can be found using the dot product. (Note that the angle between the zero vector and another vector is not defined.)

The Angle Between Two Vectors



Example 3 – Finding the Angle Between Two Vectors

Find the angle between $\mathbf{u} = \langle 4, 3 \rangle$ and $\mathbf{v} = \langle 3, 5 \rangle$.

Solution:

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}$$
$$= \frac{\langle 4, 3 \rangle \cdot \langle 3, 5 \rangle}{\|\langle 4, 3 \rangle\| \|\langle 3, 5 \rangle\|}$$
$$= \frac{27}{5\sqrt{34}}$$

This implies that the angle between the two vectors is

$$\theta = \arccos \frac{27}{5\sqrt{34}} \approx 22.2^{\circ}$$

as shown in Figure 6.37.



Figure 6.37

cont'd

The Angle Between Two Vectors

Definition of Orthogonal Vectors

The vectors **u** and **v** are **orthogonal** when $\mathbf{u} \cdot \mathbf{v} = 0$.

The terms *orthogonal* and *perpendicular* mean essentially the same thing—meeting at right angles.

Even though the angle between the zero vector and another vector is not defined, it is convenient to extend the definition of orthogonality to include the zero vector.

In other words, the zero vector is orthogonal to every vector **u** because $\mathbf{0} \cdot \mathbf{u} = 0$.

Example 4 – Determining Orthogonal Vectors

Are the vectors

$$\mathbf{u} = \langle 2, -3 \rangle$$
 and $\mathbf{v} = \langle 6, 4 \rangle$

orthogonal?

Solution:

Begin by finding the dot product of the two vectors.

$$\mathbf{u} \cdot \mathbf{v} = \langle 2, -3 \rangle \cdot \langle 6, 4 \rangle$$
$$= 2(6) + (-3)(4)$$
$$= 0$$

Because the dot product is 0, the two vectors are orthogonal, as shown in Figure 6.39.





cont'd



You have already seen applications in which two vectors are added to produce a resultant vector. Many applications in physics and engineering pose the reverse problem decomposing a given vector into the sum of two vector components.

Consider a boat on an inclined ramp, as shown in Figure 6.40.

The force **F** due to gravity pulls the boat *down* the ramp and *against* the ramp.



These two orthogonal forces, w_1 and w_2 , are vector components of **F**. That is,

 $F = w_1 + w_2$.

Vector components of ${\bf F}$

The negative of component \mathbf{w}_1 represents the force needed to keep the boat from rolling down the ramp, and \mathbf{w}_2 represents the force that the tires must withstand against the ramp.

A procedure for finding w1 and w2 is shown below.

```
Definition of Vector Components
```

Let **u** and **v** be nonzero vectors such that

 $\mathbf{u} = \mathbf{w}_1 + \mathbf{w}_2$

where \mathbf{w}_1 and \mathbf{w}_2 are orthogonal and \mathbf{w}_1 is parallel to (or a scalar multiple of) \mathbf{v} , as shown in Figure 6.41. The vectors \mathbf{w}_1 and \mathbf{w}_2 are called vector components of \mathbf{u} . The vector \mathbf{w}_1 is the **projection** of \mathbf{u} onto \mathbf{v} and is denoted by

 $\mathbf{w}_1 = \operatorname{proj}_{\mathbf{v}} \mathbf{u}.$

The vector \mathbf{w}_2 is given by

 $\mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1.$







Figure 6.41

From the definition of vector components, you can see that it is easy to find the component w_2 once you have found the projection of **u** onto **v**. To find the projection, you can use the dot product, as follows.

 $\mathbf{u} = \mathbf{w}_{1} + \mathbf{w}_{2}$ $\mathbf{u} = c\mathbf{v} + \mathbf{w}_{2}$ $\mathbf{u} \cdot \mathbf{v} = (c\mathbf{v} + \mathbf{w}_{2}) \cdot \mathbf{v}$ $\mathbf{u} \cdot \mathbf{v} = c\mathbf{v} \cdot \mathbf{v} + \mathbf{w}_{2} \cdot \mathbf{v}$ $\mathbf{u} \cdot \mathbf{v} = c\mathbf{v} \cdot \mathbf{v} + \mathbf{w}_{2} \cdot \mathbf{v}$ $\mathbf{w}_{2} \text{ and } \mathbf{v} \text{ are orthogonal.}$ $\mathbf{u} \cdot \mathbf{v} = c||\mathbf{v}||^{2} + 0$

So,
$$c = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}$$

and

$$\mathbf{w}_1 = \operatorname{proj}_{\mathbf{v}} \mathbf{u} = c\mathbf{v} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\mathbf{v}.$$

Projection of u onto v

Let **u** and **v** be nonzero vectors. The projection of **u** onto **v** is given by

$$\operatorname{proj}_{\mathbf{v}} \mathbf{u} = \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right) \mathbf{v}$$

Example 5 – Decomposing a Vector into Components

Find the projection of

 $\mathbf{u} = \langle 3, -5 \rangle$ onto $\mathbf{v} = \langle 6, 2 \rangle$.

Then write u as the sum of two orthogonal vectors, one of which is $\mathrm{proj}_v u$.

Example 5 – Solution

The projection of onto is

$$\mathbf{w}_1 = \operatorname{proj}_{\mathbf{v}} \mathbf{u} = \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right) \mathbf{v} = \left(\frac{8}{40}\right) \langle 6, 2 \rangle = \left\langle \frac{6}{5}, \frac{2}{5} \right\rangle$$

as shown in Figure 6.42.



Example 5 – Solution

The other component, \mathbf{w}_2 , is

$$\mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 3, -5 \rangle - \left\langle \frac{6}{5}, \frac{2}{5} \right\rangle = \left\langle \frac{9}{5}, -\frac{27}{5} \right\rangle.$$

So,

 $\mathbf{u} = \mathbf{w}_1 + \mathbf{w}_2$ $= \left\langle \frac{6}{5}, \frac{2}{5} \right\rangle + \left\langle \frac{9}{5}, -\frac{27}{5} \right\rangle$ $= \langle 3, -5 \rangle$

cont'd