AP B Physics	Name	Per	
Summer Assignment	Date		
We have read the AP Student contract	t and submitted a signed c	opy to an AP Counselor	
We have read the policies and expecta We understand and accept these polic	ations for AP Physics. cies.		
Student Signature:		Date	
Parent / Guardian Name (print)			
Parent / Guardian Signature:		Date	

- I. As is evident in the AP Physics Syllabus, we must cover a large number of topics before the test in May. This necessitates a very fast pace. This summer homework will allow us to start on the Physics subject matter immediately when school begins. This assignment is an introduction to Chapter 2 in the textbook and a math review to brush up on valuable skills, and perhaps a means to assess whether you are correctly placed in Advanced Placement Physics.
- II. Physics, and AP Physics in particular, requires an exceptional proficiency in algebra, trigonometry, and geometry. In addition to the science concepts Physics often seems like a course in applied mathematics. The following assignment includes mathematical problems that are considered routine in AP Physics. This includes knowing several key metric system conversion factors and how to employ them.
- III. The attached pages contain a brief review, hints, and example problems. It is hoped that combined with your previous math knowledge this assignment is merely a review and a means to brush up before school begins in the fall. Please read the text and instructions throughout.

IV. What is due the first day of school?

- A. Signed Class Expectations Sheet
 - 1. **Read the above statements.**
 - 2. Complete the section at the top of this form and obtain appropriate signatures.
- B. Problems 1 to 5 of the Math Skills Worksheet (next 3 pages of this packet)
- C. AP Physics Textbook Assignment

Check out the Physics Text (by Giancoli) at registration.

- Read & Outline Chapter 2, sections 2-1 to 2-6.
- Make sure to include the Problem Solving section on page 28.
- At the back of each chapter is a set of Questions followed by a set of Problems. Completely answer Questions # 1 11 on page 38 and do Problems # 1 14 on page 39 at the back of chapter 2. Be sure to show ALL your work when completing the Problems. Also be sure to include units with ALL numerical values that have units.

Math Skills Worksheet

 The following are ordinary physics problems. Place the answer in scientific notation when appropriate and simplify the units (Scientific notation is used when it takes less time to write than the ordinary number does. As an example 200 is easier to write than 2.00x10², but 2.00x10⁸ is easier to write than 200,000,000). Do your best to cancel units, and attempt to show the simplified units in the final answer.

a.
$$T_{s} = 2\pi \sqrt{\frac{4.5 \times 10^{-2} kg}{2.0 \times 10^{3} kg/s^{2}}} =$$
b.
$$F = \left(9.0 \times 10^{9} \frac{N \cdot m^{2}}{C^{2}}\right) \frac{(3.2 \times 10^{-9} C)(9.6 \times 10^{-9} C)}{(0.32m)^{2}} =$$
c.
$$\frac{1}{R_{p}} = \frac{1}{4.5 \times 10^{2} \Omega} + \frac{1}{9.4 \times 10^{2} \Omega} \qquad R_{p} =$$
d.
$$K_{max} = \left(6.63 \times 10^{-34} J \cdot s\right) \left(7.09 \times 10^{14} s\right) - 2.17 \times 10^{-19} J =$$
e.
$$\gamma = \sqrt{\frac{1}{\sqrt{1 - \frac{2.25 \times 10^{8} m/s}{3.00 \times 10^{8} m/s}}} =$$

2. Often problems on the AP exam are done with variables only. Solve for the variable indicated. Don't let the different letters confuse you. Manipulate them algebraically as though they were numbers.

a.
$$K = \frac{1}{2}kx^2$$
 , $x =$
 f. $B = \frac{\mu_o}{2\pi} \frac{I}{r}$
 , $r =$

 ...
 ...
 g. $x_m = \frac{m\lambda L}{d}$
 , $d =$

 ...
 ...
 g. $x_m = \frac{m\lambda L}{d}$
 , $d =$

 b. $T_p = 2\pi \sqrt{\frac{\ell}{g}}$
 , $g =$
 h. $pV = nRT$
 , $T =$

 ...
 ...
 i. $\sin \theta_c = \frac{n_1}{n_2}$
 , $\theta_c =$

 ...
 ...
 i. $\sin \theta_c = \frac{1}{n_2}$
 , $v =$

 ...
 ...
 ...
 k. $\frac{1}{f} = \frac{1}{2}mv^2$
 , $v =$

 ...
 ...
 ...
 ...
 ...

 e. $x = x_o + v_o t + \frac{1}{2}at^2$
 , $t =$
 ...
 ...

APB Physics, Summer Assignment

3. Science uses the *KMS* system (*SI*: System Internationale). *KMS* stands for kilogram, meter, second. These are the units of choice of physics. The equations in physics depend on unit agreement. So you must convert to *KMS* in most problems to arrive at the correct answer.

kilometers (<i>km</i>) to meters (<i>m</i>) centimeters (<i>cm</i>) to meters (<i>m</i>)	minutes (<i>min</i>) to seconds (<i>s</i>) hours (<i>hr</i>) to seconds (<i>s</i>)	gram (<i>g</i>) to kilogram (<i>kg</i>) Celsius (°C) to Kelvin (<i>K</i>)				
millimeters (<i>mm</i>) to meters (<i>m</i>) nanometers (<i>nm</i>) to meters (<i>m</i>)	days (<i>d</i>) to seconds (<i>s</i>) vears (<i>yr</i>) to seconds (<i>s</i>)	atmospheres (<i>atm</i>) to Pascals (<i>Pa</i>) liters (<i>L</i>) to cubic meters (m^3)				
micrometers (μm) to meters (m)						
Other conversions will be taught as they become necessary.						

What if you don't know the conversion factors? Colleges want students who can find their own information (so do employers). Hint: Try a good dictionary and look under "measure" or "measurement". Or the Internet? Enjoy.

a.	4008 g	=	_ kg	h.	25.0 <i>µm</i>	=	m
b.	1.2 <i>km</i>	=	_ m	i.	2.65 <i>mm</i>	=	m
c.	823 nm	=	_ m	j.	8.23 <i>m</i>	=	_ km
d.	298 <i>K</i>	=	°C	k.	5.4 <i>L</i>	=	_ <i>m</i> ³
e.	0.77 <i>m</i>	=	_ cm	I.	40.0 <i>cm</i>	=	m
f.	8.8x10 ⁻⁸ <i>m</i>	=	_ mm	m.	6.23x10 ⁻⁷ m	=	nm
g.	1.2 <i>atm</i>	=	Pa	n.	1.5x10 ¹¹ <i>m</i>	=	_ km

- 4. Solve the following geometric problems.
 - a. Line **B** touches the circle at a single point. Line **A** extends through the center of the circle.

С

θ

45°

θ

30°

30°

- i. What is line **B** in reference to the circle?
- ii. How large is the angle between lines **A** and **B**?
- b. What is angle C?



c. What is angle θ ?

How large is *θ*?

APB Physics, Summer Assignment

d.

- The radius of a circle is 5.5 cm, e.
 - What is the circumference in meters? i.
- What is its area in square meters? ii. 4 What is the area under the curve at the right? f. 2012 5. Using the generic triangle to the right, Right Triangle Trigonometry and Pythagorean Theorem solve the following. Your calculator must be in degree mode. С b θ а $\boldsymbol{\theta} = 55^{\circ}$ and $\boldsymbol{c} = 32 m$, solve for \boldsymbol{a} and \boldsymbol{b} . **a** = 250 *m* and **b** = 180 *m*, solve for **θ** and **c**. g. j.

 - $\boldsymbol{\theta} = 45^{\circ}$ and $\boldsymbol{a} = 15$ m/s, solve for **b** and **c**. h.
 - \boldsymbol{b} = 17.8 *m* and $\boldsymbol{\theta}$ = 65°, solve for \boldsymbol{a} and \boldsymbol{c} . i.
- k. $a = 25 \ cm$ and $c = 32 \ cm$, solve for b and θ .
- Ι. **b** =65 cm and **c** = 104 cm, solve for **a** and **θ**.