


AP Biology


Lab Review












Description

dialysis tubing filled with starchglucose solution in beaker filled with KI solution

potato cores in sucrose solutions




- Concepts
 - **♦**semi-permeable membrane
 - **♦** diffusion
 - **♦**osmosis
 - **♦**solutions
 - hypotonic
 - hypertonic
 - isotonic
 - water potential



Conclusions

water moves from high concentration of water (hypotonic=low solute) to low concentration of water (hypertonic=high solute)

Solute concentration & size of molecule affect movement through semi-permeable membrane



ESSAY 1992


A laboratory assistant prepared solutions of 0.8 M, 0.6 M, 0.4 M, and 0.2 M sucrose, but forgot to label them. After realizing the error, the assistant randomly labeled the flasks containing these four unknown solutions as flask A, flask B, flask C, and flask D.


Design an experiment, based on the principles of diffusion and osmosis, that the assistant could use to determine which of the flasks contains each of the four unknown solutions.


Include in your answer:

- a description of how you would set up and perform the experiment;
 - a. the results you would expect from your experiment; and
 - b.an explanation of those results based on the principles involved.

Be sure to clearly state the principles addressed in your discussion.









AP Biology 2004-2005


- Description
 - measured factors affecting enzyme activity
 - $Arr H_2O_2 \xrightarrow{\text{catalase}} H_2O + O_2$
 - **♦** measured rate of O₂ production




- Concepts
 - **♦**substrate
 - **♦**enzyme
 - enzyme structure
 - **♦**product
 - denaturation of protein
 - experimental design
 - rate of reactivity
 - reaction with enzyme vs. reaction without enzyme
 - optimum pH or temperature
 - test at various pH or temperature values



- Conclusions
 - enzyme reaction rate is affected by:
 - pH
 - temperature
 - substrate concentration
 - enzyme concentration




calculate rate?





ESSAY 2000

The effects of pH and temperature were studied for an enzyme-catalyzed reaction. The following results were obtained.




- a. How do (1) temperature and (2) pH affect the activity of this enzyme? In your answer, include a discussion of the relationship between the structure and the function of this enzyme, as well as a discussion of ho structure and function of enzymes are affected by temperature and pH.
- b. Describe a controlled experiment that could have produced the data shown for <u>either</u> temperature or pH. Be sure to state the hypothesis that was tested here.






- Description
 - cell stages of mitosis
 - exam slide of onion root tip
 - count number of cells in each stage to determine relative time spent in each stage
 - crossing over in meiosis
 - farther gene is from centromere the greater
 - number of crossovers
 - observed crossing over in fungus, *Sordaria* arrangement of ascospores







Fertilization

- Concepts
 - **♦**mitosis
 - interphase
 - prophase
 - metaphase
 - anaphase
 - telophase
 - **♦**meiosis
 - meiosis 1
 - meiosis 2
 - crossing over
 - tetrad in prophase 1






- Conclusions
 - **♦**Mitosis
 - longest phase = interphase
 - each subsequent phase is shorter in duration
 - **♦**Meiosis
 - 4:4 arrangement in ascospores
 - no crossover
 - any other arrangement
 - crossover
 - * 2:2:2:2 or 2:4:2




Sordaria analysis






B. Crossing over during meiosis







ESSAY 1987

Discuss the process of cell division in animals. Include a description of mitosis and cytokinesis, and of the other phases of the cell cycle. Do not include meiosis.


ESSAY 2004

Meiosis reduces chromosome number and rearranges genetic information.


- a. Explain how the reduction and rearrangement are accomplished in meiosis.
- b. Several human disorders occur as a result of defects in the meiotic process. Identify ONE such chromosomal abnormality; what effects does it have on the phenotype of people with the disorder? Describe how this abnormality could result from a defect in meiosis.
- c. Production of offspring by parthenogenesis or cloning bypasses the typical meiotic process. Describe either parthenogenesis or cloning and compare the genomes of the offspring with those of the parents.


AP Biology

B. Calibrate the Spectrophotometer ("blanking")




C. Read the Sample






AP Biology


- Description
 - determine rate of photosynthesis under different conditions
 - light vs. dark
 - boiled vs. unboiled chloroplasts
 - chloroplasts vs. no chloroplasts
 - **♦**use DPIP in place of NADP⁺
 - DPIP_{ox} = blue
 - DPIP_{red} = clear
 - measure light transmittance
 - paper chromatography to separate plant pigments



Photosystem I


- Concepts
 - photosynthesis
 - **♦**Photosystem 1
 - NADPH
 - chlorophylls & other plant pigments
 - chlorophyll a
 - chlorophyll b
 - xanthophylls
 - carotenoids
 - experimental design
 - control vs. experimental







- Conclusions
 - **♦**Pigments
 - pigments move at different rates based on solubility in solvent
 - Photosynthesis
 - light & unboiled chloroplasts produced highest rate of photosynthesis



every 5 minutes for 30 minutes.

Which is the control? #2 (DPIP + chloroplasts + light)

ESSAY 2004 (part 1)

A controlled experiment was conducted to analyze the effects of darkness and boiling on the photosynthetic rate of incubated chloroplast suspensions. The dye reduction technique was used. Each chloroplast suspension was mixed with DPIP, an electron acceptor that changes from blue to clear when it is reduced. Each sample was placed individually in a spectrophotometer and the percent transmittance was recorded. The three samples used were prepared as follows.

Sample 1 —chloroplast suspension + DPIP

Sample 2 —chloroplast suspension surrounded by foil wrap to provide a dark environment + DPIP

Sample 3 —chloroplast suspension that has been boiled + DPIP


Data are given in the table on the next page.

- a. Construct and label a graph showing the results for the three samples.
- b.Identify and explain the control or controls for this experiment.
- c.The differences in the curves of the graphed data indicate that there were differences in the number of electrons produced in the three samples during the experiment. Discuss how electrons are generated in photosynthesis and why the three samples gave different transmittance results.

ESSAY 2004 (part 2)


Time (min)	Light, Unboiled % transmittance Sample 1	Dark, Unboiled % transmittance Sample 2	Light, Boiled % transmittance Sample 3
0	28.8	29.2	28.8
5	48.7	30.1	29.2
10	57.8	31.2	29.4
15	62.5	32.4	28.7
20	66.7	31.8	28.5

AP Biology




AP Biology 2004-2005


- Description
 - ◆using respirometer to measure rate of
 O₂ production by pea seeds
 - non-germinating peas
 - germinating peas
 - effect of temperature
 - control for changes in pressure & temperature in room








- Concepts
 - respiration
 - experimental design
 - control vs. experimental
 - function of KOH
 - function of vial with only glass beads




Conclusions

- **♦**↓temp = ↓respiration
- **♦**↑germination = ↑respiration

- Place three respirometers in a water bath at 10°C with the tips of the pipettes resting on a sling above the water level. Allow several minutes for them to equilibrate.
- Lower the tips of the pipettes into the water.
- Take an initial reading from each respirometer.
- Take readings from each respirometer at 5 minute intervals for the next 15 minutes.
- Use the second set of three vials to measure the rate of respiration in a 25°C bath.



calculate rate?




ESSAY 1990

The results below are measurements of cumulative oxygen consumption by germinating and dry seeds. Gas volume measurements were corrected for changes in temperature and pressure.

Cumulative Oxygen Consumed (mL)							
Time (minutes)	0	10	20	30	40		
Germinating seeds 22°C	0.0	8.8	16.0	23.7	32.0		
Dry Seeds (non-germinating) 22°C	0.0	0.2	0.1	0.0	0.1		
Germinating Seeds 10°C	0.0	2.9	6.2	9.4	12.5		
Dry Seeds (non-germinating) 10°C	0.0	0.0	0.2	0.1	0.2		


- a. Plot the results for the germinating seeds at 22°C and 10°C.
- b. Calculate the rate of oxygen consumption for the germinating seeds at 22°C, using the time interval between 10 and 20 minutes.
- c. Account for the differences in oxygen consumption observed between:
 - 1. germinating seeds at 22°C and at 10°C
 - 2. germinating seeds and dry seeds.
- d. Describe the essential features of an experimental apparatus that could be used to measure oxygen consumption by a small organism. Explain why each of these features is necessary.



been transformed by the amp^R gene will grow.


AP Biology 2004-2005


- Description
 - **♦**Transformation
 - insert foreign gene in bacteria by using engineered plasmid
 - also insert ampicillin resistant gene on same plasmid as selectable marker
 - **♦**Gel electrophoresis
 - cut DNA with restriction enzyme
 - fragments separate on gel based on size



DNA I DNA II Hind III

- Concepts
 - **♦**transformation
 - plasmid
 - **selectable** marker
 - ampicillin resistance
 - restriction enzyme
 - **♦**gel electrophoresis
 - DNA is negatively charged
 - smaller fragments travel faster






AP Biology

Lab 6: Transformation


- Conclusions
 - can insert foreign DNA using vector
 - ampicillin becomes selecting agent
 - no transformation = no growth on amp⁺ plate




Lab 6: Gel Electrophoresis


Conclusions

1. Make gel.



Obtain prepared DNA samples.






DNA = negatively charged

 Separate fragments by electrophoresis.


correlate distance

to size

6. Prepare a standard curve. Determine fragment sizes.



 Stain DNA fragments and measure distances.




smaller fragments travel faster & therefore farther

AP Biol

004-2005

ESSAY 1995

The diagram below shows a segment of DNA with a total length of 4,900 base pairs. The arrows indicate reaction sites for two restriction enzymes (enzyme X and enzyme Y).



Explain how the principles of gel electrophoresis allow for the separation of DNA fragments


- <u>a.</u> <u>Describe</u> the results you would expect from electrophoretic separation of fragments from the following treatments of the DNA segment above. Assume that the digestion occurred under appropriate conditions and went to completion.
- b. DNA digested with only enzyme X
 - I. DNA digested with only enzyme Y
 - II. DNA digested with enzyme X and enzyme Y combined
 - III. Undigested DNA
 - **IV.** Explain both of the following:
- c. The mechanism of action of restriction enzymes
 - 1. The different results you would expect if a mutation occurred at the recognition site for enzyme Y.

ESSAY 2002

The human genome illustrates both continuity and change.


- <u>a. Describe</u> the essential features of <u>two</u> of the procedures/techniques below. For each of the procedures/techniques you describe, <u>explain</u> how its application contributes to understanding genetics.
 - The use of a bacterial plasmid to clone and sequence a human gene
 - Polymerase chain reaction (PCR)
 - Restriction fragment polymorphism (RFLP analysis)
- b. All humans are nearly identical genetically in coding sequences and have many proteins that are identical in structure and function. Nevertheless, each human has a unique DNA fingerprint. Explain this apparent contradiction.

Lab 7: Genetics (Fly Lab)



3 Red-eyed:1 Sepia-eyed offspring

- Description
 - given fly of unknown genotype use crosses to determine mode of inheritance of trait



Concepts

- phenotype vs. genotype
- dominant vs. recessive
- **♦**P, F1, F2 generations
- **♦**sex-linked
- monohybrid cross
- dihybrid cross
- **♦**test cross
- chi square





Recessive Dominant



Conclusions: Can you solve these?


F ₁ RESULTS	OBSERVED PHENOTYPES AND NUMBERS
	Red eyes
o ⁷ MALES	12
Q FEMALES	8



	OBSERVED PHENOTYPES AND NUMBERS						
F ₂ RESULTS	Red eyes	Sepia eyes					
o³ MALES	19	4					
Q FEMALES	12	٥					

Case 2

Case 1



OBSERVED



	OBSERVED PHENOTYPES AND NUMBERS					
F ₂ RESULTS	Red eyes	White eyes				
o ⁷ MALES	12					
Q FEMALES	21					

ESSAY 2003 (part 1)

In fruit flies, the phenotype for eye color is determined by a certain locus. E indicates the dominant allele and e indicates the recessive allele. The cross between a male wild type fruit fly and a female white eyed fruit fly produced the following offspring

	Wild-Type	Wild-Type	White-eyed	White-Eyed	Brown-Eyed
	Male	Female	Male	Female	Female
F-1	0	45	55	0	1

The wild-type and white-eyed individuals from the F1 generation were then crossed to produce the following offspring.

	Wild-Type	Wild-Type	White-eyed	White-Eyed	Brown-Eyed
	Male	Female	Male	Female	Female
F-2	23	31	22	24	0

- a. Determine the genotypes of the original parents (P generation) and explain your reasoning. You may use Punnett squares to enhance your description, but the results from the Punnett squares must be discussed in your answer.
- b. Use a Chi-squared test on the F2 generation data to analyze your prediction of the parental genotypes. Show all your work and explain the importance of your final answer.
- c. The brown-eyed female of the F1 generation resulted from a mutational change. Explain what a mutation is, and discuss two types of mutations that might have produced the brown-eyed female in the F1 generation.

ESSAY 2003 (part 2)

Probability (p)	Degrees of Freedom (df)							
	1 2 3 4 5							
.05	3.84	5.99	7.82	9.49	11.1			

The formula for Chi-squared is:

$$X^2 = \sum \frac{(observed - expected)^2}{expected}$$


size of population & gene pool

Main distributions of population in California, 1997

An earthquake that kills three people out of a population of 10 million has little effect on the composition of the gene pool.




random vs. non-random mating






- Description
 - **♦**simulations were used to study effects of different parameters on frequency of alleles in a population
 - selection
 - heterozygous advantage
 - genetic drift



Concepts

- **◆**Hardy-Weinberg equilibrium
 - p + q = 1
 - $p^2 + 2pq + q^2 = 1$
 - required conditions
 - large population
 - random mating
 - no mutations
 - no natural selection
 - no migration
- **♦**gene pool
- heterozygous advantage
- genetic drift
 - founder effect
 - bottleneck






- Conclusions
 - recessive alleles remain hidden in the pool of heterozygotes
 - even lethal recessive alleles are not completely removed from population
 - know how to solve H-W problems!
 - ◆to calculate <u>allele frequencies</u>, use p + q = 1
 - to calculate genotype frequencies or how many individuals use p 2pq + q² = 1

ESSAY 1989


Do the following with reference to the Hardy-Weinberg model.

- a. Indicate the conditions under which allele frequencies (p and q) remain constant from one generation to the next.
- b. Calculate, showing all work, the frequencies of the alleles and frequencies of the genotypes in a population of 100,000 rabbits of which 25,000 are white and 75,000 are agouti. (In rabbits the white color is due to a recessive allele, w, and agouti is due to a dominant allele, W.)
- c. If the homozygous dominant condition were to become lethal, what would happen to the allelic and genotypic frequencies in the rabbit population after two generations?




AP Biology 2004-2005

- Description
 - test the effects of environmental factors on rate of transpiration
 - temperature
 - humidity
 - air flow (wind)
 - light intensity




- Concepts
 - transpiration
 - **♦**stomates
 - guard cells
 - **♦**xylem
 - adhesion
 - cohesion
 - H bonding




AP Biology

2004-2005

- Conclusions
 - **♦**↑transpiration
 - ↑ wind
 - ↑ light
 - **♦**↓transpiration
 - ↑ humidity





ESSAY 1991

A group of students designed an experiment to measure transpiration rates in a particular species of herbaceous plant. Plants were divided into four groups and were exposed to the following conditions.

Group I:Room conditions (light, low humidity, 20°C, little air movement.)

Group II:Room conditions with increased humidity.


Group III:Room conditions with increased air movement (fan)


Group IV:Room conditions with additional light


The cumulative water loss due to transpiration of water from each plant was measured at 10-minute intervals for 30 minutes. Water loss was expressed as milliliters of water per square centimeter of leaf surface area. The data for all plants in Group I (room conditions) were averaged. The average cumulative water loss by the plants in Group I is presented in the table below.

Average Cumulative Water Loss by the Plants in Group I						
Time (minutes)	Average Cumulative Water Loss (mL H ₂ O/cm ²)					
10	3.5 x 10 ⁻⁴					
20	7.7 x 10 ⁻⁴					
30	10.6 x 10 ⁻⁴					


- 1. Construct and label a graph using the data for Group I. Using the same set of axes, draw and label three additional lines representing the results that you would predict for Groups II, III, and IV.
- 2. Explain how biological and physical processes are responsible for the difference between each of your predictions and the data for Group I.
- 3. Explain how the concept of water potential is used to account for the movement of water from the plant stem to the atmosphere during transpiration.














- Description
 - study factors that affect heart rate
 - body position
 - level of activity
 - determine whether an organism is an endotherm or an ectotherm by measuring change in pulse rate as temperature changes
 - Daphnia

Concepts

- thermoregulation
- **◆**endotherm
- **◆**ectotherm
- **♦**Q₁₀



 Daphnia can adjust their temperature to the environment, as temperature in environment increases, their body temperature also increases which increases their heart rate



환 16X

8X

10

1etabolic

Daphnia

20


30

Temperature (°C)

50



- Conclusions
 - **◆**Activity increase heart rate
 - in a fit individual pulse & blood pressure are lower & will return more quickly to resting condition after exercise than in a less fit individual
 - ◆Pulse rate changes in an ectotherm as external temperature changes




ESSAY 2002


In mammals, heart rate during periods of exercise is linked to the intensity of exercise.

- <u>a. Discuss</u> the interactions of the respiratory, circulatory, and nervous systems during exercise.
- b. Design a controlled experiment to determine the relationship between intensity of exercise and heart rate.
- c. On the axes provided below, <u>indicate</u> results you expect for both the control and the experimental groups for the controlled experiment you described in part B. Remember to label the axes.

AP F









held together with tape

- Description
 - set up an experiment to study behavior in an organism
 - Betta fish agonistic behavior
 - Drosophila mating behavior
 - pillbug kinesis



AP Biology 2004-2005

- Concepts
 - **♦**innate vs. learned behavior
 - experimental design
 - control vs. experimental
 - hypothesis
 - choice chamber
 - temperature
 - humidity
 - light intensity
 - salinity
 - other factors



Hypothesis development


♦Poor:


I think pillbugs will move toward the wet side of a choice chamber.

Better:

If pillbugs prefer a moist environment, then when they are randomly placed on both sides of a wet/dry choice chamber and allowed to move about freely for 10 minutes, most will be found on the wet side.

Experimental design





Aligned "doorways" held together with tape

EXPERIMENT


CONTROL

	Left compartment	Right compartment	Left compartment	Right compartment
MOISTURE	wet	dry	wet	wet
LIGHT	25 Watts	same	same	same
FILTERPAPER	#1 Whatman	same	same	same
TEMPERATURE	20°C	same	same	same
NUMBER OF BUG AT START	S 5	same	same	same

Sample size



Sample size too small.



Sample size better.

AP Biology 2004-2005

ESSAY 1997

A scientist working with *Bursatella leachii*, a sea slug that lives in an intertidal habitat in the coastal waters of Puerto Rico, gathered the following information about the distribution of the sea slugs within a ten-meter square plot over a 10-day period.

time of day	12 mid	4am	8am	12 noon	4pm	8pm	12 mid
average distance between individuals	8.0	8.9	44.8	174.0	350.5	60.5	8.0


- a. For the data above, provide information on each of the following:
 - Summarize the pattern.
 - Identify <u>three</u> physiological or environmental variables that could cause the slugs to vary their distance from each other.
 - <u>Explain</u> how each variable could bring about the observed pattern of distribution.
- b.Choose <u>one</u> of the variables that you identified and <u>design</u> a controlled experiment to test your hypothetical explanation. <u>Describe</u> results that would support or refute your hypothesis.

ESSAY 2002

The activities of organisms change at regular time intervals. These changes are called biological rhythms. The graph depicts the activity cycle over a 48-hour period for a fictional group of mammals called pointy-eared bombats, found on an isolated island in the temperate zone. High

Midnight

a. Describe the cycle of activity for the bombats. Discuss how three of the following factors might affect the physiology and/or



- b. temperature
 - food availability
 - presence of predators

 - Propose a hypothesis regarding the effect of light on the cycle of activity in bombats. Describe a controlled experiment that could be performed to test this hypothesis, and the results you would expect.


Midnight


Midnight


Time of Day


Noon


Dissolved O₂ availability












- Description
 - ◆measure primary productivity by measuring O₂ production
 - **♦** factors that affect amount of dissolved O₂
 - temperature
 - as ↑water temperature, its ability to hold O₂ decreases
 - photosynthetic activity
 - in bright light, aquatic plants produce more O₂
 - decomposition activity
 - as organic matter decays, microbial respiration consumes O₂
 - mixing & turbulence
 - wave action, waterfalls & rapids aerate H₂O & ↑O₂
 - salinity
 - as water becomes more salty, its ability to hold O₂ decreases


- Concepts
 - **♦** dissolved O₂
 - primary productivity
 - measured in 3 ways:
 - amount of CO₂ used
 - rate of sugar (biomass) formation
 - rate of O₂ production
 - net productivity vs. gross productivity
 - respiration



Conclusions

- ♦ \uparrow temperature = \downarrow dissolved O_2
- ♦ ↑ light = ↑ photosynthesis = $↑ O_2$ production
- **♦**O₂ loss from respiration
- **♦**↑respiration = \downarrow dissolved O₂ (consumption of O₂)





ESSAY 2001

A biologist measured dissolved oxygen in the top 30 centimeters of a moderately eutrophic (mesotrophic) lake in the temperate zone. The day was bright and sunny and the wind was calm. The results of the observation are presented below.

- a. Using the graph paper provided, plot the results that were obtained. Then, using the same set of axes, draw and label an additional line/curve representing the results that you would predict had the day been heavily overcast.
- b. Explain the biological processes that are operating in the lake to produce the observed data. Explain also how these processes would account for your prediction of results for a heavily overcast day.
- c. Describe how the introduction of high levels of nutrients such as nitrates and phosphates into the lake would affect subsequent observations. Explain your predictions.

hour	6am	8am	10am	noon	2pm	4pm	6pm	8pm	10pm	mid
[O ₂] mg/L	0.9	1.7	3.1	4.9	6.8	8.1	7.9	6.2	4.0	2.4

ESSAY 2004B

In most aquatic environments, primary production is affected by light available to the community of organisms.

Using measurements of dissolved oxygen concentration to determine primary productivity, design a controlled experiment to test the hypothesis that primary productivity is affected by either the intensity of light <u>or</u> the wavelength of light. In your answer, be sure to include the following.

- A statement of the specific hypothesis that you are testing
- A description of your experimental design (Be sure to include a description of what data you would collect and how you would present and analyze the data using a graph.)
- A description of results that would support your hypothesis