10.3 Percent Composition and Chemical Formulas

Is your shirt made of 100

Connecting to Your World

percent cotton or wool, or is the fabric a combination of two or more fibers? A tag sewed into the seam of the shirt usually tells you what fibers



were used to make the cloth and the percent of each. It helps to know the percents of the components in the shirt because they affect how warm it is, whether it will need to be ironed, and how it should be cleaned. In this section you will learn how the percents of the elements in a compound are important in chemistry.

The Percent Composition of a Compound

If you have had experience with lawn care, you know that the relative amount, or the percent, of each nutrient in fertilizer is important. In spring, you may use a fertilizer that has a relatively high percent of nitrogen to "green" the grass. In fall, you may want to use a fertilizer with a higher percent of potassium to strengthen the root system. Knowing the relative amounts of the components of a mixture or compound is often useful.

The relative amounts of the elements in a compound are expressed as the percent composition or the percent by mass of each element in the compound. The percent composition of a compound consists of a percent value for each different element in the compound. As you can see in Figure 10.13, the percent composition of K_2CrO_4 is K = 40.3%, Cr = 26.8%, and O = 32.9%. These percents must total 100% (40.3% + 26.8% + 32.9% = 100%). C The percent by mass of an element in a compound is the number of grams of the element divided by the mass in grams of the compound, multiplied by 100%.

% mass of element = $\frac{\text{mass of element}}{\text{mass of compound}} \times 100\%$

Percent Composition from Mass Data Imagine you are a chemist who has just finished the synthesis of a new compound. You have purified your product and stored the crystalline solid in a vial. Now you must verify the composition of your new compound and determine its molecular formula. You use analytical procedures to determine the relative masses of each element in the compound and calculate the percent composition.

> Figure 10.13 Potassium chromate (K₂CrO₄) is composed of 40.3% potassium, 26.8% chromium, and 32.9% oxygen. Interpreting Diagrams How does this percent composition differ from the percent composition of potassium dichromate (K₂Cr₂O₇), a compound composed of the same three elements?

Guide for Reading C Key Concepts


- How do you calculate the percent by mass of an element in a compound?
- What does the empirical formula of a compound show?
- How does the molecular formula of a compound compare with the empirical formula?

Vocabulary

percent composition empirical formula

Reading Strategy

Comparing and Contrasting When you compare and contrast things, you examine how they are alike and different. As you read, list the similarities and differences between empirical and molecular formulas.



Section 10.3 Percent Composition and Chemical Formulas 305

– Section Resources ———

Print

- Guided Reading and Study Workbook, Section 10.3
- Core Teaching Resources, Section 10.3 **Review, Interpreting Graphics**
- Laboratory Manual, Lab 13
- Transparencies, T110–T112

Technology

 Interactive Textbook with ChemASAP, Problem-Solving 10.33, 10.34, 10.36, 10.38; Assessment 10.3

• Go Online, Section 10.3

1 FOCUS

Objectives

10.3.1 Describe how to calculate the percent by mass of an element in a compound.

10.3.2 Interpret an empirical formula

10.3.3 Distinguish between empirical and molecular formulas.

Guide for Reading

Build Vocabulary

L2

L2

Paraphrase Have students read the definition of percent composition on this page. Then, have them define the term according to the definition in the dictionary. Their definitions should indicate that percent composition refers to the relative size of parts that make up 100 percent of something.

Reading Strategy

Relate Text and Visuals As they read the chapter, students should examine each visual as it is referenced in the text. Have them read each caption and answer any question.

2 INSTRUCT

Connecting to Your World

Ask, What should be the total of the percents listed on the label? (100%) If the shirt were 25% nylon, what percent would be cotton? (75%)

The Percent Composition of a Compound [1] **Use Visuals**

Figure 10.13 Have students study the figure and read the text on percent composition. Point out that the three numbers in each circle graph add up to a total of 100%. Ask, Which compound is a better source of potassium? (K_2CrO_4)

Answers to

Figure 10.13 The percent composition of K₂Cr₂O₇ is 26.5% K, 35.4% Cr, and 38.1% O.

Sample Problem 10.9

Answers

- 32. Mass of compound = 9.03 g + 3.48 g = 12.51 g; 9.03 g Mg/12.51 g compound × 100% = 72.2% Mg; 3.48 g N/12.51 g compound × 100% = 27.8% N
- **33.** mass of O = 14.2 g 13.2 g = 1.0 g O; 1.0 g O/14.2 g × 100% = 7.0% O; 13.2 g Hg/14.2 g × 100% = 93.0% Hg

L2


Practice Problems Plus

1. What is the percent composition of the compound formed when 2.70 g of aluminum combine with oxygen to form 5.10 g of aluminum oxide? (52.9% Al, 47.1% O)

2. Interactive Textbook with ChemASAP contains the following problem: Calculate the percent composition when 13.3 g Fe combine completely with 5.7 g O. (70% Fe, 30% O)

Handbook


For a math refresher and practice, direct students to percents, page R72.



- Facts and Figures -

Parts per Million and Parts per Billion

Percents are used to show relative parts of mixtures as well as the composition of a compound. But when an extremely small amount of a substance is present in a large amount of another substance, it might not be practical to use percents (parts per one hundred) to show the makeup of the mixture. Instead, concentrations of extremely dilute solutions are sometimes measured in units of parts per million (ppm) or parts per billion (ppb). For example, the composition of a mixture that consists of 1 gram of a substance per 10⁶ grams of water (or 1 milligram of substance per liter of water) can be expressed as 1 ppm.



Section 10.3 Percent Composition and Chemical Formulas 307

Differentiated Instruction –

L3

Gifted and Talented

Have students research the formulas of the three different oxides of iron. Ask, **Which of the oxides contains a higher percent of iron?** (Of FeO (77.7% Fe), Fe₂O₃ (69.9% Fe), and Fe₃O₄ (72.3% Fe), FeO has the highest percent of iron.)

Answers to...

Checkpoint Divide the mass of the element in one mole of the compound by the molar mass and multiply by 100%.

Quick LAB

Percent Composition

Objective After completing this activity, students will be able to:

• determine the percent of water in a hydrate.

Skills Focus observing, calculating

Prep Time 20 minutes Class Time 30 minutes

Safety Students should wear safety goggles and tie back loose hair. Caution students that while heating test tubes, they should not aim the opening of the tube toward anyone. Tell them to move the test tube in the flame and not to heat one spot excessively. CAUTION! Be sure that students allow the tubes to cool completely before they touch them. Hot glass looks exactly like cold glass!

Teaching Tips For best results, students should do a second heating and cooling of each sample to determine whether all of the water has been driven off.

Expected Outcome See data table at the bottom of the page.

Think About It

1.-3. See data table.

4. The hydrated salt of sodium sulfate lost the greatest percent. The hydrated salt of calcium chloride lost the smallest percent.

For Enrichment

L3

Have students design and conduct a similar experiment to determine the percent of oxygen in potassium chlorate. Tell students that when potassium chlorate is heated, potassium chloride and oxygen are produced, 2KClO₃ \rightarrow 2KCl + $3O_2$. For classroom safety, no more than 5 g of potassium chlorate should be used. Results should show that potassium chlorate is approximately 39% oxygen.

Quick LAB

Percent Composition

Purpose

L2


To measure the percent of water in a series of crystalline compounds called hydrates.

Materials

- centigram balance
- Bunsen burner
- 3 medium-sized test tubes
- test tube holder
- test tube rack
- spatula
- hydrated salts of copper(II) sulfate, calcium chloride, and sodium sulfate



- 1. Label each test tube with the name of a salt. Measure and record the masses.
- 2. Add 2-3 g of salt (a good-sized spatula full) to the appropriately labeled test tube. Measure and record the mass of each test tube and salt.
- **3.** Hold one of the tubes at a 45° angle and gently heat its contents over the burner, slowly passing it in and out of the flame. Note any change in the appearance of the solid salt.
- 4. As moisture begins to condense in the upper part of the test tube, gently heat the entire length of the tube. Continue heating until all of the moisture is driven from the tube. This may take 2-3 minutes. Repeat Steps 3 and 4 for the other two tubes.
- 5. Allow each tube to cool. Then measure and record the mass of each test tube and the heated salt.



Analyze and Conclude

- 1. Set up a data table so that you can subtract the mass of the empty tube from the mass of the salt and the test tube, both before and after heating.
- 2. Calculate the difference between the mass of each salt before and after heating. This difference represents the amount of water lost by the hydrate on heating.
- 3. Calculate the percent by mass of water lost by each compound.
- 4. Which compound lost the greatest percent by mass of water? The smallest?

Percent Composition as a Conversion Factor You can use percent composition to calculate the number of grams of any element in a specific mass of a compound. To do this, multiply the mass of the compound by a conversion factor based on the percent composition of the element in the compound. Suppose you want to know how much carbon and hydrogen are contained in 82.0 g of propane. In Sample Problem 10.10, you found that propane is 81.8% carbon and 18% hydrogen. That means that in a 100-g sample of propane, you would have 81.8 g of carbon and 18 g of hydrogen. You can use the ratio $81.8 \text{ g C}/100 \text{ g C}_3 H_8$ to calculate the mass of carbon contained in 82.0 g of propane (C_3H_8) .

$$82.0 \text{ g-C}_{3}\text{H}_{8} \times \frac{81.8 \text{ g C}}{100 \text{ g-C}_{3}\text{H}_{8}} = 67.1 \text{ g C}$$

Using the ratio 18 g H/100 g C_3H_8 , you can calculate the mass of hydrogen.

82.0 g-
$$C_{3}H_{8} \times \frac{18 \text{ g H}}{100 \text{ g-}C_{3}H_{8}} = 15 \text{ g H}$$


The sum of the two masses equals 82 g, the sample size, to two significant figures (67.1 g C + 15 g H = 82 g C_3H_8).

Checkpoint How many grams of hydrogen are contained in a 100-g sample of propane?

308 Chapter 10

Data Table with Sample Data

	CuSO ₄ • 5H ₂ O	CaCl ₂ • 2H ₂ O	Na ₂ SO ₄ • 10H ₂ O
Test tube + hydrate (before heating)	23.88 g	23.60 g	23.92 g
Empty test tube	21.19 g	21.25 g	21.17 g
Mass of hydrate	2.69 g	2.35 g	2.75 g
Test tube + salt (after heating)	22.88 g	23.07 g	22.71 g
Empty test tube	21.19 g	21.25 g	21.17 g
Mass of anhydrous salt	1.69 g	1.82 g	1.54 g
Mass of water lost	1.00 g	0.53 g	1.21 g
Percent water (experimental)	37.2%	22.6%	44.0%
Percent water (theoretical)	36.1%	24.5%	55.9%



Empirical Formulas

A useful formula for cooking rice is to use one cup of rice and two cups of water. If a larger amount of rice is needed, you could double or triple the amounts, for example, two cups of rice and four cups of water. The formulas for some compounds also show a basic ratio of elements. Multiplying that ratio by any factor can produce the formulas for other compounds.

The percent composition of your newly synthesized compound is the data you need to calculate the basic ratio of the elements contained in the compound. The basic ratio, called the **empirical formula**, gives the lowest whole-number ratio of the atoms of the elements in a compound. For example, a compound may have the empirical formula CO_2 . The empirical formula shows the kinds and lowest relative count of atoms or moles of atoms in molecules or formula units of a compound. Figure 10.15 shows that empirical formulas may be interpreted at the microscopic (atomic) or macroscopic (molar) level.

An empirical formula may or may not be the same as a molecular formula. For example, the lowest ratio of hydrogen to oxygen in hydrogen peroxide is 1:1. Thus the empirical formula of hydrogen peroxide is HO. The actual molecular formula of hydrogen peroxide has twice the number of atoms as the empirical formula. The molecular formula is $(HO) \times 2$, or H_2O_2 . But notice that the ratio of hydrogen to oxygen is still the same, 1:1. The empirical formula of a compound shows the smallest whole-number ratio of the atoms in the compound. The molecular formula tells the actual number of each kind of atom present in a molecule of the compound. For carbon dioxide, the empirical and molecular formulas are the same— CO_2 . Figure 10.16 shows two compounds of carbon having the same empirical formula (CH) but different molecular formulas.

Figure 10.16 Ethyne (C_2H_2) , also called acetylene, is a gas used in welder's torches. Styrene (C_8H_8) is used in making polystyrene. These two compounds have the same empirical formula. **Calculating** *What is the empirical formula of ethyne and styrene?*

Figure 10.15 A formula can be interpreted on a microscopic level in terms of atoms or on a macroscopic level in terms of moles of atoms.

Word Origins

Empirical comes from the Latin word *empiricus* meaning a doctor relying on experience alone. An empirical formula must be obtained from experimental data. Thus, an empirical formula relies on experience. Is a molecular formula also based on experimental data?





Section 10.3 Percent Composition and Chemical Formulas 309

Empirical Formulas



Empirical Formulas from Percent Composition

L2

Purpose Students are provided with an analogy that helps clarify the concepts of percent composition and empirical formulas.

Materials 3 red marbles, 6 green marbles, 3 black marbles, and 12 blue marbles

Procedure Provide pairs of students with sets of marbles. Have students express the number of different colored marbles as fractions and percents of the whole collection. Ask, What percent of the collection do the red marbles represent? (12.5%) Show them that the sums of the fractions and percents are equal to 1 and 100%, respectively. Ask, What is the ratio of red: green:black:blue marbles in lowest terms? (1:2:1:4) This activity can be extended if the different colored marbles are assumed to be atoms of different elements. Ask, What is the empirical formula of a hypothetical "compound" that consists of 25% red marbles and 75% green marbles? (The ratio of red marbles to green marbles in the empirical formula would be 1:3.)

Expected Outcomes Students express percent composition of the marbles and determine the "empirical formula" of a marble combination.

Word Origins

A molecular formula is based on experimental data in two different ways. The molar mass is determined experimentally, as is the empirical formula.

Differentiated Instruction -

Less Proficient Readers

As students read about how to determine empirical and molecular formulas, have groups of students develop numbered lists of steps they would take to determine these formulas. They should have three lists: one

1

for determining empirical formulas from percent composition; one for determining empirical formulas from mass data; and one for determining molecular formulas from the empirical formula and molar mass.



Sample Problem 10.11

Answers

36. a. 94.1 g O × 1 mol O/16.0 g O = 5.88 mol O $5.9 \text{ g H} \times 1 \text{ mol H}/1.0 \text{ g H} = 5.9 \text{ mol H}$ 5.88 mol O/5.88 = 1.00 mol O 5.9 mol H/5.88 = 1.0 mol H Empirical formula = HO **b.** 67.6 g Hg \times 1 mol Hg/200.6 g Hg = 0.337 mol Ha 10.8 g S × 1 mol S/32.1 g S = 0.336 mol S 21.6 g O × 1 mol O/16.0 g O = 1.35 mol O 0.337 mol Hg/0.336 = 1.00 mol Hg 0.336 mol S/0.336 = 1.00 mol S 1.35 mol O/0.336 = 4.02 mol O Empirical formula = $HgSO_4$ **37.** 62.1 g C × 1 mol C/12.0 g C = 5.18 mol C $13.8 \text{ g H} \times 1 \text{ mol H}/1.00 \text{ g H} = 13.8 \text{ mol H}$ 24.1 a N \times 1 mol N/14.0 a N = 1.72 mol N 5.18 mol C/1.72 = 3.01 mol C 13.8 mol H/1.72 = 8.02 mol H 1.72 mol N/1.72 = 1.00 mol N Empirical formula = C_3H_8N L2


Practice Problems Plus What is the empirical formula of

each of the following compounds? a. 36.1% Ca, 63.9% Cl (*CaCl*₂) b. 40.0% C, 6.7% H, 53.3% O (*CH*₂O) c. 3.7% H, 44.4% C, and 51.9% N (*HCN*)

Math

For a math refresher and practice, direct students to dimensional analysis, page R66.

Handbook



SAMPLE PROBLEM 10.11

– Facts and Figures –

Computing Formulas

In using percent composition to determine empirical formula, 100.0 g of compound is arbitrarily chosen because it is easy to use. If an element comprises 28.5% of the mass of a compound, for example, it makes up 28.5 g of a 100.0-g sample. Any other mass of compound can be used but computation will be more difficult.

Table 10.3

Comparison of Empirical and Molecular Formulas			
Formula (name)	Classification of formula	Molar mass	
СН	Empirical	13	
C ₂ H ₂ (ethyne)	Molecular	26 (2 $ imes$ 13)	
C ₆ H ₆ (benzene)	Molecular	78 (6 $ imes$ 13)	
CH ₂ O (methanal)	Empirical and Molecular	30	
$C_2H_4O_2$ (ethanoic acid)	Molecular	60 (2 $ imes$ 30)	
C ₆ H ₁₂ O ₆ (glucose)	Molecular	180 (6 $ imes$ 30)	

Molecular Formulas

Look at the compounds listed in Table 10.3. Ethyne and benzene have the same empirical formula—CH. Methanal, ethanoic acid, and glucose, shown in Figure 10.17 have the same empirical formula—CH₂O. But the compounds in these two groups have different molar masses. Their molar masses are simple whole-number multiples of the molar masses of the empirical formulas, CH and CH₂O. The molecular formula of a compound is either the same as its experimentally determined empirical formula, or it is a simple whole-number multiple of its empirical formula.

Once you have determined the empirical formula of your newly synthesized compound, you can determine its molecular formula, but you must know the compound's molar mass. A chemist often uses an instrument called a mass spectrometer to determine molar mass. The compound is broken into charged fragments (ions) that travel through a magnetic field. The magnetic field deflects the particles from their straight-line paths. The mass of the compound is determined from the amount of deflection experienced by the particles.

From the empirical formula, you can calculate the empirical formula mass (efm). This is simply the molar mass represented by the empirical formula. Then you can divide the experimentally determined molar mass by the empirical formula mass. This gives the number of empirical formula units in a molecule of the compound and is the multiplier to convert the empirical formula to the molecular formula. For example, recall that the empirical formula of hydrogen peroxide is HO. Its empirical formula mass is 17.0 g/mol. The molar mass of H₂O₂ is 34.0 g/mol.

$$\frac{34.0 \text{ g/mot}}{17.0 \text{ g/mot}} = 2$$

To obtain the molecular formula of hydrogen peroxide from its empirical formula, multiply the subscripts in the empirical formula by 2. (HO) \times 2 = H₂O₂.

Checkpoint How does the molecular formula for a compound relate to its empirical formula?

Figure 10.17 Methanal (formaldehyde), ethanoic acid (acetic acid), and glucose have the same empirical formula. Applying Concepts How could you easily obtain the molar mass of glucose using the molar mass of methanal?

Section 10.3 Percent Composition and Chemical Formulas **311**

CTHANOIG ADD

OgH, Da

Facts and Figures -

Some Carbohydrates Share the Same Empirical Formula

Several carbohydrates have the empirical formula CH₂O. Examples include glucose, which is abundant in plants and animals, and fructose, which is found in fruits and honey. Both of these simple structures have the molecular formula $C_6H_{12}O_6$, but they differ in structure.

Molecular Formulas Use Visuals

Table 10.3 Have students examine the patterns shown in the table. Then have them add entries for these pairs of compounds: nitrogen dioxide (NO_2) and dinitrogen tetroxide (N_2O_4) , diphosphorus pentoxide (P_2O_5) and tetraphosphorus decoxide (P_4O_{10}) . Students can determine the empirical formula for each pair.

11

Answers to...

Figure 10.17 Multiply the molar mass of methanal by 6.

Checkpoint The molecular formula of a compound is the same as the empirical formula or it is a whole-number multiple of it.

Sample Problem 10.12

Answers

- **38.** molar mass/efm = 62/31 = 2 molecular formula = $2(CH_3O) =$ $C_2H_6O_2$
- **39. a.** same empirical formula (CH₂O) **b.** different empirical formulas

L2

[1]

Practice Problems Plus

1. What is the molecular formula of a compound with the empirical formula CCIN and a molar mass of **184.5?** $(C_3Cl_3N_3)$ 2. What is the molecular formula of a compound that is 56.6% K, 8.7% C, and 34.7% O? (K₂CO₃)

B ASSESS

L2 **Evaluate Understanding**

Have students list the steps they would take to calculate the molecular formula in each of the following situations:

- The empirical formula and molar mass are known.
- The percent composition and molar mass are known.

Reteach

Point out to students that when they know the percent composition and molar mass of a compound, they must first use the percent composition to calculate the empirical formula. They can then calculate the empirical formula mass and compare it to the molar mass of the molecular compound to determine the molecular formula.

Elements Handbook

CaO: 71.5% CaCO₃: 40.1% Ca(OH)₂:54.1% CaSO₄ • 2H₂O: 23.3% Ca₃(PO₄)₂: 38.8%



If your class subscribes to the Interactive Textbook, use it to review key concepts in Section 10.3.

, with **ChemASAP**

	SAMPLE PROBLEM 10.12Finding the Molecular Formula of a CompoundCalculate the molecular formula of a compound whose molar mass is60.0 g/mol and empirical formula is CH4N.			
	1 Analyze List the knowns and the unknown. Knowns Unknown • empirical formula = CH ₄ N • molecular formula = ? • molar mass = 60.0 g/mol			
	Calculate Solve for the unknown. First calculate the empirical formula mass. Then divide the molar mass by the empirical formula mass to obtain a whole number. To get the molecular formula, multiply the formula subscripts by this value. Empirical formula efm Molar mass/efm Molecular formula			
3 Textbook Problem-Solving 10.38 Solve Problem 38 with the help of an interactive guided	CH_4N 30.0 $60.0/30.0 = 2$ $C_2H_8N_2$ 3 Evaluate Does the result make sense? The molecular formula has the molar mass of the compound. Practice Problems			
	 38. Find the molecular formula of ethylene glycol, which is used as antifreeze. The molar mass is 62 g/mol and the 39. Which pair of molecules has the same empirical formula? a. C₂H₄O₂, C₆H₁₂O₆ b. NaCrO₄, Na₂Cr₂O₇ 			

empirical formula is CH₃O.

10.3 Section Assessment

with ChemASAP

tutorial.

- 40. 🕞 Key Concept How do you calculate the percent by mass of an element in a compound?
- 41. 🕞 Key Concept What information can you obtain from an empirical formula?
- 42. 👝 Key Concept How is the molecular formula of a compound related to its empirical formula?
- 43. Calculate the percent composition of the compound that forms when 222.6 g N combines completely with 77.4 g O.
- 44. Calculate the percent composition of calcium acetate (Ca($C_2H_3O_2$)₂).
- 45. The compound methyl butanoate smells like apples. Its percent composition is 58.8% C, 9.8% H, and 31.4% O and its molar mass is 102 g/mol. What is its empirical formula? What is its molecular formula?

312 Chapter 10

- 46. What is an empirical formula? Which of the following molecular formulas are also empirical formulas? a. ribose $(C_5H_{10}O_5)$

 - **b.** ethyl butyrate ($C_6H_{12}O_2$)
 - c. chlorophyll ($C_{55}H_{72}MgN_4O_5$)

d. DEET (C₁₂H₁₇ON)

Elements Handbook

Calcium Select three important compounds that contain calcium from among those discussed on page R11 of the Elements Handbook. Determine the percent of calcium in each.

Interactive Textbook Assessment 10.3 Test yourself on the concepts in Section 10.3.

with ChemASAP

Section 10.3 Assessment

- 40. Divide the mass of an element in the compound by the mass of the compound; then multiply by 100%.
- **41.** The empirical formula gives the lowest whole-number ratio of atoms in the compound.
- 42. The molecular formula of a compound is a simple whole-number multiple of the empirical formula.
- 43. 74.2% N, 25.8% O **44.** 25.4% Ca, 30.4% C, 3.8% H, 40.5% O
- **45.** C₅H₁₀O₂ is both its empirical and molecular formula.
- **46.** An empirical formula has the lowest whole-number ratio of elements.
 - a.molecular b.molecular
 - c. molecular and empirical
 - d. molecular and empirical

Technology & Society

Drug Testing

A test to identify an abused substance in the body must be extremely accurate. A false-positive result could ruin a career. A false-negative result could endanger lives. The best method currently available to test for drug abuse is the gas chromatography/mass spectrometer system, or GC/MS. Gas chromatography separates a chemical mixture and identifies its components. Mass spectrometry uses masses to verify the identification. Used together, the GC/MS testing is nearly 100% reliable. Interpreting Diagrams What is the purpose of the separation column?

The separation column is packed with a

vaporized sample. Components of the mixture move at different rates along the column and

then pass through a detector.

material that interacts physically with the

Detector

KI SIL LINEY

eated metal block

The sample to be tested. such as urine or blood serum, is vaporized and mixed with an unreactive carrier gas such as helium. The resulting gas mixture is forced through a separation column.

Facts and Figures -

Drugs in the Blood

Current drug tests can detect even small traces of drugs in the blood or urine. For example, THC (the metabolic product of marijuana use) remains detectable in the body for at least 18 hours for an occasional user and up to 30 days for a habitual user. It can be detected in concentrations as small as 50 ng/mL of solution.



The separated component

enters a mass spectrometer

Technology and Society 313

where its identity is confirmed.


4

Each peak on a gas chromatogram

shows the retention time for a

component. Retention time is used to

identify compounds.



Drug Testing Discuss

12

L2

Chromatography Discuss with students the basic concepts of chromatography. Chromatography is based on the relative attraction of the material in the separation column (stationary phase) for the materials in the medium that moves through it (mobile phase). The amount of attraction is based on the bonding in the molecules in the different phases. If the bonds in one phase are polar and the bonds in the other phase are nonpolar, the phases have little attraction for each other, and the mobile phase passes quickly through the stationary phase. If the molecules in both phases are either polar or nonpolar, the attraction between the molecules in the phases causes this fraction of the mobile phase to move more slowly through the stationary phase.

Demo TEACHER

Paper Chromatography

Purpose Students observe an example of a mixture being separated into its components.

Materials coffee filter paper, water soluble marker, beaker

Procedure Cut a strip of filter paper. About one-third of the distance from the bottom of the paper, make a mark on the paper. Place water in the beaker to a depth of 2 cm. Place the end of the filter paper closest to the mark in the beaker, making sure the mark is above the water line. When the water rises almost to the top of the paper, remove the paper from the water, and observe what happened.

Expected Outcomes The components of the ink separate, based on the attraction of the components for the paper.

Answers to ...

Interpreting Diagrams The column concentrates and separates the components of the mixture by causing each to move at a different rate.