


DIASTAT® AcuDial™ (diazepam rectal gel) Administration Instructions



1 Put person on their side where they can't fall.



2 Get medicine.



3 Get syringe.

Note: Seal Pin is attached to the cap.



4 Push up with thumb and pull to remove cap from syringe.

Be sure Seal Pin is removed with the cap.



5 Lubricate rectal tip with lubricating jelly.



6 Turn person on side facing you.



7 Bend upper leg forward to expose rectum.



8 Separate buttocks to expose rectum.



9 Gently insert syringe tip into rectum.

Note: Rim should be snug against rectal opening.



10 Slowly count to 3 while gently pushing plunger in until it stops.



11 Slowly count to 3 before removing syringe from rectum.



12 Slowly count to 3 while holding buttocks together to prevent leakage.



13 **ONCE DIASTAT® IS GIVEN**
Keep person on side facing you, note time given and continue to observe.

DISPOSAL INSTRUCTIONS FOR DIASTAT ACUDIAL

14a

- Pull on plunger until it is completely removed from the syringe body.
- Point tip over sink or toilet.

- Replace plunger into syringe body, gently pushing plunger until it stops.
- Flush toilet or rinse sink with water until gel is no longer visible.

SINK OR TOILET

This step is for Diastat® AcuDial™ users only
At the completion of step 14a:

- Discard all used materials in the garbage can.
- Do not reuse.
- Discard in a safe place away from children.

DISPOSAL FOR DIASTAT 2.5 MG
At the completion of step 13:

14b

- Discard all used materials in the garbage can.
- Do not reuse.
- Discard in a safe place away from children.

Call for Help if any of the Following Occur

- Seizure(s) continues 15 minutes after giving DIASTAT or per the doctor's instructions:

- Seizure behavior is different from other episodes
- You are alarmed by the frequency or severity of the seizure(s)
- You are alarmed by the color or breathing of the person
- The person is having unusual or serious problems

Local emergency number: _____

Doctor's number: _____

(please be sure to note if your area has 911)

Information for emergency squad: Time DIASTAT given: _____ Dose: _____

IMPORTANT SAFETY INFORMATION

DIASTAT® AcuDial™ (diazepam rectal gel) is a gel formulation of diazepam intended for rectal administration in the management of selected refractory patients with epilepsy, on stable regimens of AEDs, who require intermittent use of diazepam to control bouts of increased seizure activity for patients 2 years and older.

In clinical trials with DIASTAT®, the most frequent adverse event was somnolence (23%). Less frequent adverse events reported were dizziness, headache, pain, vasodilation, diarrhea, ataxia, euphoria, incoordination, asthma, rash, abdominal pain, nervousness and rhinitis (1%-5%). Please see full DIASTAT AcuDial Prescribing Information.