

Disaster Medical Operations — Part 2

CERT Basic Training
Unit 4





Unit 3 Review

he Ric

- 3 "Killers"
 - Airway obstruction
 - Excessive bleeding
 - Shock









CERT Sizeup

- Gather Facts
- 2. Assess Damage
- 3. Consider Probabilities
- Assess Your Situation
- 5. Establish Priorities
- 6. Make Decisions
- Develop Plan of Action
- Take Action
- 9. Evaluate Progress



REMEMBER:

CERT SIZEUP IS A
CONTINUAL
PROCESS







Unit Objectives had been

- Take appropriate sanitation measures to help protect public health
- Perform head-to-toe patient assessments
- Establish a treatment area
- Apply splints to suspected fractures and sprains
- Employ basic treatments for other injuries





Unit Topics

- No Re
- Public Health
 Considerations
- Functions of Disaster Medical Operations
- Establishing Medical Treatment Areas
- Conducting Head-to-Toe Assessments
- Treating Burns
- Wound Care


- Treating Fractures,
 Dislocations, Sprains,
 and Strains
- Nasal Injuries
- Treating Cold-Related Injuries
- Treating Heat-Related Injuries
- Bites and Stings





Public Health Considerations

- Maintaining proper hygiene
- Maintaining proper sanitation
- Purifying water (if necessary)
- Preventing spread of disease








Maintaining Hygiene

- Wash hands frequently
 - Or use alcohol-based hand sanitizer
- Wear non-latex exam gloves
- Wear N95 mask and goggles
- Keep dressings sterile
- Avoid contact with body fluids
 - "If it is warm, wet, and not yours, don't touch it!"







Maintain Sanitation

- Control disposal of bacterial sources
- Put waste products in plastic bags
 - Tie off bags and mark them as medical waste
- Bury human waste





Water Sanitation Methods

- Boil water for 1 minute
- Water purification tablets
- Non-perfumed liquid bleach
 - 8 drops/gal of water
 - 16 drops/gal if water is cloudy
 - Let stand for 30 minutes before use



Functions of Disaster Medical Operations

- Triage
- Treatment
- Transport
- Morgue
- Supply



Establish a Medical Treatment Area

- Select site and set up treatment area as soon as injured survivors are confirmed
- When determining best location(s) for treatment area, consider:
 - Safety of rescuers and survivors
 - Most effective use of resources









Treatment Area Site Selection

- The site selected should be:
 - In a safe area, free of hazards and debris
 - Upwind, uphill, and upstream (if possible) from hazard zone(s)
 - Accessible by transportation vehicles
 - Expandable





The treatment site should be uphill and upwind from the hazard.





Most Effective Use of CERT Resources

- To help meet the challenge of limited resources,
 CERT may need to establish:
 - Decentralized medical treatment location (more than one location)
 - Centralized medical treatment location (one location)





Treatment Area Layout


- Four treatment areas:
 - "I" for Immediate care
 - "D" for Delayed care
 - "M" for Minor injuries/walking wounded
 - "DEAD" for the morgue







Treatment Area Layout



Treatment area layout, showing the organization for the incident site, triage, transportation, and morgue





Treatment Area Organization


- Assign treatment leader to each treatment area
- Document thoroughly
 - Available identifying information
 - Description (age, sex, body build, estimated height)
 - Clothing
 - Injuries
 - Treatment
 - Transfer location





Head-to-Toe Assessment

- Objectives of head-totoe assessment:
 - Determine extent of injuries
 - Determine type of treatment needed
 - Document injuries








DCAP-BTLS

- Deformities
- Contusions
- Abrasions
- Punctures
- Burns
- Tenderness
- Lacerations
- Swelling







Where and When

- Light damage: assess in place
- Moderate damage: move to treatment area first
- Assess and tag everyone
- Both verbal and hands on





Conducting Head-to-Toe Assessment

- Pay careful attention
- Look, listen, and feel
- Check own hands for patient bleeding
- If you suspect a spinal injury in unconscious survivors, treat accordingly
- Check PMS in all extremities
- Look for medical identification





Order of Assessment

- 1. Head
- 2. Neck
- 3. Shoulders
- 4. Chest
- 5. Arms
- 6. Abdomen
- 7. Pelvis
- 8. Legs



Closed-Head, Neck, Spinal Injuries

- Do no harm
 - Minimize movement of head and neck
- Keep spine in straight line
- Stabilize head





Treating Burns h

- Conduct thorough sizeup
- Treat with first aid
 - Cool burned area
 - Cover with sterile cloth to reduce risk of infection





Burn Severity

- to He
- Factors that affect burn severity:
 - Temperature of burning agent
 - Period of time survivor exposed
 - Area of body affected
 - Size of area burned
 - Depth of burn






Burn Classifications -

- Superficial: epidermis
- Partial Thickness: dermis and epidermis

Full Thickness: subcutaneous layer and all

layers above







Burn Treatment: DOS

- When treating a burn survivor, DO:
 - Cool skin or clothing if they are still hot
 - Cover burn loosely with dry, sterile dressings to keep air out, reduce pain, and prevent infection
 - Elevate burned extremities





Burn Treatment: DON'Ts

- When treating a burn survivor, DO NOT:
 - Use ice
 - Apply antiseptics, ointments, or other remedies
 - Remove shreds of tissue, break blisters, or remove adhered particles of clothing





Treatment for Chemical Burns

- Remove cause of burn + affected clothing/jewelry
- If irritant is dry, gently brush away as much as possible
 - Always brush away from eyes, survivor, and you
- Flush with lots of cool running water
- Apply cool, wet compress to relieve pain
- Cover wound loosely with dry, sterile or clean dressing
- Treat for shock if appropriate







Inhalation Burns Signs and Symptoms

- Sudden loss of consciousness
- Evidence of respiratory distress or upper airway obstruction
- Soot around mouth or nose
- Singed facial hair
- Burns around face or neck




Figure 27-26 A singed mustache and burns to the tip of the tongue signal danger of airway burns or burns to the eyes.





Wound Care

- In The
- Control bleeding
- Clean wound
- Apply dressing and bandage





Cleaning and Bandaging Wounds

- Clean by irrigating with clean, room temperature water
 - NEVER use hydrogen peroxide
 - Irrigate but do not scrub
- Apply dressing and bandage
 - Dressing applied directly to wound
 - Bandage holds dressing in place





Rules of Dressing

- If active bleeding:
 - Redress OVER existing dressing
- If no active bleeding:
 - Remove bandage and dressing to flush wound
 - Check for infection every 4-6 hours





Signs of Infection

- Signs of possible infection
 - Swelling around wound site
 - Discoloration
 - Discharge from wound
 - Red striations from wound site









Amputations

- No. He
- Control bleeding; treat shock
- If amputated body part is found:
 - Save tissue parts, wrapped in clean material and placed in plastic bag
 - Keep tissue parts cool, but NOT directly on ice
 - Keep severed part with survivor





Impaled Objects had been

- When foreign object is impaled in patient's body:
 - Immobilize affected body part
 - Do not attempt to move or remove
 - Try to control bleeding at entrance wound
 - Clean and dress wound, making sure to stabilize impaled object






Fractures, Dislocations, Sprains, Strains

- Immobilize injury and joints immediately above and below injury site
- If uncertain of injury type, treat as fracture





Types of Fractures








Treating Open Fractures

- Do not draw exposed bone ends back into tissue
- Do not irrigate wound
- Cover wound with sterile dressing
- Splint fracture without disturbing wound
- Place moist dressing over bone end





Displaced and Nondisplaced Fractures







Dislocations

- No Ric
- Dislocation is injury to ligaments around joint
 - So severe that it permits separation of bone from its normal position in joint
- Treatment
 - Immobilize; do NOT relocate
 - Check PMS before and after splinting/ immobilization





Signs of Sprain has Too

- Tenderness at site
- Swelling and bruising
- Restricted use or loss of use








Splinting

he Ric







Splinting Guidelines -

- Support injured area above and below injury
- 2. Assess PMS in extremity
- 3. Splint injury in position that you find it
- 4. Don't try to realign bones or joints
- 5. Fill voids to stabilize and immobilize
- 6. Immobilize above and below injury
- 7. After splinting, reassess PMS





Nasal Injuries



Causes

- Blunt force to nose
- Skull fracture
- Nontrauma conditions, e.g., sinus infections, high blood pressure, and bleeding disorders

Cautions


- Large blood loss from nosebleed can lead to shock
- Actual blood loss may not be evident because survivor will swallow some amount of blood





Treatment of Nasal Injuries

- Control nasal bleeding:
 - Pinch nostrils or put pressure on upper lip under nose
 - Have survivor sit with head forward, NOT back
- Ensure that airway remains open
- Keep survivor calm







Cold-Related Injuries


- Hypothermia :
 - Occurs when body's temperature drops below normal
- Frostbite:
 - Occurs when extreme cold shuts down blood flow to extremities, causing tissue death





Symptoms of Hypothermia

- Body temperature of 95° F or lower
- Redness or blueness of skin
- Numbness and shivering
- Slurred speech
- Unpredictable behavior
- Listlessness







Hypothermia Treatment

- Remove wet clothing
- Wrap survivor in blanket
- Protect survivor from weather
- Provide food and drink to conscious survivors
- Do not attempt to massage to warm body
- Place unconscious survivor in recovery position
- Place survivor in warm bath





Symptoms of Frostbite

- Skin discoloration
- Burning or tingling sensation
- Partial or complete numbness







Frostbite Treatment

- Immerse injured area in warm (NOT hot) water
 - Warm slowly!
- Do NOT allow part to re-freeze
- Do NOT attempt to use massage
- Wrap affected body parts in dry, sterile dressing







Heat-Related Injuries

Heat cramps:

- Muscle spasms brought on by over-exertion in extreme heat
- Heat exhaustion:
 - Occurs when exercising or working in extreme heat results in loss of body fluids
- Heat stroke:
 - Survivor's temperature control system shuts down
 - Body temperature rises so high that brain damage and death may result





Symptoms of Heat Exhaustion

- Cool, moist, pale or flushed skin
- Heavy sweating
- Headache
- Nausea or vomiting
- Dizziness
- Exhaustion





Symptoms of Heat Stroke

- Hot, red skin
- Lack of perspiration
- Changes in consciousness
- Rapid, weak pulse and rapid, shallow breathing





Treatment of Heat-Related Injuries

- Remove from heat to cool environment
- Cool body slowly
- Have the survivor drink water, SLOWLY
- No food or drink if survivor is experiencing vomiting, cramping, or is losing consciousness





Treatment for Bites/Stings

- If bite or sting is suspected, and situation is non-emergency:
 - Remove stinger if still present by scraping edge of credit card or other stiff, straightedged object across stinger
 - Wash site thoroughly with soap and water
 - Place ice on site for 10 minutes on and 10 minutes off





Anaphylaxis

- to Tie
- Check airway and breathing
- Calm individual
- Remove constrictive clothing and jewelry
- Find and help administer survivor's Epi-pen
- Watch for signs of shock and treat appropriately







Unit Summary

- he He
- Public health concerns related to sanitation, hygiene, and water purification
- Organization of disaster medical operations
- Establishing treatment areas
- Conducting head-to-toe assessments
- Treating wounds, fractures, sprains, and other common injuries





Homework Assignment

- Read unit to be covered in next session
- Bring necessary supplies for next session
- Wear appropriate clothes for next session
- Practice complete head-to-toe assessment on friend or family member





