



Georgia Standards of Excellence Frameworks

Mathematics

GSE Pre-Calculus

Unit 3: Trigonometry of General Triangles



Richard Woods, Georgia's School Superintendent
"Educating Georgia's Future"

Unit 3

Trigonometry of General Triangles

Table of Contents

OVERVIEW	3
STANDARDS ADDRESSED IN THIS UNIT	3
KEY STANDARDS	3
RELATED STANDARDS	4
STANDARDS FOR MATHEMATICAL PRACTICE	4
ENDURING UNDERSTANDINGS	4
ESSENTIAL QUESTIONS	5
CONCEPTS/SKILLS TO MAINTAIN	5
SELECTED TERMS AND SYMBOLS	5
CLASSROOM ROUTINES	6
STRATEGIES FOR TEACHING AND LEARNING	7
EVIDENCE OF LEARNING	8
TASKS	9
FINDING A NEW AREA FORMULA FOR TRIANGLES LEARNING TASK	10
PROVING THE LAW OF COSINES	16
PROVING THE LAW OF SINES	25
THE HINGE THEOREM	33
CULMINATING TASK: COMBINING LOTS	40

OVERVIEW

In this unit students will:

- Expand the use of trigonometric functions beyond right triangles into more general triangles.
- Develop the trigonometric formula for area of triangle.
- Use the Laws of Sines and Cosines to solve problems.

Although the units in this instructional framework emphasize key standards and big ideas at specific times of the year, routine topics such as estimation, mental computation, and basic computation facts should be addressed on an ongoing basis. Ideas related to the eight practice standards should be addressed constantly as well. To assure that this unit is taught with the appropriate emphasis, depth, and rigor, it is important that the tasks listed under “Evidence of Learning” be reviewed early in the planning process. A variety of resources should be utilized to supplement this unit. This unit provides much needed content information, but excellent learning activities as well. The tasks in this unit illustrate the types of learning activities that should be utilized from a variety of sources.

STANDARDS ADDRESSED IN THIS UNIT

Mathematical standards are interwoven and should be addressed throughout the year in as many different units and activities as possible in order to emphasize the natural connections that exist among mathematical topics. Students will build on their previous experience with trigonometric functions to include applications of the Laws of Sines and Cosines.

KEY STANDARDS

Apply trigonometry to general triangles

MGSE9-12.G.SRT.9 Derive the formula $A = (1/2)ab \sin(C)$ for the area of a triangle by drawing an auxiliary line from a vertex perpendicular to the opposite side.

MGSE9-12.G.SRT.10 Prove the Laws of Sines and Cosines and use them to solve problems.

MGSE9-12.G.SRT.11 Understand and apply the Law of Sines and the Law of Cosines to find unknown measurements in right and non-right triangles (e.g., surveying problems, resultant forces).

RELATED STANDARDS

Define trigonometric ratios and solve problems involving right triangles

MGSE9-12.G.SRT.6 Understand that by similarity, side ratios in right triangles are properties of the angles in the triangle, leading to definitions of trigonometric ratios for acute angles.

MGSE9-12.G.SRT.7 Explain and use the relationship between the sine and cosine of complementary angles.

MGSE9-12.G.SRT.8 Use trigonometric ratios and the Pythagorean Theorem to solve right triangles in applied problems.

STANDARDS FOR MATHEMATICAL PRACTICE

Refer to the Comprehensive Course Overview for more detailed information about the Standards for Mathematical Practice.

1. Make sense of problems and persevere in solving them.
2. Reason abstractly and quantitatively.
3. Construct viable arguments and critique the reasoning of others.
4. Model with mathematics.
5. Use appropriate tools strategically.
6. Attend to precision.
7. Look for and make use of structure.
8. Look for and express regularity in repeated reasoning.

ENDURING UNDERSTANDINGS

- Derive a formula for the area of a triangle using two sides and a non-included angle
- Calculate the area of a general triangle using $A = \frac{1}{2}ab(\sin C)$
- Verify the Law of Sines and the Law of Cosines for a general triangle.
- Apply the Law of Sines and the Law of Cosines to solve problems.

ESSENTIAL QUESTIONS

- How can I calculate the area of any triangle given only two sides and a non-included angle?
- How can I apply trigonometric relationships to non-right triangles?
- What is the least amount of information that is sufficient to find all six parts of a triangle?

CONCEPTS/SKILLS TO MAINTAIN

It is expected that students will have prior knowledge/experience related to the concepts and skills identified below. It may be necessary to pre-assess in order to determine if time needs to be spent on conceptual activities that help students develop a deeper understanding of these ideas.

- number sense
- computation with whole numbers and decimals, including application of order of operations
- calculating the area of a triangle
- solving trigonometric equations
- using inverse trigonometric functions to solve problems
- constructing altitudes in a triangle
- performing operations with trigonometric functions

SELECTED TERMS AND SYMBOLS

The following terms and symbols are often misunderstood. These concepts are not an inclusive list and should not be taught in isolation. However, due to evidence of frequent difficulty and misunderstanding associated with these concepts, instructors should pay particular attention to them and how their students are able to explain and apply them.

The definitions below are for teacher reference only and are not to be memorized by the students. Students should explore these concepts using models and real life examples. Students should understand the concepts involved and be able to recognize and/or demonstrate them with words, models, pictures, or numbers.

The website below is interactive and includes a math glossary.

<http://intermath.coe.uga.edu/dictionary/homepg.asp>

Definitions and activities for these and other terms can be found on the Intermath website. Links to external sites are particularly useful.

- **Altitude of a Triangle:** The perpendicular distance between a vertex of a triangle and the side opposite that vertex. Sometimes called the height of a triangle. Also, sometimes the line segment itself is referred to as the altitude.

- **Hinge Theorem:** If two sides of one triangle are congruent to two sides of another triangle, and the included angle of the first is larger than the included angle of the second, then the third side of the first triangle is longer than the third side of the second triangle. (Wikipedia)
- **Included Angle:** The angle between two given sides of a triangle
- **Law of Cosines:** The square of the length of any side of a triangle equals the sum of the squares of the lengths of the other two sides minus twice the product of the lengths of the other two sides and the cosine of the angle between them. (Swokowski, Cole)
- **Law of Sines:** In any triangle, the ratio of the sine of an angle to the side opposite that angle is equal to the ratio of the sine of another angle to the side opposite that angle (Swokowski, Cole)
- **Oblique Triangle:** A triangle that is not a right triangle
- **Vertex of a Triangle:** The common endpoint of the two legs that serve as the sides of a triangle

CLASSROOM ROUTINES

The importance of continuing the established classroom routines cannot be overstated. Daily routines must include such obvious activities as estimating, analyzing data, describing patterns, and answering daily questions. They should also include less obvious routines, such as how to select materials, how to use materials in a productive manner, how to put materials away, how to access classroom technology such as computers and calculators. An additional routine is to allow plenty of time for children to explore new materials before attempting any directed activity with these new materials. The regular use of routines is important to the development of students' number sense, flexibility, fluency, collaborative skills and communication. These routines contribute to a rich, hands-on standards based classroom and will support students' performances on the tasks in this unit and throughout the school year.

STRATEGIES FOR TEACHING AND LEARNING

- Students should be actively engaged by developing their own understanding.
- Mathematics should be represented in as many ways as possible by using graphs, tables, pictures, symbols and words.
- Interdisciplinary and cross curricular strategies should be used to reinforce and extend the learning activities.
- Appropriate manipulatives and technology should be used to enhance student learning.
- Students should be given opportunities to revise their work based on teacher feedback, peer feedback, and metacognition which includes self-assessment and reflection.
- Students should write about the mathematical ideas and concepts they are learning.
- Consideration of all students should be made during the planning and instruction of this unit. Teachers need to consider the following:

What level of support do my struggling students need in order to be successful with this unit?

In what way can I deepen the understanding of those students who are competent in this unit?

What real life connections can I make that will help my students utilize the skills practiced in this unit?

- A discussion of precision before this unit will help students with the standard for mathematical practice that states, “They calculate accurately and efficiently, express numerical answers with a degree of precision appropriate for the problem context.” An excessive number of decimal places may not be appropriate for real world problems. Significant digits may be discussed or the class may agree that when measurements are uncertain, 3 or 4 significant digits should be used.

EVIDENCE OF LEARNING

By the conclusion of this unit, students should be able to demonstrate the following competencies:

- Demonstrate the process of deriving the general formula for the area of a generic triangle
- Verify the Law of Sines
- Verify the Law of Cosines
- Apply the Law of Sines to solve for missing pieces of a generic triangle
- Apply the Law of Cosines to solve for missing pieces of a generic triangle
- Understand and apply the Law of Sines and the Law of Cosines to solve problems in the context of a real-world situation.

TASKS

The following tasks represent the level of depth, rigor, and complexity expected of all Pre-Calculus students. These tasks or tasks of similar depth and rigor should be used to demonstrate evidence of learning. It is important that all elements of a task be addressed throughout the learning process so that students understand what is expected of them. While some tasks are identified as a performance task, they may also be used for teaching and learning (learning task).

Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

Task Name	Task Type <i>Grouping Strategy</i>	Content Addressed
Finding a New Area Formula for Triangles	Learning Task <i>Individual/Partner Task</i>	Determining the area of triangles with the aid of trigonometry.
Proving the Law of Cosines	Learning Task <i>Individual/Partner Task</i>	Derivation of the Law of Cosines
Proving the Law of Sines	Learning Task <i>Individual/Partner Task</i>	Derivation of the Law of Sines
The Hinge Theorem	Extension Task <i>Partner/Small Group Task</i>	Investigating the Hinge Theorem using trigonometry.
Culminating Task: Combining Lots	Performance Task <i>Individual/Partner Task</i>	Utilize the standards in order to determine missing measurements.

TASKS

The following tasks represent the level of depth, rigor, and complexity expected of all Pre-Calculus students. These tasks or tasks of similar depth and rigor should be used to demonstrate evidence of learning. It is important that all elements of a task be addressed throughout the learning process so that students understand what is expected of them. While some tasks are identified as a performance task, they may also be used for teaching and learning (learning task).

FINDING A NEW AREA FORMULA FOR TRIANGLES LEARNING TASK

MGSE9-12.G.SRT.9 -Derive the formula $A = (1/2)ab \sin(C)$ for the area of a triangle by drawing an auxiliary line from a vertex perpendicular to the opposite side

Introduction:

This task is designed to help a student create a new formula for calculating the area of a triangle by combining the familiar formula for area with the trigonometric ratios that they have learned previously.

FINDING A NEW AREA FORMULA FOR TRIANGLES


A developer needs to find the area of some plots of land he is interested in buying. Each plot is owned by a different person and neither owner knows the actual area of the land. The diagram below illustrates the plots he wants to buy but he wants to know the area before buying it.

Your task is to calculate the total areas of the plots.

1. Recall that the formula for the area of a triangle is $A = \frac{1}{2}bh$


where b is the length of the base and h is the height, perpendicular to the base. Can the formula be used in this situation? Why or why not?

The area formula cannot be used with these triangles with the given information. The height must be measured perpendicular from the base, and such a measure is not given.



2. What information would be helpful in determining the area of the triangles? (Hint: Can you draw another line that would help?)

Students should recognize that they can draw a perpendicular through point C and point D. This is a good place to revisit constructions, particularly constructing a perpendicular through a given point.



3. Calculate the height of $\triangle ABC$. Find the area. Use your strategy to calculate the area of $\triangle ABD$. What is the total area of the two properties?

Since students should be in a trigonometry frame of mind, they should think of using trig ratios to find the height of the triangle. Although the task is designed to lead this way, you should encourage other methods of solution and see where that takes your class. Remember, the ultimate goal is to encourage student's mathematical thinking and problem solving.

$$\sin \angle ABC = \frac{h}{BC}$$

To find the height of ABC: $\sin 80^\circ = \frac{h}{800}$

$$h = 800 \sin 80^\circ$$

$$h = 787.85 \text{ yds}$$

Thus the area of the triangle is: $\frac{1}{2} AB \times h = \frac{1}{2} (610)(787.85) = 240293.09 \text{ yds}^2$

$$\sin \angle ABD = \frac{h}{BD}$$

To find the height of ABD: $\sin 59^\circ = \frac{h}{610}$

$$h = 610 \sin 59^\circ$$

$$h = 522.87 \text{ yds}$$

Thus the area of triangle ABC is: $\frac{1}{2} \overline{AB} \times h = \frac{1}{2} (610)(522.87) = 159475.35 \text{ yds}^2$

The total area of the two plots is: $240293.09 + 159475.35 = 399768.44 \text{ yds}^2$

Or more realistically about 399800 yds²

4. After working on this problem, would it be possible to generalize this method for use on any triangle?

Construct any triangle. Label the angles A, B, and C. Label the side opposite from A with a , the side opposite B with b , and the side opposite C with c .

Construct any altitude from a vertex.

Now calculate the height of your triangle.

This is perhaps the most important part of the task. Here the student will take a concrete idea (area) and generalize it to fit any oblique triangle. Do not emphasize memorizing the formula, as that is a low functioning task. Instead, challenge students to understand where the formula comes from and be able to construct the formula from scratch.

Now that you know the height of the triangle, you can write a general formula for the area of any triangle. Use $A = \frac{1}{2}bh$ as a starting point.

Students should have any one of the following 3 formulas:

$$A = \frac{1}{2}ac(\sin B)$$

$$A = \frac{1}{2}ab(\sin C)$$

$$A = \frac{1}{2}bc(\sin A)$$

In addition, students should be able to generalize all three of these formulas by understanding that the area of a triangle equals one-half the product of the lengths of any two sides and the sine of the included angle.

5. Compare your formula with another student. Did you get the same thing? Could you both be right?

This is where students justify their answers and explain to other students their methods and findings. Students should arrive at the conclusion that any of the three formulas above are correct.

6. Would it be possible to develop a third formula for the area? If so, find it. If not, explain why not.

The pairs should then attempt to find the third iteration of the above formulas, depending on the formulas they found already.


FINDING A NEW AREA FORMULA FOR TRIANGLES LEARNING TASK

A developer needs to find the area of some plots of land he is interested in buying. Each plot is owned by a different person and neither owner knows the actual area of the land. The diagram below illustrates the plots he wants to buy but he wants to know the area before buying it.

Your task is to calculate the total areas of the plots.

1. Recall that the formula for the area of a triangle is $A = \frac{1}{2}bh$

where b is the length of the base and h is the height, perpendicular to the base. Can the formula be used in this situation? Why or why not?



2. What information would be helpful in determining the area of the triangles? (Hint: Can you draw another line that would help?)
3. Calculate the height of $\triangle ABC$. Find the area. Use your strategy to calculate the area of $\triangle ABD$. What is the total area of the two properties?

4. After working on this problem, would be possible to generalize this method for use on any triangle?

Construct any triangle. Label the angles A, B, and C. Label the side opposite from A with a, the side opposite B with b, and the side opposite C with c.

Construct any altitude from a vertex.

Now calculate the height of your triangle.

Now that you know the height of the triangle, you can write a general formula for the area of any triangle. Use $A = \frac{1}{2}bh$ as a starting point.

5. Compare your formula with another student. Did you get the same thing? Could you both be right?
6. Would it be possible to develop a third formula for the area? If so, find it. If not, explain why not.

PROVING THE LAW OF COSINES

MGSE9-12.G.SRT.10 Prove the Laws of Sines and Cosines and use them to solve problems.

MGSE9-12.G.SRT.11 Understand and apply the Law of Sines and the Law of Cosines to find unknown measurements in right and non-right triangles

Introduction

The purpose of this task is to guide students through the derivation of the Law of Cosines. The teacher should spend the extra time and effort in helping students understand the conceptual foundation for the Law of Cosines, and not just memorizing the formula. A few problems have been provided at the end of the task, but the classroom teacher should provide ample opportunities to practice and improve fluency.

PROVING THE LAW OF COSINES


During a baseball game an outfielder caught a ball hit to dead center field, 400 feet from home plate. If the distance from home plate to first base is 90 feet, how far does the outfielder have to throw the ball to get it to first base?

1. Model the problem with a picture. Be sure to label information that you know.




2. Do you have enough information to solve the problem? If not, what is missing?

We know two sides of the triangle. We also know the angle at home base is 45° (The 'diamond' is a square. The diagonal of a square bisects the angle at the vertex.)



Typically, you have solved triangles that are right triangles. This is a case where we do not have a right triangle to solve. We know two sides and one included angle. In this task, you will develop a method for solving triangles like this using trigonometry. We will come back to the baseball example later. For now, consider the triangle below. Follow these steps to derive a way to solve for c knowing just that much information. For this example, assume we know measurements for segments a , b , and angle C .



3. What does segment h represent? What are its properties and what does it do to the large triangle?

h is the altitude of the triangle. It is perpendicular to the base and divides the larger triangle into two right triangles.

Comment:

Get students to realize that the altitude does not necessarily bisect the angle!

4. Write an equation that represents c^2 . Explain the method you used.

According to the Pythagorean Theorem:

$$c^2 = h^2 + (a - x)^2 = h^2 + a^2 - 2ax + x^2$$

5. Now write an equation that represents h^2 in terms of b and x . Substitute this expression into the expression you wrote in #4. Expand and simplify.

According to the Pythagorean Theorem:

$$h^2 = b^2 - x^2$$

When you substitute this into the c^2 equation you get:

$$c^2 = b^2 + a^2 - 2ax$$

6. Now write an expression that represents x in terms of the angle C . Substitute this expression into the equation you wrote in #5. Simplify completely.

Students should use the cosine ratio here to arrive at:

$$\cos C = \frac{x}{b}$$
$$x = b(\cos C)$$

Substituting this in gives: $c^2 = b^2 + a^2 - 2ab(\cos C)$, one of the versions of the Law of Cosines.

Your answer to #6 is one of three formulas that make up the **Law of Cosines**. Each of the formulas can be derived in the same way you derived this one by working with each vertex and the other heights of the triangle.

Law of Cosines

Let a , b , and c be the lengths of the legs of a triangle opposite angles A , B , and C . Then,

$$a^2 = b^2 + c^2 - 2bc(\cos A)$$

$$b^2 = a^2 + c^2 - 2ac(\cos B)$$

$$c^2 = a^2 + b^2 - 2ab(\cos C)$$

These formulas can be used to solve for unknown lengths and angles in a triangle.

Students should also be able to verbally articulate the generalization of these Laws. It should go something like this: The square of the length of any side of a triangle is equal to the sum of the squares of the other two sides minus two times the product of the lengths of the other two sides and the cosine of their included angle.

7. Solve the baseball problem at the beginning of this task using the Law of Cosines.

In this problem $a = 90$, $b = 400$ and $C = 45^\circ$.

Using the Law of Cosines:

$$c^2 = a^2 + b^2 - 2ab\cos C$$

$$c^2 = 90^2 + 400^2 - 2(90)(400)\cos 45^\circ$$

$$c \approx 342$$

The distance is about 342 feet from dead centerfield to 1st base.

Here are a few problems to help you apply the Law of Cosines.

8. Two airplanes leave an airport, and the angle between their flight paths is 40° . An hour later, one plane has traveled 300 miles while the other has traveled 200 miles. How far apart are the planes at this time?

$$a^2 = b^2 + c^2 - 2bccosA$$

Let a be the missing side, $b = 300$, $c = 200$ and


$A = 40$.

$$a^2 = 300^2 + 200^2 - 2(300)(200)\cos(40)$$

$$a^2 = 38074.66683$$

$$a \approx 195.13 \text{ miles}$$

The planes are approximately 195 miles apart.



Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

9. A triangle has sides of 8 and 7 and the angle between these sides is 35° . Solve the triangle. (Find all missing angles and sides.)

$$a^2 = b^2 + c^2 - 2bccosA$$

$$b = 7, c = 8 \text{ and } A = 35.$$

$$a^2 = 7^2 + 8^2 - 2(7)(8)cos(35)$$

$$a^2 \approx 21.255$$

$$a \approx 4.61$$

To find angle C

$$c^2 = a^2 + b^2 - 2abcosC$$


$$8^2 = 4.61^2 + 7^2 - 2(4.61)(7)(cosC)$$

$$64 = 70.2521 - 64.54(cosC)$$

$$0.0969 = cos C$$

$$84.44 = C$$

$$B = 180 - 84.44 - 35 = 60.56$$



10. Three soccer players are practicing on a field. The triangle they create has side lengths of 18, 14, and 15 feet. At what angles are they standing from each other?

To find angle C

$$c^2 = a^2 + b^2 - 2abcosC$$

$$15^2 = 14^2 + 18^2 - 2(14)(18)(cosC)$$

$$225 = 520 - 504(cosC)$$

$$0.5853 = cos C$$

$$54.17^\circ = C$$

To find angle B

$$b^2 = a^2 + c^2 - 2accosB$$

$$18^2 = 14^2 + 15^2 - 2(14)(15)(cosB)$$


$$324 = 421 - 420(cosC)$$

$$0.231 = cos C$$

$$76.65^\circ = C$$

To find angle A

$$A = 180 - 54.17 - 76.65 = 49.18^\circ$$



Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

11. Is it possible to know two sides of a triangle and the included angle and not be able to solve for the third side?

Using the Law of Cosines, you will always be able to find the third side of the triangle.

PROVING THE LAW OF COSINES


During a baseball game an outfielder caught a ball hit to dead center field, 400 feet from home plate. If the distance from home plate to first base is 90 feet, how far does the outfielder have to throw the ball to get it to first base?

1. Model the problem with a picture. Be sure to label information that you know.



2. Do you have enough information to solve the problem? If not, what is missing?

Typically, you have solved triangles that are right triangles. This is a case where we do not have a right triangle to solve. We know two sides and one included angle. In this task, you will develop a method for solving triangles like this using trigonometry. We will come back to the baseball example later. For now, consider the triangle below. Follow these steps to derive a way to solve for c knowing just that much information. For this example, assume we know measurements for segments a , b , and angle C .



3. What does segment h represent? What are its properties and what does it do to the large triangle?

4. Write an equation that represents c^2 . Explain the method you used.
5. Now write an equation that represents h^2 in terms of b and x . Substitute this expression into the expression you wrote in #4. Expand and simplify.
6. Now write an expression that represents x in terms of the angle C . Substitute this expression into the equation you wrote in #5. Simplify completely.

Your answer to #6 is one of three formulas that make up the **Law of Cosines**. Each of the formulas can be derived in the same way you derived this one by working with each vertex and the other heights of the triangle.

Law of Cosines

Let a , b , and c be the lengths of the legs of a triangle opposite angles A , B , and C . Then,

$$a^2 = b^2 + c^2 - 2bc(\cos A)$$

$$b^2 = a^2 + c^2 - 2ac(\cos B)$$

$$c^2 = a^2 + b^2 - 2ab(\cos C)$$

These formulas can be used to solve for unknown lengths and angles in a triangle.

7. Solve the baseball problem at the beginning of this task using the Law of Cosines.

GSE Pre-Calculus • Unit 3

11. Is it possible to know two sides of a triangle and the included angle and not be able to solve for the third side?

PROVING THE LAW OF SINES

MGSE9-12.G.SRT.10 Prove the Laws of Sines and Cosines and use them to solve problems.


MGSE9-12.G.SRT.11 Understand and apply the Law of Sines and the Law of Cosines to find unknown measurements in right and non-right triangles

Introduction

The purpose of this task is to guide students through the process of deriving the Law of Sines. Emphasis should be placed on conceptual understanding rather than memorizing a formula.

PROVING THE LAW OF SINES

When using the Law of Cosines it is necessary to know the angle included between two sides. However, there are times the angle that is known is not the angle included between two known sides. And there are other times where we might know two angles and only one side. Consider the following triangle: The angle labels should be closer to the same size as the side labels.



1. Construct an altitude through C . Label it h .
2. Write an equation for h in terms of angle B . Also write an equation for h in terms of angle A .

Students should be familiar with altitudes from the last task.

$$\sin B = \frac{\text{opp}}{\text{hyp}} = \frac{h}{a} \qquad \sin A = \frac{\text{opp}}{\text{hyp}} = \frac{h}{b}$$

$$h = a \sin B \qquad h = b \sin A$$

3. Write a new equation relating the two equations together.

By noticing that $h=h$ and then substituting in, we get:

$$a \sin B = b \sin A$$

$$\frac{a}{\sin A} = \frac{b}{\sin B}$$

4. Now construct another altitude through B . Label it j .
5. Write an equation for j in terms of angle A . Also write an equation for j in terms of angle C .

$$\sin A = \frac{\text{opp}}{\text{hyp}} = \frac{j}{c} \qquad \sin C = \frac{\text{opp}}{\text{hyp}} = \frac{j}{a}$$

$$j = c \sin A \qquad j = a \sin C$$

6. Write a new equation relating the two equations together.

By noticing that $j=j$ and then substituting in, we get:

$$c \sin A = a \sin C$$

$$\frac{a}{\sin A} = \frac{c}{\sin C}$$

7. Using the Transitive Property of Equality, write an equation relating the equations in #3 and #6.

Thus by the transitive property: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$


What you have arrived at is known as the **Law of Sines**.

Law of Sines

Let a , b , and c be the lengths of the legs of a triangle opposite angles A , B , and C . Then,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Here is an example of applying the Law of Sines. In the triangle to the right the measurements for angle B and A are missing and the length of AC . Follow the steps below to solve for the measure using the Law of Sines.



Students should focus on understanding and being able to explain the definition of the law of sines. The ratio of the sine of an angle to the side opposite the angle is equal to the ratio of the sine of another angle to the side opposite that angle.

This is the information that we know so far.

$$\frac{6.7}{\sin 33} = \frac{5.4}{\sin A} = \frac{b}{\sin B}$$

- Solve the first two ratios to find A .

$$5.4 \sin 33 = 6.7 \sin A$$

$$\sin A = \frac{5.4 \sin 33}{6.7} = .4389$$

$$A = 26.04$$

- Find angle B . (Remember that the sum of the interior angles in a triangle is always 180 degrees)

$$B = 180 - 26 - 33 = 121^\circ$$

Comment:

If students use the Law of Sines to solve this, they will get the wrong answer. This is a good opportunity to discuss the range of the \sin^{-1} function and why it isn't a good idea to use the law of sines to find the largest angle in a triangle.

Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

- Use either the first or second ratio with the last ratio to solve for side b .

$$\frac{6.7}{\sin 33} = \frac{b}{\sin 121}$$

$$\frac{6.7}{\sin 33} * \sin 121 = b$$

$$b = 10.55$$

Here are some problems that will help you practice applying the Law of Sines. Begin each problem by drawing a picture to model the situation. Then solve the problem.


- A surveyor is near a river and wants to calculate the distance across the river. He measures the angle between his observations of two points on the shore, one on his side and one on the other side, to be 28° . The distance between him and the point on his side of the river can be measured and is 300 feet. The angle formed by him, the point on his side of the river, and the point on the opposite side of the river is 128° . What is the distance across the river? Remember that the distance across the river should be the shortest distance.

Angle $c = 24^\circ$

$$\frac{300}{\sin 24^\circ} = \frac{d}{\sin 28^\circ}$$

$$\frac{300 \sin 28^\circ}{\sin 24^\circ} = d$$

$$d = 346 \text{ feet}$$



Georgia Department of Education


Georgia Standards of Excellence Framework

GSE Pre-Calculus • Unit 3

9. Two people are observing a hot air balloon from different places on level ground. The angles of elevation are 20° and 50° . The two people are 7 miles apart and the balloon is between them. How high is the balloon off the ground?

First find the distance from the balloon to the person with the 50 degree line of sight.

$$\frac{7}{\sin 110^\circ} = \frac{a}{\sin 20^\circ}$$
$$a = 2.548$$



Now use side a to find the height.


$$\sin 50^\circ = \frac{h}{2.548}$$

$$h = 1.95$$

The balloon is about 1.95 miles off the ground

PROVING THE LAW OF SINES

When using the Law of Cosines it is necessary to know the angle included between two sides. However, there are times the angle that is known is not the angle included between two known sides. And there are other times where we might know two angles and only one side. Consider the following triangle:



1. Construct an altitude through C . Label it h .
2. Write an equation for h in terms of angle B . Also write an equation for h in terms of angle A .
3. Write a new equation relating the two equations together.
4. Now construct another altitude through B . Label it j .

Georgia Department of Education

Georgia Standards of Excellence Framework

GSE Pre-Calculus • Unit 3

5. Write an equation for j in terms of angle A . Also write an equation for j in terms of angle C .
6. Write a new equation relating the two equations together.
7. Using the Transitive Property of Equality, write an equation relating the equations in #3 and #6.


What you have arrived at is known as the **Law of Sines**.

Law of Sines

Let a , b , and c be the lengths of the legs of a triangle opposite angles A , B , and C . Then,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Here is an example of applying the Law of Sines. In the triangle to the right the measurements for angle B and A are missing and the length of AC . Follow the steps below to solve for the measure using the Law of Sines.



This is the information that we know so far.

$$\frac{6.7}{\sin 33} = \frac{5.4}{\sin A} = \frac{b}{\sin B}$$

- Solve the first two ratios to find A .
- Find angle B . (Remember that the sum of the interior angles in a triangle is always 180 degrees)
- Use either the first or second ratio with the last ratio to solve for side b . Here are some problems that will help you practice applying the Law of Sines. Begin each problem by drawing a picture to model the situation. Then solve the problem.

Georgia Department of Education

Georgia Standards of Excellence Framework

GSE Pre-Calculus • Unit 3

8. A surveyor is near a river and wants to calculate the distance across the river. He measures the angle between his observations of two points on the shore, one on his side and one on the other side, to be 28° . The distance between him and the point on his side of the river can be measured and is 300 feet. The angle formed by him, the point on his side of the river, and the point on the opposite side of the river is 128° . What is the distance across the river? Remember that the distance across the river should be the shortest distance.
9. Two people are observing a hot air balloon from different places on level ground. The angles of elevation are 20° and 50° . The two people are 7 miles apart and the balloon is between them. How high is the balloon off the ground?

THE HINGE THEOREM

MGSE9-12.G.SRT.10 Prove the Laws of Sines and Cosines and use them to solve problems.

MGSE9-12.G.SRT.11 Understand and apply the Law of Sines and the Law of Cosines to find unknown measurements in right and non-right triangles


Introduction:

Although the Hinge Theorem is not part of the standards, this task makes a connection to concepts already known by the student and the Law of Sines. This exploration works well with construction software if it is available.

THE HINGE THEOREM

From your previous math experience you know that the measures of two sides and a non-included angle will not necessarily work together to create a triangle. The Hinge Theorem is a geometric theorem that focuses on this idea. You may have explored this idea when you studied congruent triangles.

Consider the two triangles below. Given sides of 7 cm and 4.2 cm with a non-included angle of 30° , there are two triangles that can be created. This is why angle-side-side is not a congruency theorem for triangles.



In trigonometry we consider this to be the ambiguous case for solving triangles.


Looking at the two triangles above, why do you think this theorem might be called the Hinge Theorem?

BC seems to swing back and forth (like a hinge) to create the 2 different triangles.

What is the relationship between the measure of $\angle ABC$ in the first figure and $m\angle ABC$ in the second figure?

They are supplementary.

Consider triangle ABC to the right. Note that $m\angle A = 30^\circ$, $AC = 7$ cm and $BC = 3.5$ cm. Are there still 2 possible triangles? Explain your answer.



No, since the angle opposite the 30° angle is half the adjacent side, this must be a 30° - 60° - 90° triangle. BC is too short to “swing”.

Redraw triangle ABC so that $m\angle A = 30^\circ$, $AC = 7$ cm and $BC = 9$ cm. Are there still 2 triangles? Explain your answer.


No, BC is too long to swing to the other side of AC and still have a 30° angle.

Redraw triangle ABC so that $m\angle A = 30^\circ$, $AC = 7$ cm and $BC = 2$ cm. Are there still 2 triangles? Explain your answer.

No, since BC is shorter than 3.5, it is too short to reach line AB , so no triangle can be created.

1. Order the sides of the triangle to the right from longest to shortest. (The figure is NOT drawn to scale.)

$$a > c > b$$



2. How did you decide the order of the sides?

The longest side is opposite the largest angle.

The shortest side is always opposite the smallest angle.

Georgia Department of Education

Georgia Standards of Excellence Framework

GSE Pre-Calculus • Unit 3

3. Sketch the information given and decide if it is possible to create a triangle with the given information. Explain your answer.

Triangle 1: $A = 40^\circ$, $a = 8$ cm and $b = 5$ cm.

To make a right triangle the side would need to be 5.14 cm. The side of 5 is too short so this will not make a triangle.

Triangle 2: $A = 150^\circ$, $a = 5$ cm and $b = 8$ cm

The side opposite the obtuse angle should be the longest side in the triangle. But it's shorter than the other given side. Therefore, you cannot make a triangle with these measures.

Be sure to take time to analyze the data you are given when you are solving a triangle to determine if you might have a set of data that will not yield a triangle. This could save you a lot of work!

Remember that the calculator only yields inverse sine values between -90° and 90° , so it will never let you know if there are obtuse angles in your triangles.

4. Solve the following triangle. Determine if there is no solution, one solution or two solutions.

Measure of angle $A = 35^\circ$, $a = 10$ cm and $b = 16$ cm

*If this were going to make a right triangle, a would have to be equal to $16\sin 35^\circ$.
 $16\sin 35^\circ = 9.18$*

Since a is greater than that but still less than 16, there will be two solutions.

$$\frac{10}{\sin 35} = \frac{16}{\sin B}$$

$$B = 66.6^\circ \text{ or } 113.4^\circ$$

$$C = 180 - 66.6 - 35 = 78.4^\circ \quad \text{or} \quad C = 180 - 113.4 - 35 = 31.6^\circ$$

$$\frac{10}{\sin 35} = \frac{c}{\sin 78.4}$$
$$c = 17.1$$

$$\frac{10}{\sin 35} = \frac{c}{\sin 31.6}$$
$$c = 9.13$$

Georgia Department of Education

Georgia Standards of Excellence Framework

GSE Pre-Calculus • Unit 3

5. A ship traveled 60 miles due east and then adjusted its course 15 degrees northward. After traveling for a while the ship turned back towards port. The ship arrived in port 139 miles later. How far did the ship travel on the second leg of the journey? What angle did the ship turn through when it headed back to port?

$$\frac{139}{\sin 165} = \frac{60}{\sin C}$$


$$C = 6.4 \text{ degrees}$$

$$A = 180 - 165 - 6.4 = 8.6$$

$$\frac{139}{\sin 165} = \frac{a}{\sin 8.6}$$

A



$$a = 80.2 \text{ miles}$$



THE HINGE THEOREM

From your previous math experience you know that the measures of two sides and a non-included angle will not necessarily work together to create a triangle. The Hinge Theorem is a geometric theorem that focuses on this idea. You may have explored this idea when you studied congruent triangles.

Consider the two triangles below. Given sides of 7 cm and 4.2 cm with a non-included angle of 30° , there are two triangles that can be created. This is why angle-side-side is not a congruency theorem for triangles.




In trigonometry we consider this to be the ambiguous case for solving triangles.

Looking at the two triangles above, why do you think this theorem might be called the Hinge Theorem?

What is the relationship between the measure of $\angle ABC$ in the first figure and $m\angle ABC$ in the second figure?

Consider triangle ABC to the right. Note that $m\angle A = 30^\circ$, $AC = 7$ cm and $BC = 3.5$ cm. Are there still 2 possible triangles? Explain your answer.




Redraw triangle ABC so that $m\angle A = 30^\circ$, $AC = 7$ cm and $BC = 9$ cm. Are there still 2 triangles? Explain your answer.

Redraw triangle ABC so that $m\angle A = 30^\circ$, $AC = 7$ cm and $BC = 2$ cm. Are there still 2 triangles? Explain your answer.

Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

1. Order the sides of the triangle to the right from longest to shortest. (The figure is NOT drawn to scale.



2. How did you decide the order of the sides?

3. Sketch the information given and decide if it is possible to create a triangle with the given information. Explain your answer.

Triangle 1: $A = 40^\circ$, $a = 8$ cm and $b = 5$ cm.

Triangle 2: $A = 150^\circ$, $a = 5$ cm and $b = 8$ cm

Be sure to take time to analyze the data you are given when you are solving a triangle to determine if you might have a set of data that will not yield a triangle. This could save you a lot of work!

Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

Remember that the calculator only yields inverse sine values between -90° and 90° , so it will never let you know if there are obtuse angles in your triangles.

4. Solve the following triangle. Determine if there is no solution, one solution or two solutions.

Measure of angle $A = 35^\circ$, $a = 10$ cm and $b = 16$ cm

5. A ship traveled 60 miles due east and then adjusted its course 15 degrees northward. After traveling for a while the ship turned back towards port. The ship arrived in port 139 miles later. How far did the ship travel on the second leg of the journey? What angle did the ship turn through when it headed back to port?

CULMINATING TASK: COMBINING LOTS


Standards:

MGSE9-12.G.SRT.9 Derive the formula $A = (1/2)ab \sin(C)$ for the area of a triangle by drawing an auxiliary line from a vertex perpendicular to the opposite side.

MGSE9-12.G.SRT.10 Prove the Laws of Sines and Cosines and use them to solve problems.

MGSE9-12.G.SRT.11 Understand and apply the Law of Sines and the Law of Cosines to find unknown measurements in right and non-right triangles (e.g., surveying problems, resultant forces).

An antique map was found in the attic of a local courthouse. It shows some measurements from a local farm that was divided into 5 parts. Some of the measurements have faded with age, so you must find the remaining measurements, as well as calculate the total area and perimeter of the outside of the property. All lengths are measured in miles.



From the Pythagorean Theorem, $\overline{AC} = 5$

Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

From the Law of Cosines, $\overline{AD}^2 = 5^2 + 3.1623^2 - 2(5)(3.1623)\cos 71.5651^\circ = 25$
 $\overline{AD} = 5$

From the Law of Sines: $\frac{7.8102}{\sin \angle EDA} = \frac{5}{\sin 38.8845^\circ}$
 $\angle EDA = 101.3117^\circ$ *(The obtuse solution)*

From subtraction: $\angle EAD = 180^\circ - (38.8845^\circ + 101.3117^\circ) = 39.8038^\circ$

From Law of Sines: $\frac{\overline{ED}}{\sin 39.8038^\circ} = \frac{5}{\sin 38.8845^\circ}$
 $\overline{ED} = 5.0988$

From Law of Sines: $\frac{7.8102}{\sin 120.9638^\circ} = \frac{5.831}{\sin \angle AEF}$
 $\angle AEF = 39.8062^\circ$

By Subtraction: $\angle EAC = 180^\circ - (39.8062^\circ + 120.9638^\circ) = 19.23^\circ$

From Law of Sines: $\frac{7.8102}{\sin 120.9638^\circ} = \frac{\overline{EF}}{\sin 19.23^\circ}$
 $\overline{EF} = 3$

From Subtraction: $\angle GAF = 90^\circ - (39.8038^\circ + 19.23^\circ) = 30.9662^\circ$

From Subtraction: $\angle AFG = 180^\circ - 120.9638^\circ = 59.0362^\circ$

From Law of Sines: $\frac{5.831}{\sin 90^\circ} = \frac{\overline{AG}}{\sin 59.0362^\circ}$
 $\overline{AG} = 5$

From Law of Sines: $\frac{5.831}{\sin 90^\circ} = \frac{\overline{FG}}{\sin 30.9638^\circ}$
 $\overline{FG} = 3$

It should be noted that the methods here are only one possible way to find the missing parts. Encourage multiple solutions to this problem

Perimeter = 3+4+3.1623+5.0988+3+3+5=26.2611 km

Georgia Department of Education
Georgia Standards of Excellence Framework
GSE Pre-Calculus • Unit 3

Area: $\Delta ABC = \frac{1}{2}(3)(4) = 6$
 $\Delta ADC = \frac{1}{2}(5)(3.1623) \sin 71.5651 = 7.5$
 $\Delta ADE = \frac{1}{2}(7.8102)(5.0988) \sin 38.8845 = 12.5$
 $\Delta AEF = \frac{1}{2}(3)(5.831) \sin 120.9638 = 7.5$
 $\Delta AGF = \frac{1}{2}(5)(3) = 7.5$

Total Area: $6 + 7.5 + 12.5 + 7.5 + 7.5 = 41 \text{ km}^2$

CULMINATING TASK: COMBINING LOTS

An antique map was found in the attic of a local courthouse. It shows some measurements from a local farm that was divided into 5 parts. Some of the measurements have faded with age, so you must find the remaining measurements, as well as calculate the total area and perimeter of the outside of the property. All lengths are measured in miles.

