

## CALCULUS BC

## SECTION II, Part A

Time—30 minutes

Number of questions—2

**A GRAPHING CALCULATOR IS REQUIRED FOR THESE QUESTIONS.**

1. People enter a line for an escalator at a rate modeled by the function  $r$  given by

$$r(t) = \begin{cases} 44\left(\frac{t}{100}\right)^3\left(1 - \frac{t}{300}\right)^7 & \text{for } 0 \leq t \leq 300 \\ 0 & \text{for } t > 300, \end{cases}$$

where  $r(t)$  is measured in people per second and  $t$  is measured in seconds. As people get on the escalator, they exit the line at a constant rate of 0.7 person per second. There are 20 people in line at time  $t = 0$ .


- (a) How many people enter the line for the escalator during the time interval  $0 \leq t \leq 300$  ?
- (b) During the time interval  $0 \leq t \leq 300$ , there are always people in line for the escalator. How many people are in line at time  $t = 300$  ?
- (c) For  $t > 300$ , what is the first time  $t$  that there are no people in line for the escalator?
- (d) For  $0 \leq t \leq 300$ , at what time  $t$  is the number of people in line a minimum? To the nearest whole number, find the number of people in line at this time. Justify your answer.
-

## CALCULUS BC

## SECTION II, Part B

Time—1 hour

Number of questions—4

**NO CALCULATOR IS ALLOWED FOR THESE QUESTIONS.**Graph of  $g$ 

3. The graph of the continuous function  $g$ , the derivative of the function  $f$ , is shown above. The function  $g$  is piecewise linear for  $-5 \leq x < 3$ , and  $g(x) = 2(x - 4)^2$  for  $3 \leq x \leq 6$ .
- (a) If  $f(1) = 3$ , what is the value of  $f(-5)$ ?
- (b) Evaluate  $\int_1^6 g(x) \, dx$ .
- (c) For  $-5 < x < 6$ , on what open intervals, if any, is the graph of  $f$  both increasing and concave up? Give a reason for your answer.
- (d) Find the  $x$ -coordinate of each point of inflection of the graph of  $f$ . Give a reason for your answer.
-

$t$ (years)	2	3	5	7	10
$H(t)$ (meters)	1.5	2	6	11	15


4. The height of a tree at time  $t$  is given by a twice-differentiable function  $H$ , where  $H(t)$  is measured in meters and  $t$  is measured in years. Selected values of  $H(t)$  are given in the table above.
- (a) Use the data in the table to estimate  $H'(6)$ . Using correct units, interpret the meaning of  $H'(6)$  in the context of the problem.
- (b) Explain why there must be at least one time  $t$ , for  $2 < t < 10$ , such that  $H'(t) = 2$ .
- (c) Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate the average height of the tree over the time interval  $2 \leq t \leq 10$ .
- (d) The height of the tree, in meters, can also be modeled by the function  $G$ , given by  $G(x) = \frac{100x}{1+x}$ , where  $x$  is the diameter of the base of the tree, in meters. When the tree is 50 meters tall, the diameter of the base of the tree is increasing at a rate of 0.03 meter per year. According to this model, what is the rate of change of the height of the tree with respect to time, in meters per year, at the time when the tree is 50 meters tall?
-

CALCULUS BC  
SECTION II, Part B

Time—60 minutes

Number of problems—4

No calculator is allowed for these problems.

Graph of  $f$ 

3. The figure above shows the graph of the piecewise-linear function  $f$ . For  $-4 \leq x \leq 12$ , the function  $g$  is defined by  $g(x) = \int_{-2}^x f(t) dt$ .
- (a) Does  $g$  have a relative minimum, a relative maximum, or neither at  $x = 10$ ? Justify your answer.
  - (b) Does the graph of  $g$  have a point of inflection at  $x = 4$ ? Justify your answer.
  - (c) Find the absolute minimum value and the absolute maximum value of  $g$  on the interval  $-4 \leq x \leq 12$ . Justify your answers.
  - (d) For  $-4 \leq x \leq 12$ , find all intervals for which  $g(x) \leq 0$ .
-

4. Consider the differential equation  $\frac{dy}{dx} = x^2 - \frac{1}{2}y$ .

(a) Find  $\frac{d^2y}{dx^2}$  in terms of  $x$  and  $y$ .


(b) Let  $y = f(x)$  be the particular solution to the given differential equation whose graph passes through the point  $(-2, 8)$ . Does the graph of  $f$  have a relative minimum, a relative maximum, or neither at the point  $(-2, 8)$ ? Justify your answer.

(c) Let  $y = g(x)$  be the particular solution to the given differential equation with  $g(-1) = 2$ . Find

$$\lim_{x \rightarrow -1} \left( \frac{g(x) - 2}{3(x+1)^2} \right).$$
 Show the work that leads to your answer.

(d) Let  $y = h(x)$  be the particular solution to the given differential equation with  $h(0) = 2$ . Use Euler's method, starting at  $x = 0$  with two steps of equal size, to approximate  $h(1)$ .

---



5. The inside of a funnel of height 10 inches has circular cross sections, as shown in the figure above. At height  $h$ , the radius of the funnel is given by  $r = \frac{1}{20}(3 + h^2)$ , where  $0 \leq h \leq 10$ . The units of  $r$  and  $h$  are inches.
- (a) Find the average value of the radius of the funnel.
  - (b) Find the volume of the funnel.
  - (c) The funnel contains liquid that is draining from the bottom. At the instant when the height of the liquid is  $h = 3$  inches, the radius of the surface of the liquid is decreasing at a rate of  $\frac{1}{5}$  inch per second. At this instant, what is the rate of change of the height of the liquid with respect to time?
-

**CALCULUS BC**  
**SECTION II, Part A**

Time—30 minutes

Number of problems—2

**A graphing calculator is required for these problems.**

1. The rate at which rainwater flows into a drainpipe is modeled by the function  $R$ , where  $R(t) = 20 \sin\left(\frac{t^2}{35}\right)$  cubic feet per hour,  $t$  is measured in hours, and  $0 \leq t \leq 8$ . The pipe is partially blocked, allowing water to drain out the other end of the pipe at a rate modeled by  $D(t) = -0.04t^3 + 0.4t^2 + 0.96t$  cubic feet per hour, for  $0 \leq t \leq 8$ . There are 30 cubic feet of water in the pipe at time  $t = 0$ .
- (a) How many cubic feet of rainwater flow into the pipe during the 8-hour time interval  $0 \leq t \leq 8$ ?
- (b) Is the amount of water in the pipe increasing or decreasing at time  $t = 3$  hours? Give a reason for your answer.
- (c) At what time  $t$ ,  $0 \leq t \leq 8$ , is the amount of water in the pipe at a minimum? Justify your answer.
- (d) The pipe can hold 50 cubic feet of water before overflowing. For  $t > 8$ , water continues to flow into and out of the pipe at the given rates until the pipe begins to overflow. Write, but do not solve, an equation involving one or more integrals that gives the time  $w$  when the pipe will begin to overflow.
-

**CALCULUS BC**  
**SECTION II, Part B****Time—60 minutes****Number of problems—4****No calculator is allowed for these problems.**

$t$ (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

3. Johanna jogs along a straight path. For  $0 \leq t \leq 40$ , Johanna's velocity is given by a differentiable function  $v$ . Selected values of  $v(t)$ , where  $t$  is measured in minutes and  $v(t)$  is measured in meters per minute, are given in the table above.

(a) Use the data in the table to estimate the value of  $v'(16)$ .

(b) Using correct units, explain the meaning of the definite integral  $\int_0^{40} |v(t)| dt$  in the context of the problem.

Approximate the value of  $\int_0^{40} |v(t)| dt$  using a right Riemann sum with the four subintervals indicated in the table.

(c) Bob is riding his bicycle along the same path. For  $0 \leq t \leq 10$ , Bob's velocity is modeled by  $B(t) = t^3 - 6t^2 + 300$ , where  $t$  is measured in minutes and  $B(t)$  is measured in meters per minute.

Find Bob's acceleration at time  $t = 5$ .


(d) Based on the model  $B$  from part (c), find Bob's average velocity during the interval  $0 \leq t \leq 10$ .

---



4. Consider the differential equation  $\frac{dy}{dx} = 2x - y$ .

(a) On the axes provided, sketch a slope field for the given differential equation at the six points indicated.



- (b) Find  $\frac{d^2y}{dx^2}$  in terms of  $x$  and  $y$ . Determine the concavity of all solution curves for the given differential equation in Quadrant II. Give a reason for your answer.
- (c) Let  $y = f(x)$  be the particular solution to the differential equation with the initial condition  $f(2) = 3$ . Does  $f$  have a relative minimum, a relative maximum, or neither at  $x = 2$ ? Justify your answer.
- (d) Find the values of the constants  $m$  and  $b$  for which  $y = mx + b$  is a solution to the differential equation.
-

5. Consider the function  $f(x) = \frac{1}{x^2 - kx}$ , where  $k$  is a nonzero constant. The derivative of  $f$  is given by

$$f'(x) = \frac{k - 2x}{(x^2 - kx)^2}.$$

- (a) Let  $k = 3$ , so that  $f(x) = \frac{1}{x^2 - 3x}$ . Write an equation for the line tangent to the graph of  $f$  at the point whose  $x$ -coordinate is 4.
- (b) Let  $k = 4$ , so that  $f(x) = \frac{1}{x^2 - 4x}$ . Determine whether  $f$  has a relative minimum, a relative maximum, or neither at  $x = 2$ . Justify your answer.
- (c) Find the value of  $k$  for which  $f$  has a critical point at  $x = -5$ .
- (d) Let  $k = 6$ , so that  $f(x) = \frac{1}{x^2 - 6x}$ . Find the partial fraction decomposition for the function  $f$ .  
Find  $\int f(x) dx$ .
-