AP STATISTICS REVIEW: FIRST SEMESTER

Name	:				Date:			
A. M	ultiple choice	e.						
1.		ouped into the f This data is	following class	es: Subcompac	t, compact, midsize,	and		
	A. ordinal	B. nominal	C. ratio	D. interval	E. binomial			
2.		-	-	. The machine operator checks every tenth et. This is an example of				
	A. random	sampling	B. systema	tic sampling	C. cluster sampli	ng		
	D. stratifie	d sampling	E. quota sa	mpling.				
3.	What is the		ouying two of		wo defective calculat and ending up with	ors.		
	A. 20%	B. 25%	C. 33%	D. 36%	E. 50%			
4.	award prize		00, and \$200 to	the lucky win	\$1.00 per ticket and ners. What is the			
	A. Loss of	\$1 B. I	Loss of \$0.66	C. I	Loss of \$0.50			
	D. Loss of	\$0.34	E. Gain of	\$0.20				
5.				_	re accepted, what is tall be accepted?	the		
	A. 25%	B. 30%	C. 33%	D. 38%	E. 44%			
6.	and the nur average nu	nber of days the	ey worked duri orked per perso	ng one month is on is 20. If 12 o	are chosen at random determined. The of the people worked cople work?			
	A. 20	B. 21	C. 22	D. 23	E. 24			

7.	In a sample of 12 people in an exercise class, their ages are given as follows:										
	26	37 20 2	2 29 31 30	29 24 3	3 35 28						
	What is the inter-quartile range of the ages?										
	A. 0	B. 2.5	C. 5	D. 7	E. 8.5						
8.	-	probability of			are face cards. What is the vithout replacement and	e					
	A. 0.1%	B. 0.5%	C. 1%	D. 2%	E. 6%						
9.	children wit	h probabilitie werage numb	es of 20%, 20%, 3	30%, 20%, 89	ve 0, 1, 2, 3, 4, or 5%, and 2% respectively, enth, couples getting						
	A. 1.0	B. 1.8	C. 2.0	D. 2.2	E. 2.8						
10	only 3 calls		5 calls a day. W	hat is the pro	bability that they will hav	e					
	A. 20%	B. 14%	C. 12%	D. 86%	F. 26.5%						
11		nced coin is f than 51 head		. What is the	approximate probability of	of					
	A. 62%	B. 46%	C. 54%	D. 38%	E. 50%						
12	12. A tennis player is successful in getting an "ace" on 12% of his serves. If he serves 50 times in his next match, how many aces can he expect to serve?										
	A. 12	B. 2	C. 20	D. 14	E. 6						
13	13. A few weeks before city council elections, a sociologist must conduct a poll to predict potential winners. With a limited budget, she must conduct the poll in a single location and on one day. In which of the following locations is he most likely to obtain the fairest sample?										
	A. A superr	narket	B. A church	n C.	A golf club						
	D. A movie	theater	E. A beauty	salon salon							

	that all 4 of	these people ar	e smokers?		
	A. 0.312	B. 0.133	C. 0.040	D. 0.032	E. 0.020
15.		_	-		board the plane was 7 nany children were ab
	A. 15	B. 7	C. 10	D. 12	E. 8
16.		's Theorem, wh			ion of 4. Using data whose value lies
	A. $\frac{1}{3}$	B. $\frac{1}{2}$	C. $\frac{5}{9}$	D. $\frac{7}{10}$	E. $\frac{4}{5}$
17.		ient of variation value of the me	-	s 0.40. If the st	andard deviation is 5,
	A. 12.5	B. 10	C. 7.5	D. 4.5	E. 2
18.	-	r population of what <i>must</i> be t	_	•	the mean is 10 and the dian?
	A. It must	be less than 25.	В. І	t must be less t	than 10.
	C. It must	be greater than	10. D. I	t must lie betw	veen 10 and 25.
	A coach for				in overtime. What is on sample of 50 hockers
19.		_		time ii a rando	
19.	expected nu	_	C. 5	D. 6	E. 7
	expected nu games are p A. 3 A six-sided	blayed? B. 4	C. 5	D. 6	

B. Free Response.

1.	The probability that merchandise stolen from a store will be recovered is 15%. Suppose a random sample of 8 stores, from which merchandise has been stolen, is chosen.							
	a. Find the probability that exactly 2 of these stores will recover their merchandise.							
	b. Find the probability that at most one store will recover their merchandise.							
	c. Find the expected number of stores that will recover their merchandise in this sample.							
	d. Find the standard deviation of the probability distribution.							
2.	In a certain lottery game, there are ten finalists for the \$100,000 grand prize. One grand prize winner is to be selected at random to receive this prize. If each contestant paid \$5 for their ticket, what is each player's expected value of their ticket?							

3.	At the Stager School of Nursing, 85% of incoming freshman are female. Recent records indicate that 70% of entering female students will graduate with a BSN degree, while 90% of the male students will obtain a BSN degree. If an incoming freshman nursing student is selected at random, find
	a. P(student will graduate, given the student is female).
	b. P(student will graduate and student is a female).
	c. P(student will graduate, given student is male).
	d. P(student will graduate and student is male)>
	e. P(student will graduate).
4.	In question 3 above, 10 incoming freshman are randomly selected. Find the probability that
	a. exactly 7 will be female students.
	b. at least 7 will be female students.
	c. at most 2 will be male students.

5. In Chemistry class, weights are assigned to required activities as follows:

Participation = 15%, Exam 1 = 20%, Exam 2 = 20%, Exam 3 = 20%, Labs = 25%

Each activity is graded on a 100 point scale. Mary earned 70 points on participation, 80 points on exam 1, 64 points on exam 2, 77 points on exam 3, and 96 points on labs. What is Mary's overall average in Chemistry?

- 6. Identify each of the following samples by naming the sampling technique used (cluster, convenience, simple random, stratified, systematic).
 - a. Measure the length of time every fifth person coming into a bank waits for teller service over a period of two days.
 - b. Take a sample of 5 Zip codes from the Chicago metropolitan region and use all the elementary schools from each of the Zip code regions. Determine the number of students enrolled in first grade in each of the schools selected.
 - c. Divide the users of a computer online service into different age groups and then select a random sample from each age group to survey the amount of time they are using the internet each week.
 - d. Survey five friends about their opinion of the student cafeteria.
 - e. Pick a random sample of students enrolled at your school and determine the number of times each has used guidance services.
 - f. Check every 10th item on an assembly line for defects.

7.	Identify the statistical display (Time plot, Bar graph, Pareto chart, Ogive, or Circle graph) which is the most appropriate for each situation.a. Identify the frequency of events or categories in decreasing order of frequency of occurrence.										
	b. Are useful for both quantitative or qualitative data. With qualitative data, the frequency or percentage of occurrence can be displayed. With quantitative data, the measurement itself can be displayed.										
	c. Display how a total is dispersed into several categories.										
	d. Display how data change over time. It is best if the units of time are consistent.										
	e. Display cumulative frequencies.										
8.	3. Find the mean and 10% trimmed mean for the annual snowfalls (in inches) for a city in northern Wisconsin. Which is the most representative of the average annual snowfall? Why?										
	24	37	28	13	38	29	112	21	40	36	

- 9. The average number of customers entering a store in a 20 minute period is 6 customers.
 - a. What is the probability that, in any 20 minute period, four customers will enter the store?
 - b. What is the probability that, in any 20 minute period, at most four customers will enter the store?
 - c. What is the probability that, in a given hour, at least 15 customers will enter the store?
- 10. Let the random variable X represent the number of telephone lines in use by a technical support center of a software manufacturer at noon each day. The probability distribution of X is shown in the table below.

X 0		1	2	3	4	5
P(X)	0.35	0.20	0.15	0.15	0.10	0.05

- a. Calculate the expected value of X.
- b. Using past records, the staff at technical support randomly selected 20 days and found an average of 1.25 telephone lines were in use at noon on those days. The staff proposes to select another random sample of 1,000 days and compute the average number of telephone lines that were in use at noon on those days. How do you expect the average from this new sample to compare to that of the first sample? Justify your response.

c. The median of a random variable is defined as any value x such that $P(X \le x) \ge 0.5$ and $P(X \ge x) \ge 0.5$. For the probability distribution shown in the table above, determine the median value of X.

d. In a sentence or two, comment on the relationship between the mean and the median relative to the shape of this distribution.