



ALLERGIC REACTION

ALGORITHM FOR MANAGEMENT OF MEDICAL EMERGENCIES

Injury and Illness Protocol Legend



Children may experience a delayed allergic reaction up to 2 hours following exposure to allergen (i.e. food ingestion, bee sting, etc.)



References:

- Scott H. Sicherer, MD, Todd Mahr, MD, and The Section on Allergy and Immunology. (2010). *Clinical report—Management of food allergy in the school setting*. Retrieved April 24, 2020, from <https://pediatrics.aappublications.org/content/pediatrics/early/2010/11/29/peds.2010-2575.full.pdf>.
- American Academy of Pediatrics & National Association of School Nurses. (2014). *PedFACTs: Pediatric first aid for caregivers and teachers (2nd ed.)*. Burlington, MA : Jones & Bartlett Publishers.
- Cosby, M.F, Lyons, E., Prestidge, L. (2019). Students with acute illness and injury. In J. Selekman, R.A. Shannon, R. A., & C.F. Yonkaitis (Eds.), *School nursing: A comprehensive text* (pp. 415-455). Philadelphia, PA: F. A. Davis Company.
- National Association of School Nurses & Epinephrine Policies and Protocols Workgroup of the National Association of School Nurses. (2014). *Sample protocol for treatment of symptoms of anaphylaxis - Epinephrine autoinjector administration by school health professionals and trained personnel for school age children – Kindergarten - Grade 12*. Retrieved April 15, 2020, from https://higherlogicdownload.s3.amazonaws.com/NASN/3870c72d-fff9-4ed7-833f-215de278d256/UploadedImages/PDFs/Epinephrine%20Policies%20Protocols%20and%20Reporting/Sample_Anaphylaxis_Epinephrine_Administration_Protocol.pdf.
- Schoessler, S., & White, M.V. (2013). Recognition and treatment of anaphylaxis in the school setting: The essential role of the school nurse. *The Journal of School Nursing*, 29(6) 407-415.