

Derivatives**Definition and Notation**

If $y = f(x)$ then the derivative is defined to be $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$.

If $y = f(x)$ then all of the following are equivalent notations for the derivative.

$$f'(x) = y' = \frac{dy}{dx} = \frac{d}{dx} f(x) = Df(x)$$

If $y = f(x)$ all of the following are equivalent notations for derivative evaluated at $x = a$.

$$f'(a) = y'|_{x=a} = \frac{dy}{dx} \bigg|_{x=a} = Df(a)$$

Interpretation of the Derivative

1. If $y = f(x)$ then,
 - a. $m = f'(a)$ is the slope of the tangent line to $y = f(x)$ at $x = a$ and the equation of the tangent line at $x = a$ is given by $y = f(a) + f'(a)(x - a)$.
 - b. $f'(a)$ is the instantaneous rate of change of $f(x)$ at $x = a$.
2. If $f(x)$ is the position of an object at time x then $f'(x)$ is the velocity of the object at $x = a$.

Basic Properties and Formulas

If $f(x)$ and $g(x)$ are differentiable functions (the derivative exists), c and n are any real numbers,

1. $(cf)' = cf'(x)$
2. $(f \pm g)' = f' \pm g'$
3. $(fg)' = f'g + fg'$ - Product Rule
4. $\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$ - Quotient Rule
5. $\frac{d}{dx}(c) = 0$
6. $\frac{d}{dx}(x^n) = nx^{n-1}$ - Power Rule
7. $\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$
This is the Chain Rule

Common Derivatives

$$\begin{aligned} \frac{d}{dx}(\sin x) &= \cos x & \frac{d}{dx}(\sec x) &= \sec x \tan x \\ \frac{d}{dx}(\cos x) &= -\sin x & \frac{d}{dx}(\csc x) &= -\csc x \cot x \\ \frac{d}{dx}(\tan x) &= \sec^2 x & \frac{d}{dx}(\cot x) &= -\csc^2 x \\ \frac{d}{dx}(\sec x) &= \sec x \tan x & \frac{d}{dx}(\sin^{-1} x) &= \frac{1}{\sqrt{1-x^2}} \\ \frac{d}{dx}(\csc x) &= -\csc x \cot x & \frac{d}{dx}(\cos^{-1} x) &= -\frac{1}{\sqrt{1-x^2}} \\ \frac{d}{dx}(\tan^{-1} x) &= \frac{1}{1+x^2} & \frac{d}{dx}(\ln x) &= \frac{1}{x}, x > 0 \\ & & \frac{d}{dx}(\log_e x) &= \frac{1}{x \ln a}, x > 0 \end{aligned}$$

Chain Rule Variants

The chain rule applied to some specific functions.

1. $\frac{d}{dx}[(f(x))^n] = n[f(x)]^{n-1} f'(x)$
2. $\frac{d}{dx}(e^{f(x)}) = f'(x)e^{f(x)}$
3. $\frac{d}{dx}(\ln[f(x)]) = \frac{f'(x)}{f(x)}$
4. $\frac{d}{dx}(\sin[f(x)]) = f'(x)\cos[f(x)]$
5. $\frac{d}{dx}(\cos[f(x)]) = -f'(x)\sin[f(x)]$
6. $\frac{d}{dx}(\tan[f(x)]) = f'(x)\sec^2[f(x)]$
7. $\frac{d}{dx}(\sec[f(x)]) = f'(x)\sec[f(x)]\tan[f(x)]$
8. $\frac{d}{dx}(\tan^{-1}[f(x)]) = \frac{f'(x)}{1+[f(x)]^2}$

Higher Order Derivatives

The Second Derivative is denoted as

$$f''(x) = f^{(2)}(x) = \frac{d^2 f}{dx^2} \text{ and is defined as}$$

$f''(x) = (f'(x))'$, i.e. the derivative of the first derivative, $f'(x)$.

The n^{th} Derivative is denoted as

$$f^{(n)}(x) = \frac{d^n f}{dx^n} \text{ and is defined as}$$

$f^{(n)}(x) = (f^{(n-1)}(x))'$, i.e. the derivative of the $(n-1)^{\text{st}}$ derivative, $f^{(n-1)}(x)$.

Implicit Differentiation

Find y' if $e^{3x+9y} + x^3 y^2 = \sin(y) + 11x$. Remember $y = y(x)$ here, so products/quotients of x and y will use the product/quotient rule and derivatives of y will use the chain rule. The "trick" is to differentiate as normal and every time you differentiate a y you tack on a y' (from the chain rule). After differentiating solve for y' .

$$\begin{aligned} e^{3x+9y}(2-9y') + 3x^2 y^2 + 2x^3 y y' &= \cos(y) y' + 11 \\ 2e^{3x+9y} - 9y' e^{3x+9y} + 3x^2 y^2 + 2x^3 y y' &= \cos(y) y' + 11 \\ (2x^3 y - 9e^{3x+9y} - \cos(y)) y' &= 11 - 2e^{3x+9y} - 3x^2 y^2 \\ y' &= \frac{11 - 2e^{3x+9y} - 3x^2 y^2}{2x^3 y - 9e^{3x+9y} - \cos(y)} \end{aligned}$$

Critical Points

$x = c$ is a critical point of $f(x)$ provided either

1. $f'(c) = 0$ or
2. $f'(c)$ doesn't exist.

Increasing/Decreasing - Concave Up/Concave Down

Concave Up/Concave Down

1. If $f''(x) > 0$ for all x in an interval I then $f(x)$ is concave up on the interval I .
2. If $f''(x) < 0$ for all x in an interval I then $f(x)$ is concave down on the interval I .

Increasing/Decreasing

1. If $f'(x) > 0$ for all x in an interval I then $f(x)$ is increasing on the interval I .
2. If $f'(x) < 0$ for all x in an interval I then $f(x)$ is decreasing on the interval I .
3. If $f'(x) = 0$ for all x in an interval I then $f(x)$ is constant on the interval I .

Inflection Points

$x = c$ is an inflection point of $f(x)$ if the concavity changes at $x = c$.

Extrema

Absolute Extrema

1. $x=c$ is an absolute maximum of $f(x)$ if $f(c) \geq f(x)$ for all x in the domain.
2. $x=c$ is an absolute minimum of $f(x)$ if $f(c) \leq f(x)$ for all x in the domain.

Fermat's Theorem

If $f(x)$ has a relative (or local) extrema at $x=c$, then $x=c$ is a critical point of $f(x)$.

Extreme Value Theorem

If $f(x)$ is continuous on the closed interval $[a, b]$ then there exist numbers c and d so that,
 1. $a \leq c, d \leq b$, 2. $f(c)$ is the abs. max. in $[a, b]$, 3. $f(d)$ is the abs. min. in $[a, b]$.

Finding Absolute Extrema

To find the absolute extrema of the continuous function $f(x)$ on the interval $[a, b]$ use the following process.

1. Find all critical points of $f(x)$ in $[a, b]$.
2. Evaluate $f(x)$ at all points found in Step 1.
3. Evaluate $f(a)$ and $f(b)$.
4. Identify the abs. max. (largest function value) and the abs. min. (smallest function value) from the evaluations in Steps 2 & 3.

Mean Value Theorem

If $f(x)$ is continuous on the closed interval $[a, b]$ and differentiable on the open interval (a, b) then there is a number $a < c < b$ such that $f'(c) = \frac{f(b) - f(a)}{b - a}$.

Rolle's Theorem


Assume that $f(x)$ is continuous on $[a, b]$ and differentiable on (a, b) .

If $f(a) = f(b)$, then there exists a number c between a and b such that $f'(c) = 0$.

Related Rates

Sketch picture and identify known/unknown quantities. Write down equation relating quantities and differentiate with respect to t using implicit differentiation (i.e. add on a derivative every time you differentiate a function of t). Plug in known quantities and solve for the unknown quantity.

Ex. A 15 foot ladder is resting against a wall. The bottom is initially 10 ft away and is being pushed towards the wall at $\frac{1}{4}$ ft/sec. How fast is the top moving after 12 sec?



x' is negative because x is decreasing. Using Pythagorean Theorem and differentiating.

$$x^2 + y^2 = 15^2 \Rightarrow 2xx' + 2yy' = 0$$

After 12 sec we have $x = 10 - 12(\frac{1}{4}) = 7$ and so $y = \sqrt{15^2 - 7^2} = \sqrt{176}$. Plug in and solve for y' .

$$7(-\frac{1}{4}) + \sqrt{176}y' = 0 \Rightarrow y' = \frac{7}{4\sqrt{176}} \text{ ft/sec}$$

Ex. Two people are 50 ft apart when one starts walking north. The angle θ changes at 0.01 rad/min. At what rate is the distance between them changing when $\theta = 0.5$ rad?



We have $\theta = 0.01$ rad/min. and want to find x' . We can use various trig fns but easiest is,

$$\sec \theta = \frac{y}{50} \Rightarrow \sec \theta \tan \theta \theta' = \frac{y'}{50}$$

We know $\theta = 0.05$ so plug in θ' and solve.

$$\sec(0.5) \tan(0.5)(0.01) = \frac{y'}{50}$$

Remember to have calculator in radians!
 $x' = 0.3112$ ft/sec

Optimization

Sketch picture if needed, write down equation to be optimized and constraint. Solve constraint for one of the two variables and plug into first equation. Find critical points of equation in range of variables and verify that they are min/max as needed.

Ex. We're enclosing a rectangular field with 500 ft of fence material and one side of the field is a building. Determine dimensions that will maximize the enclosed area.



Maximize $A = xy$ subject to constraint of $x + 2y = 500$. Solve constraint for x and plug into area.

$$x = 500 - 2y \Rightarrow A = y(500 - 2y)$$

Differentiate and find critical point(s).
 $A' = 500 - 4y \Rightarrow y = 125$

By 2nd deriv. test this is a rel. max. and so is the answer we're after. Finally, find x .

$$x = 500 - 2(125) = 250$$

The dimensions are then 250 x 125.

Ex. Determine point(s) on $y = x^2 + 1$ that are closest to (0,2).



Minimize $f = d^2 = (x-0)^2 + (y-2)^2$ and the constraint is $y = x^2 + 1$. Solve constraint for x^2 and plug into the function.

$$x^2 = y - 1 \Rightarrow f = x^2 + (y-2)^2$$

$$= y - 1 + (y-2)^2 = y^2 - 3y + 3$$

Differentiate and find critical point(s).

$$f' = 2y - 3 \Rightarrow y = \frac{3}{2}$$

By the 2nd derivative test this is a rel. min. and so all we need to do is find x value(s).

$$x^2 = \frac{3}{2} - 1 = \frac{1}{2} \Rightarrow x = \pm \frac{1}{\sqrt{2}}$$

The 2 points are then $(\frac{1}{\sqrt{2}}, \frac{3}{2})$ and $(-\frac{1}{\sqrt{2}}, \frac{3}{2})$

Adapted from:

Visit <http://tutorial.math.lamar.edu> for a complete set of Calculus notes.

Limits

Definitions

Precise Definition: We say $\lim_{x \rightarrow a} f(x) = L$ if for every $\varepsilon > 0$ there is a $\delta > 0$ such that whenever $0 < |x - a| < \delta$ then $|f(x) - L| < \varepsilon$.

“Working” Definition: We say $\lim_{x \rightarrow a} f(x) = L$ if we can make $f(x)$ as close to L as we want by taking x sufficiently close to a (on either side of a) without letting $x = a$.

Right hand limit: $\lim_{x \rightarrow a^+} f(x) = L$. This has the same definition as the limit except it requires $x > a$.

Left hand limit: $\lim_{x \rightarrow a^-} f(x) = L$. This has the same definition as the limit except it requires $x < a$.

Relationship between the limit and one-sided limits

$\lim_{x \rightarrow a} f(x) = L \Rightarrow \lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^-} f(x) = L$
 $\lim_{x \rightarrow a^+} f(x) \neq \lim_{x \rightarrow a^-} f(x) \Rightarrow \lim_{x \rightarrow a} f(x)$ Does Not Exist

Properties

Assume $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ both exist and c is any number then,

- $\lim_{x \rightarrow a} [cf(x)] = c \lim_{x \rightarrow a} f(x)$
- $\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} \left[\frac{f(x)}{g(x)} \right] = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ provided $\lim_{x \rightarrow a} g(x) \neq 0$
- $\lim_{x \rightarrow a} [f(x)]^n = \left[\lim_{x \rightarrow a} f(x) \right]^n$
- $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$

Basic Limit Evaluations at $\pm \infty$

Note: $\text{sgn}(a) = 1$ if $a > 0$ and $\text{sgn}(a) = -1$ if $a < 0$.

- $\lim_{x \rightarrow \infty} e^x = \infty$ & $\lim_{x \rightarrow -\infty} e^x = 0$
- $\lim_{x \rightarrow \infty} \ln(x) = \infty$ & $\lim_{x \rightarrow 0^+} \ln(x) = -\infty$
- If $r > 0$ then $\lim_{x \rightarrow \infty} \frac{b}{x^r} = 0$
- If $r > 0$ and x' is real for negative x then $\lim_{x \rightarrow -\infty} \frac{b}{x^r} = 0$
- n even: $\lim_{x \rightarrow \pm \infty} x^n = \infty$ & $\lim_{x \rightarrow \pm \infty} x^n = -\infty$
- n odd: $\lim_{x \rightarrow \infty} x^n = \infty$ & $\lim_{x \rightarrow -\infty} x^n = -\infty$
- n even: $\lim_{x \rightarrow \infty} a x^n + \dots + b x + c = \text{sgn}(a) \infty$
- n odd: $\lim_{x \rightarrow \infty} a x^n + \dots + b x + c = \text{sgn}(a) \infty$
- n odd: $\lim_{x \rightarrow -\infty} a x^n + \dots + b x + c = -\text{sgn}(a) \infty$

Visit <http://tutorial.math.lamar.edu> for a complete set of Calculus notes.

© 2005 Paul Dawkins

Evaluation Techniques

Continuous Functions

If $f(x)$ is continuous at a then $\lim_{x \rightarrow a} f(x) = f(a)$

Continuous Functions and Composition

$f(x)$ is continuous at b and $\lim_{x \rightarrow a} g(x) = b$ then

$$\lim_{x \rightarrow a} f(g(x)) = f\left(\lim_{x \rightarrow a} g(x)\right) = f(b)$$

Factor and Cancel

$$\begin{aligned} \lim_{x \rightarrow 2} \frac{x^2 + 4x - 12}{x^2 - 2x} &= \lim_{x \rightarrow 2} \frac{(x-2)(x+6)}{x(x-2)} \\ &= \lim_{x \rightarrow 2} \frac{x+6}{x} = \frac{8}{2} = 4 \end{aligned}$$

Rationalize Numerator/Denominator

$$\begin{aligned} \lim_{x \rightarrow 9} \frac{3 - \sqrt{x}}{x^2 - 81} &= \lim_{x \rightarrow 9} \frac{3 - \sqrt{x}}{(x-9)(x+9)} \cdot \frac{3 + \sqrt{x}}{3 + \sqrt{x}} \\ &= \lim_{x \rightarrow 9} \frac{9 - x}{(x^2 - 81)(3 + \sqrt{x})} = \lim_{x \rightarrow 9} \frac{-1}{(x-9)(x+9)(3 + \sqrt{x})} \\ &= \frac{-1}{(18)(6)} = -\frac{1}{108} \end{aligned}$$

Combine Rational Expressions

$$\begin{aligned} \lim_{h \rightarrow 0} \left(\frac{1}{x+h} - \frac{1}{x} \right) &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{x - (x+h)}{x(x+h)} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\frac{-h}{x(x+h)} \right) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)} = -\frac{1}{x^2} \end{aligned}$$

Some Continuous Functions

Partial list of continuous functions and the values of x for which they are continuous.

- Polynomials for all x .
- Rational function, except for x 's that give division by zero.
- $\sqrt[n]{x}$ (n odd) for all x .
- $\sqrt[n]{x}$ (n even) for all $x \geq 0$.
- e^x for all x .
- $\ln x$ for $x > 0$.
- $\cos(x)$ and $\sin(x)$ for all x .
- $\tan(x)$ and $\sec(x)$ provided $x \neq \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$
- $\cot(x)$ and $\csc(x)$ provided $x \neq \dots, -2\pi, -\pi, 0, \pi, 2\pi, \dots$

Intermediate Value Theorem

Suppose that $f(x)$ is continuous on $[a, b]$ and let M be any number between $f(a)$ and $f(b)$. Then there exists a number c such that $a < c < b$ and $f(c) = M$.

Polynomials at Infinity

$p(x)$ and $q(x)$ are polynomials. To compute

$$\lim_{x \rightarrow \infty} \frac{p(x)}{q(x)}$$

factor largest power of x out of both $p(x)$ and $q(x)$ and then compute limit.

$$\lim_{x \rightarrow \infty} \frac{3x^2 - 4}{5x - 2x^2} = \lim_{x \rightarrow \infty} \frac{x^2(3 - \frac{4}{x^2})}{x^2(\frac{5}{x} - 2)} = \lim_{x \rightarrow \infty} \frac{3 - \frac{4}{x^2}}{\frac{5}{x} - 2} = -\frac{3}{2}$$

Piecewise Function

$$\lim_{x \rightarrow 1} g(x) \text{ where } g(x) = \begin{cases} x^2 + 5 & \text{if } x < -2 \\ 1 - 3x & \text{if } x \geq -2 \end{cases}$$

Compute two one sided limits,

$$\lim_{x \rightarrow 1^-} g(x) = \lim_{x \rightarrow 1^-} x^2 + 5 = 9$$

$$\lim_{x \rightarrow 1^+} g(x) = \lim_{x \rightarrow 1^+} 1 - 3x = 7$$

One sided limits are different so $\lim_{x \rightarrow 1} g(x)$

doesn't exist. If the two one sided limits had been equal then $\lim_{x \rightarrow 1} g(x)$ would have existed and had the same value.

Visit <http://tutorial.math.lamar.edu> for a complete set of Calculus notes.

© 2005 Paul Dawkins

Integrals

Definitions

Definite Integral: Suppose $f(x)$ is continuous on $[a, b]$. Divide $[a, b]$ into n subintervals of width Δx and choose x_i^* from each interval. Then $\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$.

Anti-Derivative: An anti-derivative of $f(x)$ is a function, $F(x)$, such that $F'(x) = f(x)$.

Indefinite Integral: $\int f(x) dx = F(x) + C$ where $F(x)$ is an anti-derivative of $f(x)$.

Fundamental Theorem of Calculus

Part I: If $f(x)$ is continuous on $[a, b]$ then

$$g(x) = \int_a^x f(t) dt \text{ is also continuous on } [a, b]$$

$$\text{and } g'(x) = \frac{d}{dx} \int_a^x f(t) dt = f(x).$$

Part II: $f(x)$ is continuous on $[a, b]$, $F(x)$ is an anti-derivative of $f(x)$ (i.e. $F'(x) = f(x)$) then $\int_a^b f(x) dx = F(b) - F(a)$.

Properties

$$\int f(x) \pm g(x) dx = \int f(x) dx \pm \int g(x) dx$$

$$\int_a^b f(x) \pm g(x) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$$

$$\int_a^a f(x) dx = 0$$

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$$

If $f(x) \geq g(x)$ on $a \leq x \leq b$ then $\int_a^b f(x) dx \geq \int_a^b g(x) dx$

If $f(x) \geq 0$ on $a \leq x \leq b$ then $\int_a^b f(x) dx \geq 0$

If $m \leq f(x) \leq M$ on $a \leq x \leq b$ then $m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$

Common Integrals

$$\int k dx = kx + C$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C, n \neq -1$$

$$\int x^{-1} dx = \ln|x| + C$$

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln|ax+b| + C$$

$$\int \ln u du = u \ln(u) - u + C$$

$$\int e^u du = e^u + C$$

$$\int \tan u du = \ln|\sec u| + C$$

$$\int \sec u du = \ln|\sec u + \tan u| + C$$

$$\int \sec^2 u du = \tan u + C$$

$$\int \sec u \tan u du = \sec u + C$$

$$\int \csc u \cot u du = -\csc u + C$$

$$\int \csc^2 u du = -\cot u + C$$

Standard Integration Techniques

Note that at many schools all but the Substitution Rule tend to be taught in a Calculus II class.

Substitution: The substitution $u = g(x)$ will convert $\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du$ using $du = g'(x) dx$. For indefinite integrals drop the limits of integration.

$$\text{Ex. } \int_1^2 5x^2 \cos(x^3) dx = \int_1^2 5x^3 \cos(x^3) dx = \int_1^2 \frac{5}{3} \cos(u) du$$

$$u = x^3 \Rightarrow du = 3x^2 dx \Rightarrow x^2 dx = \frac{1}{3} du$$

$$x=1 \Rightarrow u=1^3=1 \quad x=2 \Rightarrow u=2^3=8$$

$$= \frac{5}{3} \sin(u) \Big|_1^8 = \frac{5}{3} (\sin(8) - \sin(1))$$

Improper Integral

An improper integral is an integral with one or more infinite limits and/or discontinuous integrands. Integral is called convergent if the limit exists and has a finite value and divergent if the limit doesn't exist or has infinite value. This is typically a Calc II topic.

Infinite Limit

$$1. \int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$$

$$2. \int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$$

$$3. \int_{-\infty}^a f(x) dx = \lim_{t \rightarrow -\infty} \int_t^a f(x) dx + \lim_{t \rightarrow \infty} \int_t^a f(x) dx \text{ provided BOTH integrals are convergent.}$$

Approximating Definite Integrals

For given integral $\int_a^b f(x) dx$ and n (must be even for Simpson's Rule) define $\Delta x = \frac{b-a}{n}$ and divide $[a, b]$ into n subintervals $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$ with $x_0 = a$ and $x_n = b$ then,

Midpoint Rule: $\int_a^b f(x) dx \approx \Delta x [f(x_1^*) + f(x_2^*) + \dots + f(x_n^*)]$, x_i^* is midpoint $[x_{i-1}, x_i]$

Trapezoid Rule: $\int_a^b f(x) dx \approx \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + \dots + 2f(x_{n-1}) + f(x_n)]$

Average Function Value: The average value of $f(x)$ on $a \leq x \leq b$ is $f_{avg} = \frac{1}{b-a} \int_a^b f(x) dx$

Adapted from:

Visit <http://tutorial.math.lamar.edu> for a complete set of Calculus notes.

Applications of Integrals


Net Area : $\int_a^b f(x)dx$ represents the net area between $f(x)$ and the x-axis with area above x-axis positive and area below x-axis negative.



Area Between Curves : The general formulas for the two main cases for each arc,

$$y = f(x) \Rightarrow A = \int_a^b [\text{upper function}] - [\text{lower function}] dx \text{ \& } x = f(y) \Rightarrow A = \int_c^d [\text{right function}] - [\text{left function}] dy$$

If the curves intersect then the area of each portion must be found individually. Here are some sketches of a couple possible situations and formulas for a couple of possible cases.



Volumes of Revolution : The two main formulas are $V = \int A(x)dx$ and $V = \int A(y)dy$. Here is some general information about each method of computing and some examples.

Rings		Cylinders	
$A = \pi ((\text{outer radius})^2 - (\text{inner radius})^2)$		$A = 2\pi (\text{radius}) (\text{width / height})$	
Limits : x/y of right/left ring Horiz. Axis use $f(x)$, Vert. Axis use $f(y)$, $g(x)$, $A(x)$ and dx , $g(y)$, $A(y)$ and dy .		Limits : x/y of inner cyl. to x/y of outer cyl. Horiz. Axis use $f(y)$, Vert. Axis use $f(x)$, $g(y)$, $A(y)$ and dy , $g(x)$, $A(x)$ and dx .	
Ex. Axis : $y = a > 0$ 	Ex. Axis : $y = a \leq 0$ 	Ex. Axis : $y = a > 0$ 	Ex. Axis : $y = a \leq 0$
outer radius : $a - f(x)$ inner radius : $a - g(x)$	outer radius : $ a + g(x)$ inner radius : $ a + f(x)$	radius : $a - y$ width : $f(y) - g(y)$	radius : $ a + y$ width : $f(y) - g(y)$

These are only a few cases for horizontal axis of rotation. If axis of rotation is the x-axis use the $y = a \leq 0$ case with $a = 0$. For vertical axis of rotation ($x = a > 0$ and $x = a \leq 0$) interchange x and y to get appropriate formulas.

Things you MUST memorize:

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\frac{d}{dx}(fg) = fg' + gf'$$

$$\frac{d}{dx}\left(\frac{f}{g}\right) = \frac{gf' - fg'}{g^2}$$

$$\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{1}{\cot \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta} = \frac{1}{\tan \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\int a \, dx = ax + C$$

$$\int x^n \, dx = \frac{x^{n+1}}{n+1} + C, \, n \neq -1$$

$$\int \frac{1}{x} \, dx = \ln|x| + C$$

$$\int e^x \, dx = e^x + C$$

$$\int a^x \, dx = \frac{a^x}{\ln a} + C$$

$$\int \sin x \, dx = -\cos x + C$$

$$\int \cos x \, dx = \sin x + C$$

$$\int \sec^2 x \, dx = \tan x + C$$

More Things you MUST memorize:

Average value of a function

Average rate vs. Instantaneous rate

Fundamental Theorem of Calculus, parts 1 and 2

Mean Value Theorem

Absolute Value Theorem

Extreme Value Theorem

Things you SHOULD memorize:

$$\frac{d}{dx}(\cot x) = -\csc^2 x$$

$$\frac{d}{dx}(\sec x) = \sec x \tan x$$

$$\frac{d}{dx}(\csc x) = -\csc x \cot x$$

Things you SHOULD memorize:

Derivatives of inverse functions:

When $f(x)$ and $g(x)$ are inverses, then $g'(x) = \frac{1}{f'(g(x))}$

Derivatives of trig inverse functions:

$$\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}} \quad \frac{d}{dx}(\arccos x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}$$

Remember how to use l'Hopital's rule to find limits that are indeterminate $\frac{0}{0}$ or $\frac{\infty}{\infty}$. See examples below:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = \lim_{x \rightarrow 0} \frac{\cos(x)}{1} = 1$$

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x} = \lim_{x \rightarrow 0} \frac{\sin x}{1} = 0$$

Things that could be handy to know:

$$\sin 2\theta = 2 \sin \theta \cos \theta \qquad \sin^2 \theta + \cos^2 \theta = 1$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta \qquad 1 + \tan^2 \theta = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$