
Nuclear Chemistry

Nar	me:Period:Date:					
Pa	Part I: ELECTROMAGNETIC RADIATION					
1)	= a general term for any type of energy that emanates or radiates outward in all directions					
2)	(ER) = radiation moving at the speed of light, ranging from high-energy gamma rays to low energy radio waves; includes visible light					
3)	= all the forms of electromagnetic radiation					
4)	Forms of ER included in the electromagnetic spectrum arranged from HIGHEST energy to LOWEST energy:					


- a) _____ b) X-rays
- c) ultraviolet light
- d) ______e) infrared light
- f) microwaves
- g) _____
- 5) Properties that all ER has in common:
 - a) form of energy with no rest massb) travels at the ______
 - c) can travel through a vacuum
 - d) emitted by atoms after they are energized or as they decay
 - e) act like both a ______and a _____(photon; packet or bundle of energy)
- factorial radiation with sufficient energy to ionize atoms or molecules (______ energy ER)
 - detaches electrons from atoms or molecules, ionizing them
 - Ex: Alpha and beta particles, gamma rays
 - Damages _____ molecules in cells!!



7)		= radiation with insufficient energ	_ = radiation with insufficient energy to		
	ionize atoms or molecules ((energy ER)			

Part II: NUCLEAR RADIATION

- 1) Background Information:
 - a) Henri Becquerel "accidentally" discovered that uranium emitted ______
 - b) _____ studied radioactivity and completed much of the pioneer work on nuclear changes
 - _____won 2 Nobel Prizes
 - the first, shared with her husband Pierre Curie and Becquerel for discovering radioactivity
 - o the second for discovering the radioactive elements radium and polonium
- 2) _____ = particles or ER emitted from a nucleus during radioactive decay
- 3) ______ = the spontaneous break-down of a nucleus into a slightly lighter nucleus, accompanied by the emission of nuclear radiation (alpha, beta, gamma)
- 4) 3 types of nuclear radiation.
 - 1. _____ (α)= helium nucleus (4_2 He)
 - 2 protons and 2 neutrons bound together emitted from a radioactive nucleus
 - Ex: $^{232}_{90}$ Th $\rightarrow ^{228}_{88}$ Ra + $^{4}_{2}$ He
 - Ex:




- 2. _____(β)= electron ($_{-1}^{0}e$)
 - A neutron is converted into a ______and an _____ (beta particle) which is ejected at a high speed

$${}^{1}_{o}n \rightarrow {}^{1}_{1}p + {}^{0}_{-1}e$$

- When an atom emits a β particle:
 - o _____ will increase by 1 (because the_____ transmutated into an additional _____).
 - Ex: decay of the isotope of carbon-14 into nitrogen-14

$$^{14}_{6}C \rightarrow ^{14}_{7}N + ^{0}_{-1}e$$

o Ex:



	3	$(\gamma) = \text{form of high-energy ER } (0)$	
	•	high energy ER waves emitted from a nucleus as it changes to a ground energy state	from an excited state
	•	gamma does itself cause the transmutation of at	oms
		o gamma radiation is often emitted during and simultaneous to, alpha or beta	
		radioactive decay.	Penetrating Power of Nuclear
	<u>Write</u>	the nuclear symbol.	
Alpl	ha =		α ♦ ———————————————————————————————————

Beta =

Gamma =

COMPARING NUCLEAR RADIATION:

<u>Property</u>	Alpha Particles	Beta Particles	Gamma Rays
Nature	Particle consisting of protons and neutrons (Helium Nucleus)	Negatively charged ————	electromagnetic radiation
Charge			Charge
Speed	Aboutthe speed of light	the speed of light	Speed of light
Mass	atomic mass units	0.0005 atomic mass units	mass at rest
Penetrating Power	Relatively (can be stopped by a piece of paper). Causes harm through ingestion or inhalation!	100 times greater than an alpha particle (can be stopped by a sheet of aluminum foil)	(can be stopped by several centimeters of lead)
Ionizing Ability			

Paper Aluminum

PART III: THE ATOMIC MODEL (A QUICK REVIEW)

1)	= smallest particle of an element that retains the chemical properties of that element (atoms are the building blocks of matter)			
2)	An atom is composed of a densely packed central region () containing protons and neutrons with electrons outside the nucleus.			
3)	electron charge =; proton charge =; neutron charge =			
4)	= atoms of the same element having different numbers of NEUTRONS and, therefore, a different mass number and atomic mass			
5)	= the number of protons in the nucleus of an atom of an element			
6)	= the total number of protons plus neutrons in the nucleus of an isotope			
7)	= element = element 8-Helium symbols are traditionally preceded by their (lower left)			
٥١	■ Ex:			
	= mass number is written after the name the element or element's symbol			
	■ Ex: <i>element – mass number</i>			
	 Write the nuclear symbol and the hyphen notation for an isotope of uranium with a mass number of 238. 			
	Nuclear symbol = Hyphen =			
	 Write the nuclear symbol and the hyphen notation for an isotope of carbon with a mass number of 14. 			
	Nuclear symbol = Hyphen =			

IV. BALANCING NUCLEAR EQUATIONS:



Total atomic numbers and mass numbers must be EQUAL on both sides!

1. Write a nuclear equation showing the radioactive decay of polonium-218 if the decay produces an alpha particle.

$$^{218}_{~84}Po~\rightarrow~____~+~^4_2He$$

2. What type of nuclear radiation is produced when polonium-212 decays to produce lead-208?

$$^{212}_{84}Po \rightarrow ^{208}_{82}Pb + _{---}$$

3. What will decay to produce lead-206 and an alpha particle?

$$\underline{\qquad} \rightarrow {}^{206}_{82}\text{Pb} + {}^{4}_{2}\text{He}$$

4. Write a nuclear equation showing the radioactive decay of carbon-14 if the decay produces a beta particle.

$$^{14}_{6}\text{C} \rightarrow ____ + ^{0}_{-1}\text{e}$$

5. What type of nuclear radiation is produced when potassium-43 decays to produce calcium-43?

$$^{43}_{19}\text{K} \rightarrow ___ + ^{43}_{20}\text{Ca}$$

6. What will decay to produce protactinium-234 and a beta particle?

$$--- \rightarrow {}^{234}_{91}Pa + {}^{0}_{-1}e$$


Part V: NUCLEAR REACTIONS

- 1) REMEMBER: ______involve some rearrangement of _____(outside the nucleus)
- 2) ______ involve changes in particles (protons/neutrons) in an atom's _____ and thus cause a change in the itself.
 - Remember that the number of protons in an atom defines the element, so a change in _____ results in a change in the atom.

	 = a change in the identity of a nucleus because of a change in the number of its protons Ex: ²³⁸/₉₂U → ²³⁴/₉₀Th + ⁴/₂He
	• In each type of nuclear reaction a small amount of theof the reactants is converted into! (E= mc²)
3) 4	types of nuclear reactions:
a)	= the spontaneous break-down of a nucleus into a slightly lighter nucleus, accompanied by the emission of nuclear radiation (alpha, beta, or gamma)
b)) = nucleus is bombarded with alpha particles,
	protons, neutrons or other particlesScientists can change the identity of atoms by hitting them with small particles.
	• Ex: An aluminum-27 nucleus is hit by an alpha particle:
	$^{27}_{13}\text{Al} + ^{4}_{2}\text{He} \rightarrow ^{30}_{15}\text{P} + ^{1}_{0}\text{n}$
	• Ex: A sulfur-32 nucleus is hit by a neutron:
c)	a very heavy nucleus splits to form medium-weight nuclei
d)) = the process in which light weight nuclei combine to form a heavier, more stable nucleus
Pai	rt VI: RADIOACTIVE DECAY & HALF-LIFE
1) H	ow long does it take for ONE radioactive atom to decay?
•	.
•	The time it takes for a of identical radioactive atoms to decay CAN BE determined based on
2) _	= an isotope with an unstable ucleus that undergoes radioactive decay
a)	= the spontaneous break-down of a nucleus into a slightly lighter nucleus, accompanied by the emission of nuclear radiation (alpha, beta, or gamma)
•	Most, but not all, isotopes of elements with an atomic number of 83 or lower are
•	All isotopes of elements with an atomic number over 83 are

2) The rate of decay of ______ is measured in half-lives. Sulfur-38 = time needed for decay of one-half (radioactive) the atoms in a SAMPLE of radioactive material 2.48 hours Chlorine-38 4) Half-life of polonium-212 = 3×10^{-7} seconds (radioactive) 0.62 hours a) Within 3 x 10^{-7} seconds half of the atoms in a sample of Argon-38 polonium-212 will decay, changing into atoms of a (stable) ______ element and giving off 5) Half-life of uranium-238 = 4.5 billion years a) Within 4.5 billion years half of the atoms in a sample of uranium-238 will decay, changing into atoms of a different element and giving off nuclear radiation. Part VII: FISSION _____ = the process in which a very heavy nucleus splits to form medium-weight nuclei 1) ___ ■ A chain reaction occurs → each nuclear fission produces two or more , which can in turn cause more nuclear Ex: Nuclear power plants (), atomic bombs Example of a fission reaction of uranium-235: + ENERGY 235.043924 amu The mass of Reactants: the **products** 1.008665 amu is LESS than the mass of Total mass of REACTANTS 236.052589 amu the reactants! 140.914363 amu Products: 91.926270 amu 3.025995 amu Total mass of PRODUCTS 235.866628 amu MASS LOSS: 236.052589 amu - 235.866628 amu = 0.185961 amu!

- The mass of the ______is LESS than the mass of the
 - √ Why?
 - or missing mass (about 0.2 amu) is converted to a lot of energy!
- Small amounts of _____ are converted to HUGE amounts of _____ according to Einstein's formula:
 - E = energy
 - m= mass
 - $c = \text{speed of light (3 x 10}^8 \text{ m/s)}$
- _____ is the only naturally occurring element that undergoes fission, however, many synthetic nuclei split under neutron bombardment. Ex: plutonium-239
- ______ = in fission reactions, neutrons are emitted; these neutrons can split more nuclei and a chain reaction can occur
- mass of fissionable material needed to sustain a nuclear chain reaction



Part VIII: FUSION

- 1) _____ = the process in which light weight nuclei combine to form a heavier, more stable nucleus
 - Nuclear fusion produces _____energy than fission!
 - Ex: Hydrogen bombs
 - Ex: fusion powers the _____ and other _____. The reaction that occur in the sun can be represented as follows (hydrogen isotopes are fused into a helium isotope):

$$^2_1\text{H} + ^3_1\text{H} \rightarrow ^4_2\text{He} + ^1_0\text{n} + \text{ENERGY}$$

Reactants:

²H 2.0140 amu

₊ ³H 3.01605 amu

Total mass of REACTANTS

5.03005 amu


Products:


⁴He 4.00260 amu

₊ ¹₀n 1.008665 amu

Total mass of PRODUCTS

5.011265 amu






MASS LOSS: 5.03005 amu - 5.011265 amu = 0.018785 amu!

Where it went \rightarrow E = mc²

Awesome Energy of stars and H-bombs!!



Part IX: 5 BENEFICIAL USES OF RADIATION

- 1. _____radiation- cancer, X-rays, diagnostic tracers
 - Ex: Iodine-131 is used to diagnose the activity of the thyroid gland; treatment of thyroid cancer.
 - o Cobalt-60, phosphorus-32, gallium-67 and cesium-137 are used in the radiation treatment of many cancers.
 - Sodium-24 is used to detect constrictions and obstructions in the circulatory system.
 - o If a compound that contains a beta emitter is injected into a tumor, the beta particles will destroy the tumor and since the beta particles will only travel a short distance, very little healthy tissue will be destroyed.

- The rapidly dividing cancer-producing cells are most susceptible to radiation damage.
- Focused radiation beams can shrink or eliminate cancerous tumors while doing less damage to surrounding tissue.
- 2. _____: weapons
 - WWII: The U.S. Army Air Force received orders to drop these weapons anytime after August 3, 1945. On August 6, "Little Boy" fell on Hiroshima. Three days later, "Fat Man" destroyed Nagasaki.





Little Boy - Hiroshima - 0815, August 6, 1945

Size: 10 ft long

Weight: 8.900 lbs. (132 lbs >90% U-235)

(-- 2lbs underwent fission)

Height of blast: 1900 ft.

Yield: 15 - 16 K1 TNT

Casualties ~ 100.000 immediate deaths

 \sim 200,000 total deaths

Fat Man - Nagasaki - 1102, August 9, 1945

Size: ~ 10.5 ft.long, 5ft. Diameter

Weight: 10,300 lbs.. (12 lbs.. Pu-239 of which

-- 2 lbs. underwent fission)

Height of blast: 1650 ft.

Yield: 22 Kt of TNT

Casualties: ~-70,000 immediate deaths

~140,000 total deaths


- 3. ______ "carbon dating"
 - Determines the age of something (rocks, fossils, skeletal remains, etc)
 - Carbon-14 is a beta emitter that decays to nitrogen-14 with a half-life of 5,730 years.
 - In every living thing there is a constant ration of normal C-12 and radioactive C-14. You can calculate the time needed to change to change from what is expected to what is actually found.

4.	<u></u> :	mi	crowaves,	sterilize	1000
5.		•	electricity	v (TMI)	

NOTE: the following is NOT on your TEST!! @

Part X: NUCLEAR POWER PLANTS

- The secret to controlling a chain reaction is to control the neutrons.
 - o If the <u>neutrons</u> can be controlled, then the energy can be released in a <u>controlled way</u>. That's what scientists have done with nuclear power plants.
- United States has about <u>100</u> nuclear reactors, producing a little more than <u>20 %</u> of the country's electricity.
- In France, almost 80 percent of the country's electricity is generated through nuclear fission.
- How do nuclear power plants make electricity?



1. Fuel for nuclear reactors:

- a) The fuel for nuclear reactors is held in fuel rods.
- b) Fuel rods usually contain mostly non-fissionable uranium-238 and about 3% fissionable uranium-235.
- c) A critical mass of fissionable uranium-235 is present (but not nearly enough for a nuclear explosion).

2. Energy production:

- a) A chain reaction fission reaction of uranium-235 occurs in the reactor core.
- b) During the fission of uranium-235 a small amount of mass is converted into a large amount of energy.
- c) The energy generated heats water in the reactor core.
- d) Heated water in the reactor core (under high pressure so it remains a liquid) transfers heat to a second loop of water.
- e) Heated water in the second loop changes to steam.
- f) This steam spins the turbines of giant generators that produce electrical energy (electricity).

3. Cooling the reactor:

- a) Steam that has moved past the turbines is cooled by water taken from a nearby body of water
- b) Cooled steam condenses to liquid water and re-circulates.
- c) So much heat is generated that some must be released into the air through cooling towers.

4. Facilitating the reaction:

- a) Moderator = something which slows down high-speed neutrons
- b) Water often acts as the moderator in nuclear reactors.
- c) A moderator slows down high-speed neutrons making them more readily captured by other nuclei.

5. Controlling the reaction:

- a) Controlling the neutrons in a reactor allows for control of the nuclear chain reaction.
- b) Control rods = rods made of neutron absorbing material (such as boron or cadmium)
- c) Control rods are inserted between fuel rods to control the reaction.
- d) When all the control rods are pushed all the way into the reactor core, all neutrons are absorbed and the chain reaction stops.

• Nuclear power plants have CERTAIN ADVANTAGES:

- No <u>fossil fuels</u> are burned (saving fossil-fuel resources for producing plastics and medicines)
- No combustion products like carbon dioxide and sulfur dioxide to pollute the air and water.

PROBLEMS associated with nuclear power plants:

- 1. **Cost**: expensive to build and operate.
 - The electricity that's generated by nuclear power costs about **twice** as much as electricity generated through fossil fuel or hydroelectric plants.
- 2. **Supply** of fissionable uranium-235 is limited.
 - Of all the naturally occurring uranium, only about 0.75 percent is U-235. A
 vast majority is non-fissionable U-238. At current usage levels, we'll be out
 of naturally occurring U-235 in fewer than 100 years. But there's a limit to
 the amount of nuclear fuel available in the earth, just as there's a limit to
 the amount of fossil fuels.
- 3. Accidents (safety)
- 4. **Disposal** of nuclear wastes.

Accidents: Three Mile Island and Chernobyl:

• <u>Three Mile Island</u> (1979): A combination of operator error and equipment failure caused a loss of reactor core coolant. The loss of coolant led to a partial meltdown and

the release of a small amount of radioactive gas. There was no loss of life or injury to plant personnel or the general population.

- <u>Chernobyl</u>, Ukraine (1986): Human error, along with poor reactor design and engineering, contributed to a tremendous overheating of the reactor core, causing it to rupture.
 - Two explosions and a fire resulted, blowing apart the core and scattering nuclear material into the atmosphere. A small amount of this material made its way to Europe and Asia.
 - o The area around the plant is still uninhabitable.
 - The reactor has been encased in concrete, and it must remain that way for hundreds of years.
 - o Hundreds of people died.
 - o Many others felt the effect of radiation poisoning.
 - Instances of thyroid cancer, possibly caused by the release of I-13, have risen dramatically in the towns surrounding Chernobyl.

How do you get rid of this stuff?: NUCLEAR WASTES

- The fission process produces large amounts of radioactive isotopes, and some of the half-lives of radioactive isotopes are rather long.
- Those isotopes are <u>safe</u> after <u>ten half-lives</u>.
- The wastes are basically buried and guarded at the sites. High-level wastes pose a much larger problem.
 - They're temporarily being stored at the site of generation, with plans to eventually seal the material in glass and then in drums. The material will then be stored underground in Nevada.
- At any rate, the waste must be kept safe and undisturbed for at least <u>10,000 years</u>.

Radioactive Decay & Half Life Worksheet

Atoms of the same element with different numbers of neutrons are called **isotopes**. For example, all atoms of the element carbon have 6 protons, but while most carbon atoms have 6 neutrons, some have 7 or 8. Isotopes are named by giving the name of the element followed by the sum of the neutrons and protons in the isotope's nucleus. So a carbon atom with 6 protons and 6 neutrons in its nucleus is called Carbon-12. The carbon atom with 8 neutrons is called Carbon-14. It is interesting to note that Carbon-14 is radioactive while Carbon-12 and Carbon-13 are stable.

Radioactive Decay

When the nucleus of a radioactive isotope gives up its extra energy, that energy is called **ionizing radiation**. Ionizing radiation may take the form of alpha particles, beta particles, or gamma rays. Ionizing radiation is of

concern because it may cause adverse health effects. The process of emitting the radiation is called **radioactive decay**. When the nucleus of a radioactive isotope decays, emitting ionizing radiation, the nucleus is altered. It is transformed into another isotope which in many cases is a different element. This new isotope may be stable or unstable. If it is stable, the new isotope is not radioactive. If it is unstable, it also will decay, transforming its nucleus and emitting more ionizing radiation. Several decays may be required before a stable isotope is produced. This sequence is known as a **decay chain** (see Figure 1).





Figure 1. Decay Chain for an Isotope of Sulfur

Decay chains of radioactive isotopes have been studied extensively. Scientists know exactly how many decays are required for each radioactive isotope to become stable. In addition, they know how much energy will be released with each decay.



Half-Life

It is *not* possible to predict exactly when a SINGLE radioactive atom will decay. However, the time required for half of large NUMBER OF IDENTICAL radioactive atoms to decay has been determined. This time is called the half-life. Suppose, for example, a large number of atoms of a radioactive isotope with a half-life of three hours were put in a box. After three hours, one-half of those radioactive atoms would remain. The other half would have been transformed to a different isotope. After three more hours, only half of the remaining radioactive atoms (one quarter of the initial number) would still be unchanged. The concept of half-life is illustrated in Figure 2.

The half-life can vary substantially from one isotope to another, ranging from a fraction of a second to billions of years. For example, Iodine-131, an isotope with important medical applications, has a half-life of 8.04 days.

Figure 2. Radioactive Decay of Atoms with a Half-Life of Three Hours.

Calculating Half-life:

1.	Suppose we have 100 nuclei of a radioactive isotope. After one half-life, half of the nuclei will have disintegrated, leaving 50 nuclei. a. How many nuclei will be left after the second half-life?
	b. How many nuclei would you predict would remain after the third half-life?
2.	Element X has a mass of 300g and a $\frac{1}{2}$ life of 10 years. How many grams will remain after 40 years? <u>Hint:</u> first determine how many $\frac{1}{2}$ lives is 40 years.
3.	Element Z has a mass of 430g. Scientists determined it to have a ½ life of 5 years. How many grams of element Z will remain after 15 years? Show your work.
4.	Phosphourous-32 is a radioisotope with a half-life of 14.3 days. If you start with 4 g of phosphorous-32, how many grams will remain after 57.2 days? How many half-lives will have passed?

5.	Carbon-14 is a radioisotope with a half-life of 5715 years. Suppose you start with 400 mg of carbon-14. How many half-lives will have passed when 12.5 mg of carbon-14 remains? How much time will have passed when 12.5 mg carbon-14 remains?
6.	Astatine-218 is a radioisotope with a half-life of 1.6 seconds. If you start with 10 mg of astatine-218, how many
	milligrams will remain after 5 half-lives? How much time will have passed after 5 half-lives?
7.	Uranium-238 has a half-life of 4.5 billion years (4,500,000,000 years). If you start with 10 mg of uranium-238, how many milligrams will remain after five half-lives? How much time will have passed after five half-lives?

Nuclear Radiation Worksheet



Directions: Answer the following questions and balance each nuclear equation on the next page.

1)	What is ER?
2)	What type of radiation can damage DNA?
3)	What is the name of the woman who was a pioneer of nuclear chemistry?
4)	What makes the mass of C-12 and C-14 different?
5)	Write the hyphen notation for the following: atomic number 11 and mass number 23.
6)	Write the nuclear symbol notation for the following: atomic number 11 and mass number 23.
7)	What are the 3 types of nuclear radiation?
	a)
	b)
	c)
8)	What is the nuclear symbol for an alpha particle and beta particle?
9)	What could stop gamma rays? an alpha particle?
10)	What is the charge of a beta particle?
11)	What is the charge of an alpha particle?
12)	What type of nuclear radiation has the LARGEST mass?
13)	What type of nuclear radiation has the SMALLEST mass?
14)	What type of nuclear (ionizing) radiation can cause the most damage to your DNA?
15)	The spontaneous break-down of a nucleus into a slightly lighter nucleus, accompanied by the emission of nuclear radiation is known as
16)	Explain how are nuclear reactions different from chemical reactions?

Balancing Nuclear Equations Worksheet



1)
$${}^{14}_{6}C \rightarrow {}^{14}_{7}N + _{---}$$

2)
$$\longrightarrow$$
 $^{228}_{88}$ Ra + $^{4}_{2}$ He

3)
$$^{228}_{89}\text{Ac} \rightarrow \underline{\qquad} + {}^{0}_{-1}\text{e}$$

4)
$$\longrightarrow {}^{220}_{86}\text{Rn} + {}^{4}_{2}\text{He}$$

5)
$$^{234}_{91}Pa \rightarrow ^{234}_{92}U + _{---}$$

6)
$$^{232}_{90}$$
Th $\rightarrow ^{228}_{88}$ Ra + _____

7)
$$^{60}_{27}\text{Co} \rightarrow \underline{\qquad} + {}^{0}_{-1}\text{e}$$

8)
$$\longrightarrow_{20}^{40} \text{Ca} + _{-1}^{0} \text{e}$$

9)
$$^{241}_{95}\text{Am} \rightarrow _{----} + ^{4}_{2}\text{He}$$

10)
$$^{222}_{86}$$
Ra $\rightarrow ^{218}_{84}$ Po + _____

11)
$$^{40}_{19}$$
K $\rightarrow ^{40}_{20}$ Ca + _____

12)
$$^{237}_{93}\text{Np} \rightarrow _{----} + ^{4}_{2}\text{He}$$

13)
$$\rightarrow _{-1}^{0}e + _{28}^{60}Ni$$

14)
$$^{228}_{88}$$
Ra \rightarrow _____ + $^{228}_{89}$ Ac

15)
$$^{233}_{92}\text{U} \rightarrow ^{4}_{2}\text{He} + _____$$

16)
$$^{239}_{92}\text{U} \rightarrow \underline{\qquad} + ^{239}_{93}\text{Np}$$

17) _____
$$\rightarrow {}_{2}^{4}\text{He} + {}_{82}^{208}\text{Pb}$$

18)
$$\rightarrow {}^{32}_{15}P + {}^{0}_{-1}e$$



Nuclear Chemistry Worksheet


1) How long does it take one radioactive atom to decay?

2)	is the time needed for decay of one-half the atoms in a SAMPLE of
	radioactive material is the time needed for decay of one-half the atoms in a SAMPLE of
3)	is a change in the identity of a nucleus because of a change in the number of
	its protons
4)	List the 4 types of nuclear reactions.
	b)
	a) b) c) d)
5)	What is the difference between fission and fusion?
3)	what is the difference between fission and fusion:
6)	What type of nuclear reaction takes place at TMI?
7)	What type of nuclear reaction takes place at the sun?
8)	During fission and fusion reactions, what is the relationship between the MASS of the reactants and products?
	a) WHY?
9)	is the missing mass of a nuclear reaction that is converted into
10)	energy. Small amounts of mass are converted to HUGE amounts of energy according to Einstein's famous formula. What is this famous formula?
11)	is the mass of fissionable material needed to sustain a nuclear chain reaction
12)	Which would produce the MOST energy from a reaction? Fission OR Fusion
13)	Explain 3 different areas or fields that radiation is beneficial.

Class: Academic Chemistry B - Grade 11: PA Standard 3.4.A Explain concepts about the structure and properties of matter.

Unit Essential Question(s):

How are nuclear reactions different from chemical reactions?



Concept	Concept	Concept	Concept		
Nuclear Equations	Half live				
Lesson Essential Questions:	Lesson Essential Questions:	Lesson Essential Questions:	Lesson Essential Questions		
How do you balance nuclear equations?	How do you solve half life problems?				
Vocabulary:	Vocabulary:	Vocabulary:	Vocabulary:		
Radiation Electromagnetic radiation Ionizing radiation Non-ionizing radiation Marie curie Nuclear reactions Isotope Hyphen notation Nuclear symbol notation Transmutation Radioactive isotope Radioactive decay Nuclear radiation Alpha particle Beta particle Gamma ray	Half life Nuclear fission Chain reaction Critical mass Nuclear fusion Nuclear bombardment Mass defect				

Nuclear Chemistry Vocabulary:

- 1) **Radiation** = a general term for any type of energy that emanates or radiates outward in all directions
- 2) <u>Electromagnetic radiation (ER)</u> = radiation moving at the speed of light, ranging from high-energy gamma rays to low energy radio waves; includes visible light
- 3) **Electromagnetic spectrum** = all the forms of electromagnetic radiation
- 4) <u>Ionizing radiation</u> = radiation with sufficient energy to ionize atoms or molecules (higher energy ER); damages DNA
- 5) <u>Non-ionizing radiation</u> = radiation with insufficient energy to ionize atoms or molecules (lower energy ER)
- 6) <u>Marie Curie</u> = studied radioactivity and completed much of the pioneering work on nuclear changes. Won two Nobel Prizes (the first for discovering radioactivity; the second for discovering the radioactive elements radium and polonium)
- 7) <u>Nuclear reactions</u> = involve changes in particles in an atom's nucleus and thus cause a change in the atom itself
- 8) **Isotope** = atoms of the same element having different numbers of neutrons and, therefore, a different mass number and atomic mass
- 9) **Hyphen notation** = (example) U-238
- 10) **Nuclear notation** = (example)
- 11) **Transmutation** = change in the identity of a nucleus because of a change in the number of its protons
- 12) **Radioactive isotope** = an isotope with an unstable nucleus that undergoes radioactive decay
- 13) **Radioactive decay** = the spontaneous break-down of a nucleus into a slightly lighter nucleus, accompanied by the emission of nuclear radiation
- 14) Nuclear radiation = particles or ER emitted from a nucleus during radioactive decay
- 15) <u>Alpha particle</u> (α)= helium nucleus (); 2 protons and 2 neutrons bound together emitted from a radioactive nucleus
- 16) **Beta particle** (β)= electron() ejected at a high speed when a neutron changes into a proton and an electron
- 17) Gamma ray (γ) = form of high-energy ER (); often emitted during and simultaneous to, α or β radioactive decay.
- 18) **Half-life** = time needed for decay of one-half the atoms in a *sample* of radioactive material
- 19) <u>Nuclear bombardment</u> = nucleus is bombarded with alpha particles, protons, neutrons or other particles
- 20) **Nuclear fission** =process in which a heavy nucleus splits to form medium-weight nuclei; Ex: TMI
- 21) <u>Chain reaction</u> = in fission reactions, neutrons are emitted; these neutrons can split more nuclei and a chain reaction can occur
- 22) **Critical mass** = mass of fissionable material needed to sustain a nuclear chain reaction
- 23) <u>Nuclear fusion</u> = the process in which light weight nuclei combine to form a heavier, more stable nucleus; Ex: the sun
- 24) <u>Nuclear fission</u> = the process in which a very heavy nucleus splits to form medium-weight nuclei; Ex: TMI (nuclear power plant)
- 25) **Mass defect** = the missing mass of a nuclear reaction that is converted into energy.

