



MAMMOTH SCIENCE



EST

2022

PUBLISHING



# Biomolecules

- Differentiate the structures and functions of carbohydrates, lipids, proteins, and nucleic acids.
- Analyze the relationship between monomers and polymers in biological molecules.
- Evaluate the roles of dehydration synthesis and hydrolysis in the formation of biomolecules.
- Assess the elemental composition and ratios in carbohydrates, lipids, and nucleic acids.
- Evaluate the diverse functions of carbohydrates, lipids, proteins, and nucleic acids in living organisms.
- Analyze the structure and significance of phospholipids in cellular membranes.
- Assess the relationship between amino acid sequences, protein structure, and function.
- Explain the role of nucleic acids, specifically DNA and RNA, in storing and transmitting genetic information.
- Evaluate the impact of chemical reactions on compound bonds and energy changes.
- Analyze the role of enzymes as biological catalysts, including their influence on activation energy, substrate binding, and reaction rates.



# Menu

Macromolecule Characteristics

Dehydration Synthesis & Hydrolysis

Carbohydrates

Lipids

Proteins/ Enzymes

Nucleic Acids




# Macromolecules / Biomolecules



## Macromolecules

### Characteristics -

- Large organic molecules.
- Also called **polymers**.
- Made up of smaller “building blocks” called monomers.
- **Biological Macromolecules** are:
  1. Carbohydrates
  2. Lipids
  3. Proteins
  4. Nucleic acids (DNA and RNA)



# Dehydration Synthesis vs. Hydrolysis




## Dehydration Synthesis & Hydrolysis –

- Cells connect monomers to make macromolecules by a process called condensation or **dehydration synthesis** (removing a molecule of water).
- Cells break down macromolecules into monomers by a process called **hydrolysis** (adding a molecule of water)


(a) Dehydration synthesis

Monomers are joined by removal of OH from one monomer and removal of H from the other at the site of bond formation.



(b) Hydrolysis

Monomers are released by the addition of a water molecule, adding OH to one monomer and H to the other.



# Carbohydrates



## 01

### Characteristics

- Carbohydrates can range in size from small sugar molecules to large strands of sugar molecules bonded together. A biochemical test to indicate the presence of sugar – **Benedict's solution** (indicator) – turns green, yellow, orange, or red when heated.
- Elements Used to Form:
  1. Carbon
  2. Hydrogen
  3. Oxygen

## 02

### Monomer - Monosaccharide

1. **Monosaccharide**
  - ❑ Called **Simple Sugars** (*monomer*).
  - ❑ Include: Glucose, Fructose & Galactose.
  - ❑ They have the same chemical formula,  $C_6H_{12}O_6$ , but have different structures.
  - ❑ These three sugars are structural isomers:
    - **Glucose** is found in sports drinks.
    - **Galactose** is called "milk sugar"
    - **Fructose** is found in fruits.
  - ❑ If the compound name ends in – **ose** means it is a sugar.
  - ❑ In aqueous (watery) solutions, Monosaccharides **form ring structures**.
  - ❑ They are the main fuel that cells use for work.

## 03

### Disaccharide

2. **Disaccharide**
  - ❑ A **disaccharide** is made of two sugar molecules joined together.
  - ❑ They are joined in the dehydration process.
  - ❑ There is a glycosidic bond between the two sugars.

# Carbohydrates Continued



## 3. Polysaccharide

- ❑ Complex carbohydrates
- ❑ Composed of many sugar monomers linked together.
- ❑ **Polymers** of monosaccharide chains.
- ❑ Three types of polysaccharides are: starch, glycogen, and cellulose.
  - **Starch** – in **iodine** (indicator) turns dark blue or black
    - Starch is an example of a polysaccharide in plants.
    - Plant cells store starch for energy.
    - Potatoes and grains are major sources of starch in the human diet.

## ● Glycogen

- Animals store excess sugar in the form of glycogen.
- Glycogen is similar in structure to starch because both are made of glucose monomers.

## ● Cellulose

- Cellulose is the most abundant organic compound on Earth.
- It is the structural component of plants.
- It forms cable-like fibrils in the tough walls that enclose plants.
- Many animals cannot digest cellulose.
- It is also known as dietary fiber.

## ● Chitin


- Chitin is made of glucose subunits linked together in a chain.
- Partly derived from non-sugars (nitrogen).
- Composes exoskeletons of insects, lobsters, and shrimp.
- The cell walls of fungi are also made from chitin.
- Chitin fibers are also used for dissolving sutures.

# Carbohydrates




## 01


### Monosaccharide




Glucose  
Linear Form  
Fisher Projection




Glucose  
Linear Form  
Bond Angle Drawing



Glucose  
Cyclic Form  
Bond Angle Drawing




Glucose  
Cyclic Form  
Chair Conformation



Glucose  
Cyclic Form  
Haworth Projection


## 02

### Disaccharide



## 03

### Polysaccharide



# Lipids



## 01

### Characteristics

- Lipids are **hydrophobic** –“water fearing”.
- Do not mix with water.
- Includes fats, waxes, steroids, oils & triglycerides.
- Stores the most energy.
- Composed of: C, H, & O
- **Sudan** (indicator) can turn a variety of colors in the presence of lipids.
- Most plant oils tend to be low in saturated fatty acids & exist as liquids at room temperature (oils).

## 02

### Functions

- Long term energy storage
- Protection against heat loss (insulation)
- Protection against water loss
- Protection against physical shock
- Chemical messengers (hormones)
- Major component of membranes (**phospholipids**)

## 03

### Structure - Fatty acids

- **Fatty Acids:** a long chain of Hydrogen and Carbon make up lipids. The exception to the monomer/polymer rule is lipids. Lipid base units are not considered monomers. One type of lipid or fat is made up of fatty acids and glycerol molecules in a 3:1 ratio. The bonding of three fatty acids to one glycerol molecule creates a triglyceride




# Types of Lipids



## 01

Saturated


1. **Saturated Fatty Acid:** if each Carbon is bonded by single bonds to other Carbons inside the fatty acid, the fatty acid is said to be saturated.



## 02

Unsaturated


2. **Mono or Unsaturated Fatty Acid:** If a double bond is present, then it is said to be mono-saturated or unsaturated.



## 03

Polysaturated

3. **Polyunsaturated Fatty Acid** - More than one double bond
  - Most animal fats have a high proportion of saturated fatty acids & exist as solids at room temperature (butter, margarine, shortening).




# Lipids Continued



4. **Phospholipids** - Cell membranes are made of lipids called phospholipids. Controls what enters and leaves the cell

- Phospholipids have a head that is polar & attract water (**hydrophilic**).
- Phospholipids also have 2 tails that are nonpolar and do not attract water (**hydrophobic**)
- **Steroids** –
  - ❑ The carbon skeleton of steroids is bent to form 4 fused rings.
  - ❑ Cholesterol is the “base steroid” from which your body produces other steroids.



Phospholipids

# Proteins



## 01

### Characteristics

- Composed of
  - ❑ C, H, O, N, sometimes S
- **Monomer**
  - ❑ amino acids
    - 20 different (1 mod)
    - Most structural variation
- **Polymer: Polypeptide**
  - ❑ Covalent bonding links
    - Peptide Bond

## 02

### Functions

- Essential to Life – **Biuret** (indicator)  
turns purple/lavender with protein, pink with amino acid.
- **Build structure**
- **Movement:**
  - ❑ Makes up muscle tissue
- **Transport:**
  - ❑ Carries oxygen in an organism (hemoglobin).

## 03


### Functions Cont'

- **Immunity:**
  - ❑ Helps fight off foreign invaders
  - ❑ antibodies
- **Enzymes:** (more below)
  - ❑ Speed up chemical reactions
  - ❑ amylase and pepsin
- **Energy source**
  - ❑ 1 gram = 4 kcal of energy


# Amino Acid / Polypeptide



## Monomer - Amino Acid




## Polymer - Polypeptide




## Primary Structure

- The **Primary structure** is the specific sequence of amino acids in a protein. Called a **polypeptide**.



## Secondary Structure

- **Secondary protein structures** occur when protein chains coil(helix) or fold(pleated)




## Tertiary Structure


- When protein chains called polypeptides join together, the **Tertiary structure** forms because R groups interact with each other.
- Secondary structures bent and folded into a more complex 3-D arrangement of linked polypeptides
- Bonds: H-bonds, ionic, disulfide bridges (S-S)

## Quaternary Structure

- **Quaternary**  
**Structure:**  
Composed of 2 or more “subunits”.




## Summary







Quaternary structure  
complex of protein molecules



Secondary structure  
regular sub-structures



**Tertiary structure**  
three-dimensional structure

The image displays the chemical structures of five amino acids, categorized into positive and negative groups. Each structure shows the amino acid in its zwitterion form at physiological pH, with the amino group protonated and the carboxyl group deprotonated. The structures are as follows:

- Arginine (Arg) (R):** A positive amino acid with a guanidinium side chain. The side chain is shown with a positive charge on the nitrogen atom. The pKa values are 2.01 for the carboxyl group and 10.00 for the guanidinium group.
- Histidine (His) (H):** A positive amino acid with an imidazole side chain. The side chain is shown with a positive charge on the nitrogen atom. The pKa values are 1.70 for the carboxyl group and 6.04 for the imidazole group.
- Lysine (Lys) (K):** A positive amino acid with a long aliphatic side chain ending in a primary amine. The side chain is shown with a positive charge on the terminal nitrogen atom. The pKa values are 2.15 for the carboxyl group and 10.07 for the terminal amine group.
- Aspartic Acid (Asp) (D):** A negative amino acid with a side chain ending in a carboxyl group. The side chain is shown with a negative charge on the terminal carboxylate group. The pKa values are 1.95 for the carboxyl group and 3.90 for the side chain carboxyl group.
- Glutamic Acid (Glu) (E):** A negative amino acid with a side chain ending in a carboxyl group. The side chain is shown with a negative charge on the terminal carboxylate group. The pKa values are 2.16 for the carboxyl group and 4.25 for the side chain carboxyl group.

Serine (Ser) **S** pKa 2.12, pKa 9.05

Threonine (Thr) **T** pKa 2.20, pKa 9.06

Asparagine (Asn) **N** pKa 2.16, pKa 8.76

Glutamine (Gln) **Q** pKa 2.18, pKa 9.00

Cysteine (Cys) **C** pKa 1.91, pKa 10.28

Selenocysteine (Sec) **U** pKa 1.55, pKa 10

Glycine (Gly) **G** pKa 2.34, pKa 9.38

Proline (Pro) **P** pKa 1.04, pKa 10.47

Alanine (Ala) **A** C[C@H](C(=O)O)C(=O)N pKa 2.33, pKa 9.71

Isoleucine (Ile) **I** CC[C@H](C)[C@H](C(=O)O)C(=O)N pKa 2.20, pKa 9.02

Leucine (Leu) **L** CC(C)[C@H](C(=O)O)C(=O)N pKa 2.32, pKa 9.58

Methionine (Met) **M** CSCC[C@H](C(=O)O)C(=O)N pKa 2.35, pKa 9.10

Phenylalanine (Phe) **F** c1ccc(cc1)[C@H](C(=O)O)C(=O)N pKa 2.18, pKa 9.29

Tryptophan (Trp) **W** c1ccc2c(c1)c(c[nH]2)[C@H](C(=O)O)C(=O)N pKa 2.38, pKa 9.31


Tyrosine (Tyr) **Y** c1ccc(cc1)[C@H](C(=O)O)C(=O)N pKa 2.24, pKa 9.04

Valine (Val) **V** CC(C)[C@H](C(=O)O)C(=O)N pKa 2.27, pKa 9.12


# Enzymatic Activity



- Many proteins act as biological catalysts or enzymes.
    1. Thousands of different enzymes exist in the body.
    2. *Enzymes control the rate of chemical reactions by weakening bonds, thus lowering the amount of activation energy needed for the reaction.*
    3. This is accomplished by binding to the reactants.
    4. They will then twist or bend the material, lowering the energy needed to split it.
  - The enzyme contains an opening called its activation (active) site.
    1. This site is specific for the object it will hold, called the **substrate**.
    2. There are just as many enzymes as there are substrates.
    3. The enzyme system is the enzyme and substrate combined.
  - Most are proteins.
  - They decrease the activation energy of a reaction.
  - They Speed up the reaction...
  - They are specific to the substrate.
- ❑ They allow the substrate to fit into the active site like a ball in to a glove, holding it tight. (**Lock and Key Model**) / They can be reused – **NEVER DESTROYED**.




# Enzymatic Activity



Rate vs Energy Graph - lowers activation energy

## Enzyme Action & Inhibition





# Nucleic Acids



## 01


### Characteristics

- **Nucleic acids** are composed of long chains of **nucleotides** (monomer) linked by dehydration synthesis.
- Composed of the Elements: C, H, O, N & P
- **Nucleotides** include:
  - ☐ A phosphate group
  - ☐ A pentose sugar (5-carbon)
  - ☐ A nitrogenous bases:
  - ☐ Adenine (A)
  - ☐ Thymine (T) - **DNA only**
  - ☐ Uracil (U) - **RNA only**
  - ☐ Cytosine (C)
  - ☐ Guanine (G)

## 02

### Functions

- **Functions**
  1. Transmits hereditary / genetic information
  2. Leads to the formation of proteins



## 03


### 2 Polymers

- **Two types:**
  1. **Deoxyribonucleic acid** (DNA-double helix)
  2. **Ribonucleic acid** (RNA-single strand)


# Nucleic Acid Comparison




## Deoxyribonucleic Acid



## Ribonucleic Acid



MAMMOTH SCIENCE



EST

2022

PUBLISHING



# Thank you!

Do you have any questions?

[instructor@email.com](mailto:instructor@email.com)

Xxx-xxx-xxxx x-xxxx

