

## <u>Immunizations Required for School</u> (Ages and Grades of Students Admitted)

Age/Grade	Shot	Dosage	Comments
3 and 4 years old	DTaP	4-5 doses	Entering school at age 3: fourth dose required at 18 months through 3 years of age. Must receive 5 <sup>th</sup> dose as soon as possible after turning 4. Entering school at age 4: fourth or fifth dose must be no earlier than four days prior to 4 <sup>th</sup> birthday.
	Polio	3-4 doses	Entering school at age 3: third dose required at 18 months through 3 years of age. Must receive 4th dose as soon as is possible after turning 4. Entering school at age 4: last shot must be no earlier than 4 days prior to the 4 <sup>th</sup> birthday.
	НІВ	1 dose	Must have received vaccine on or after turning 15 months old unless student has received a series of three and a booster as a baby.
	MMR	1-2 dose	Entering school at age 3: must have received initial dose no earlier than four days before the 1st birthday. Entering school at age 4: must have received 2 doses of measles vaccines.
	Varicella (Chicken Pox)	1 dose	(Chicken Pox): one varicella vaccine or a validated history of having had the disease. A parent's note that states the student had chicken pox on an approximate date (month and year) is sufficient. The vaccine must have been given on or after the 1st birthday.
	Hepatitis A	2 doses	Must have received 2 doses on or after 1 <sup>st</sup> birthday. The first dose is followed by a second dose at least 6 months apart.
	Hepatitis B	3 doses	Must have received series of 3 doses. The first dose is followed by a second dose at least one month later, and a third dose at least 2 months after the second. There must be at least 4 months between the first and third dose.
	PCV7	1 dose	<b>Pneumococcal Conjugate Vaccine (PCV7)</b> Must have 1 dose after 2 years of age if they do not already have a history of having had 4 does.
3 and 4 years old at high risk*	PPC23	1 dose	Pneumococcal Polysaccharide Vaccine (PPV23) *Children with certain chronic conditions or immuno-supression conditions will also require a dose of PPV23 in addition to PCV7 2 months after the last dose of PCV7
Kindergarten and older	All the above shots (minus HIB, Hepatitis A, PCV7 and PPV23) are required.		
7 years old	Dt	3 doses	If the DTaP was not completed at an earlier age, then 3 doses are required.
K – 12 <sup>th</sup> grade	DTaP/DTP	4-5 doses	See above under DTaP.
K – 12 <sup>th</sup> grade	Measles	2 doses total	The first dose must be on or after the first birthday and the second dose no less than 30 days later.
K – 12th grade	Hepatitis B	3 doses (5 mcg/0.5ml)	Must have received a series of 3 doses.
		2 dose (10 mcg/0.1ml)	In replacement of the entire 3-dose series, adolescents between the ages of 11 and 15 years may receive the 2-dose regime. There must be a six-month interval between doses.
K – 12th grade	Varicella	1 dose	Must have received 1 dose of varicella or a documented history of the disease. Students beginning the series after age 13 must receive 2 doses with a 30-day interval.
9th grade	TdaP/TD	10 yr. booster	Must be received prior to entering 9th grade or at 10-year anniversary of series or booster, whichever comes first. It is recommended that this vaccine be given at $11 - 12$ years of age.